

Supplementary Information

Transcriptome Analysis of the Developmental Effects of Bisphenol F Exposure in Chinese Medaka (*Oryzias sinensis*)

Zhiying Liang ^{1,†}, Yafen Guo ^{1,†}, Duan Pi ^{1,†}, Xiang Li ^{1,†}, Bingying Li ¹, Yongsu Huang ¹, Xiaohong Song ², Ramji Kumar Bhandari ³ and Xuegeng Wang ^{1,*}

¹ Institute of Modern Aquaculture Science and Engineering, Guangdong-Macao Joint Laboratory for Aquaculture Breeding Development and Innovation, College of Life Sciences, South China Normal University, Guangzhou 510631, China

² The Guangxi Key Laboratory of Theory and Technology for Environmental Pollution Control, Guilin University of Technology, Guilin 541000, China

³ Department of Biology, University of North Carolina Greensboro, Greensboro, NC 27412, USA

* Correspondence: wangxuegeng@scnu.edu.cn

† These authors contributed equally to this work.




Figure S1. Results of developmental exposure. (A): Survival rates; (B): Body length (mm); (C): Embryo heart rate (bpm).




Figure S2: Percentage of larvae with abnormalities. (A): Fry with pericardial edema; (B): Fry with enlarged yolk sac; (C): Fry with spinal curvature; (D): Fry with craniofacial abnormalities; (E): Fry with decreased head-trunk angle.




Figure S4: Staphylococcus aureus infection pathway. Green indicates down-regulation DEGs, red indicates up-regulation DEGs.




Figure S6: GnRH secretion pathway. Green indicates down-regulation DEGs; red indicates up-regulation DEGs.