

Table S1. The primers designed for real-time quantitative PCR.

Primer name	Primer sequence (5'-3')
qCpCAF1-F	CAAATCAGAGAGGTATGGAACGACAAT
qCpCAF1-R	GTAGTCGGACGTGGTCTTGAAGTTG
CpActin-F	GTTATGGTTGGGATGGGACAGAAAG
CpActin-R	GGGCTTCAGTAAGGAAACAGGA
CpTublin-F	TAGTGACAAGACAGTAGGTGGAGGT
CpTublin-R	GTAGGTTCCAGTCCTCACTTCATC
AtActin-F	GTTTCGTCTTCCACTTCAG
AtActin-R	ATCATACCAGTCTCAACAC
AtFT-F	TTCCAAGTCCTAGCAACCCTCACC
AtFT-R	TTCTTCCTCCGACGCACTCTC
AtFLC-F	CCTAATTTGATCCTCAGGTTTGGG
AtFLC-R	CCGACGAAGAAAAAGTAGATAGGCAC
AtSOC1-F	GCTCTCAGTGCTTTGTGATGC
AtSOC1-R	AAGAACGTACTTGGAGCTGGC
AtELF3-F	CTGGACCATCTAGTCAGCCTTG
AtELF3-R	TGCCATGAGACTGAGATCTTCTTG

Table S2. PCR Primers used for cloning *CpCAF1* promoter and deletion fragment.

Name	Primer sequence (5'-3')	Distance to TSS
CpCAF1pro-F	G CTCTAG AGTGTAAGTGAAGAATTAAGAATTG	−1908
CpCAF1pro-P2-F	G CTCTAG AGAAACTGATGTCAAATTGGGTTAAT	−1646
CpCAF1pro-P3-F	G CTCTAG ATAAGGTTCAATACTTCTGTGACATG	−941
CpCAF1pro-R	CATG CCATGG CTTCTCTCTTTTCCTTTTTTT	+224
CpCAF1pro-P1-R	CATG CCATGG CGGGAGGGCTAGGGAAAG	+2

Note: The restriction sites are indicated in bold, *Xba*I: TCTAGA , *Nco*I: CCATGG.