



The Fight Against the Carcinogenic Epstein-Barr Virus: Gut Microbiota, Natural Medicines, and Beyond

Radwa A. Eladwy ¹, Hang Thi Vu ², Ravi Shah ¹, Chun Guang Li ¹, Dennis Chang ¹ and Deep Jyoti Bhuyan ^{1,*}

- ¹ NICM Health Research Institute, Western Sydney University, Penrith, NSW 2751, Australia
- ² Faculty of Food Science and Technology, Vietnam National University of Agriculture, Trau Quy, Gia Lam, Hanoi 100000, Vietnam
- * Correspondence: d.bhuyan@westernsydney.edu.au; Tel.: +61-2-9685-4748

Abstract: Despite recent advances in oncology, cancer has remained an enormous global health burden, accounting for about 10 million deaths in 2020. A third of the cancer cases in developing counties are caused by microbial infections such as human papillomavirus (HPV), Epstein-Barr Virus (EBV), and hepatitis B and C viruses. EBV, a member of the human gamma herpesvirus family, is a double-stranded DNA virus and the primary cause of infectious mononucleosis. Most EBV infections cause no long-term complications. However, it was reported that EBV infection is responsible for around 200,000 malignancies worldwide every year. Currently, there are no vaccines or antiviral drugs for the prophylaxis or treatment of EBV infection. Recently, the gut microbiota has been investigated for its pivotal roles in pathogen protection and regulating metabolic, endocrine, and immune functions. Several studies have investigated the efficacy of antiviral agents, gut microbial metabolites, and natural products against EBV infection. In this review, we aim to summarise and analyse the reported molecular mechanistic and clinical studies on the activities of gut microbial metabolites and natural medicines against carcinogenic viruses, with a particular emphasis on EBV. Gut microbial metabolites such as short-chain fatty acids were reported to activate the EBV lytic cycle, while bacteriocins, produced by Enterococcus durans strains, have shown antiviral properties. Furthermore, several natural products and dietary bioactive compounds, such as curcumin, epigallocatechin gallate, resveratrol, moronic acid, and andrographolide, have shown antiviral activity against EBV. In this review, we proposed several exciting future directions for research on carcinogenic viruses.

Keywords: oncogenic viruses; Epstein-Barr virus; gut microbiota; natural products; short-chain fatty acids; gut metabolites

1. Introduction

Cancer has remained a serious concern and is the second leading cause of death worldwide [1]. The transformation of normal cells into neoplasm as a result of external stimuli, including physical carcinogens (such as UV), chemical carcinogens (such as asbestos), and biological carcinogens (such as viral and bacterial infections), can lead to various cancers [2]. The early link between cancer and viral infections was studied in chickens in 1910 by the Nobel Laureate Peyton Rous [3]. Since then, several viruses with their oncogenic mechanisms have been discovered. To date, eight oncogenic viruses (both RNA and DNA viruses) have been identified, which can induce cancer by different mechanisms [4].

Oncogenic viral infections contribute to about 15% to 20% of total cancer in humans [5]. These known human oncoviruses include the hepatitis B virus (HBV), hepatitis C virus (HCB), Epstein-Bar virus (EBV), human T-lymphotropic virus 1 (HTLV-1), Kaposi sarcoma-associated herpesvirus (KSHV; also known as human herpesvirus 8), human papillomaviruses (HPV), Merkel cell polyomavirus (MCPyV), and human immunodeficiency virus (HIV) [6]. Oncoviruses are infectious agents that cause prolonged, persistent

Citation: Eladwy, R.A.; Vu, H.T.; Shah, R.; Li, C.G.; Chang, D.; Bhuyan, D.J. The Fight Against the Carcinogenic Epstein-Barr Virus: Gut Microbiota, Natural Medicines and Beyond. *Int. J. Mol. Sci.* **2023**, *24*, 1716.

https://doi.org/10.3390/ijms24021716

Academic Editors: Kamalendra Singh and Christian Lorson

Received: 8 November 2022 Revised: 8 January 2023 Accepted: 11 January 2023 Published: 15 January 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/license s/by/4.0/). infections that can induce the incidence of tumours [7–9]. For instance, HIV, rather than causing cancer directly, induces carcinogenesis of other viruses, including EBV and KSHV [6]. According to the classic carcinogenesis theory, initiators and promoters act on cells in multiple steps or 'hits,' which sets the transformation process, invasion, and metastasis [10,11]. The accumulation of these multiple hits is required for full cancer formation [12] and this whole process may take several years [12]. 'Hits' are considered genetic and defined by amplifications in oncogenes, mutations & rearrangement, or changes in tumour suppressor genes [13].

The transformation of a normal cell into a cancer cell by virus-mediated carcinogenesis involves initiation, promotion, and progression (Figure 1) [14]. An interaction between a carcinogen and the host DNA is initiation, whereas promotion is when cell proliferation occurs (which usually takes a few months to years). The final step is the spread of the tumour, known as progression. Human oncoviruses display direct and indirect cell transformation mechanisms (viral carcinogenesis) (Figure 1) [15]. The virus encodes genes that activate growth and enhances apoptosis resistance for direct carcinogenesis, therefore, modifying the DNA repair mechanism [4,16]. During viral oncogenesis, tumour suppressors like p53 and pRb are inactivated until the DNA repair process is restored, otherwise, cell death is activated (Figure 1) [16]. Our genome also tends to acquire mutation, which is increased during viral infections. For example, viral antigen EBNA-1 can cause genome instability by activating *RAG1* and *RAG2* (Figure 1) [3].



Figure 1. Transformation of normal cells into cancer cells by virus-mediated carcinogenesis.

Oncogenic viruses maintain telomere length by interfering with telomerase expression leading to uncontrollable division [17]. Viral antigens like Kaposi's sarcoma-associated herpesvirus latency-associated nuclear antigen (LANA), HPV E6, HBV HBx, and latent membrane protein 1 of the Epstein-Barr virus have been found to enhance telomerase activity [18]. MicroRNAs have been found to inhibit the translation of mRNA [19]. These microRNAs target cellular tumour suppressor genes like *WIF1*, *Bin*, and *PUMA* and inhibit cell death [20]. For indirect carcinogenesis, viruses do not live in the cancer cells but act through mechanisms such as chronic inflammation, immune suppression, and oncomodulation, leading to carcinogenic mutations in the host cells [21–23].

Standard antivirals, including acyclovir and ganciclovir, can prevent EBV and its lytic replication but have not received Food and Drug Administration (FDA) approvals [24]. Additionally, the use of acyclovir and ganciclovir is linked to thrombocytopenia, liver and kidney damage, and gastrointestinal dysfunction [25]. Therefore, discovering novel, potent, and safe antiviral agents is crucial. An analysis of medicinal chemistry annual reports from 1984 to 1995 indicated that the majority of the synthetic agents approved by the FDA are derived from medicinal plants [26], however, critical and comprehensive

reviews on the activity of these natural bioactive molecules against oncogenic viruses are limited.

Gut microbiota is the collection of eukarya, archaea, and bacteria colonising the host's gastrointestinal tract (GIT). It coevolved over thousands of years through symbiosis with the host in a complex and mutually beneficial relationship. Studies on gut microbiota have just started in the last 20 years, despite its pivotal role in human health and disease pathogenesis. Our group has recently reviewed the fundamental role of gut microbiota and its metabolites in the development and treatment of cancer [27,28]. In the current review, we summarise and discuss the activity of gut microbial metabolites and natural products against carcinogenic viruses, emphasising EBV, and their implementation in anticancer drug development in the future.

2. Epstein–Barr Virus

EBV, formally called the human gamma herpesvirus 4, is one of the most common viruses in humans [29]. It infects more than 90% of the population worldwide [30]. Primary infection during childhood is generally asymptomatic. However, the virus can cause infectious mononucleosis in 35–50% of the cases when infection occurs later in life [24]. EBV infects 90–95% of all adults globally and causes approximately 1% of all cancers and is implicated in around 2% of all cancer-related deaths [31]. EBV is associated with several human malignancies, including Burkitt lymphoma, Hodgkin disease, nasopharyngeal carcinoma (NPC), gastric cancer, T/NK lymphoma (nasal natural killer/T-cell lymphoma), AIDS, and transplantation-associated lymphomas [30]. The virus has two cycle stages, the latent and lytic cycles [24]. In latency, the virus expresses only a limited number of genes, including EBNA 1-6, LMP1, and LMP2 [24]. In the lytic cycle, EBV expresses two transcription factors, Rta (R transactivator, the product of BZLF1 gene) and Zta (Z EB replication activator, the product of BZLF1). These two proteins trigger lytic genes, which encode diffused early antigen (EA-D) and DNA polymerase (Figure 2). Positive regulators of BZLF1 and BRLF1 genes can trigger the shift from the latency to the lytic cycles of EBV. Activators of *BZLF1* and *BRLF1* genes in B cells include physiological stimuli, B-cell receptor engagement, DNA damage, TGF-β, hypoxia, and chemical agents [32].

After DNA replication, late genes are expressed, followed by the production of virus capsid antigens and membrane protein, and then they produce the mature virion [24]. The lytic infection was shown to contribute to lymphoproliferative disease [24]. Therefore, the antiviral agent that blocks the viral lytic cycle, such as resveratrol, triptolide, and berberine, can provide potential treatment or prevention strategies for EBV-associated diseases (Figure 2).

EBV latency in resting and proliferating cells is divided into types III, II, I, and 0, depending on the viral gene expression pattern [33]. It was reported that EBV induces cancers of epithelial and lymphoid origin by the transformation of quiescent B cells into lymphoblastoid cell lines (LCLs) (Figure 2) [34]. The virus life cycle is confined chiefly to the B cells, establishing premalignant latent gene expression patterns. When EBV infects a naïve B cell, it expresses the latency III or growth program [35]. In this pattern, the virus expresses EBNA 1-6, LMP1, LMP2A, and LMP2B with a complete oncogenic component of nuclear proteins EBNA2, 3A, 3B, 3C, LP, and the EBV microRNAs. The infected B cell then becomes proliferating B-cell which helps in EBV episomes replication. Notch signalling is dysregulated in this gene expression pattern, leading to nonviral lymphoid malignancies. Expression of *c-myc*, a cellular proto-oncogene, is dysregulated by upregulating cyclin Ds and E and downregulating CDK2 inhibitors like p21^{CIP1} and p27^{KIP1}. Latency III has a free pattern of gene expression resulting in the production of oncogenic proteins and eventually can cause cancer, especially in immunocompromised patients [36]. In immunosuppressed and AIDS patients, AIDS-related-non-Hodgkin lymphoma and post-transplantation lymphoproliferative disease latency III expressions are common [37].



Figure 2. The impact of resveratrol, triptolide, and berberine on different phases of the EBV life cycle (modified from Münz, 2019 [33]). The virus passes from saliva to the B lymphocytes crossing the oropharyngeal epithelium. Here, the virus causes the B cell to multiply and outspread through the B cell compartment. The infected B cells may differentiate into latency 0 or total latency III transformation. The latency III phase is highly immunogenic and *EBNA1*, *EBNA2*, *EBNA-LP*, *EBNA3* (*A*, *B*, *C*), and *LMP* (1,2) are expressed. To remain immune silent, the virus switches from latency III to latency II in which *EBNA1*, *LMP1*, and *LMP2* are expressed, and then to latency 0. For the persistence of viral infection, EBV drives latency 0 to latency I, where *EBNA1* is only expressed to allow the EBV episome to be dissociated and maintained in dividing B cells. These B cells will randomly go for the lytic cycle and lyse to produce and release the virions back to the saliva to infect more host B lymphocytes. The germinal centre differentiation pathways provide premalignant precursors for Hodgkin's lymphoma and non-Hodgkin's lymphomas, including diffuse large B-cell lymphoma and Burkit's lymphoma. The lytic cycle primarily occurs in the plasma cells.

Latency III is the most immunogenic phase of the EBV life cycle. Therefore, the cells express a pattern shift to a default program or latency II to remain partially immune silent (Figure 2). *EBNA1* and two oncogenes, *LMP1* and *LMP2A*, are expressed in latency III [35]. These two oncogenes are essential for survival and B cell proliferation [38]. *LMP1* activates NF-_kB, a key transcriptional factor in viral and nonviral lymphomagenesis which upregulates antiapoptotic molecules like A20 and Bcl2 in B cells to promote cell proliferation and malignancy [35]. *LMP1* inhibits DNA damage response and enhances telomerase activity via JNK activation to promote genetic instability [39]. Likewise, *LMP2A* helps in B cell differentiation, survival, and cell growth when the Ig receptor expression is unavailable through PI3K-AKT pathway activation [40]. AIDS-associated lymphomas and post-transplant lymphoproliferative disorders are common during latency III, while diseases like Hodgkin's disease and nasopharyngeal carcinoma are associated, during latency II [36].

Burkitt's lymphoma is common in latency I. The infected cell can switch to immunologically silent latency programs. In this expression pattern, *EBNA1* is only expressed to allow the EBV episome to be disassociated and maintained in dividing B cells [35,41] (Figure 2). EBV replicates either by B cell proliferation or by lytic replication. In a cell culture setting, some chemicals like sodium butyrate and 12-*O*-tertradecanoylphorbol (TPA) can induce the EBV lytic cycle [42]. It suggests that host cell signal transduction and epigenetic regulation play an essential role in switching the virus between latency and lytic replication. When lytic replication occurs, the virus expresses almost all the genes present in its genome [33]. For instance, genes like *BALF1*, *BCRF1*, and *BARF1* are expressed during the lytic cycle, with oncogenic consequences [36]. The early lytic EBV replication creates a tumour microenvironment responsible for the observed increased tumorigenesis [33]. Also, more TNF, CCL5, and IL-10 are produced by LCLs with an increased level of spontaneous lytic EBV reactivation, which recruits immunosuppressive myeloid cells (such as TNF α -CCL5-macrophages) and inhibits the immune control by cytotoxic lymphocytes. From a preclinical in vivo setting, EBV-associated lymphomagenesis increases with lytic EBV replication through the formation of a tumour microenvironment, and suppression of this lytic replication might be a useful strategy for the treatment of EBV-induced malignancies.

Several antiviral compounds designed for other herpesviruses have been tested on EBV. These antiviral compounds can be divided into two major categories, (a) those that interfere with the virus-encoded enzymes, and (b) the ones that interfere with the cellular processes required for virus production [24]. The former includes nucleoside analogues and non-nucleoside analogues. The other category comprises drugs targeting cellular protein. Although several antivirals were proven to inhibit EBV replication in vitro, they had limited success in clinical trials [24]. The current literature reveals that no antiviral drug has been approved to treat EBV infections [24,25,43].

3. Gut Microbiota

Microbiota refers to a vast number of interacting bacteria, fungi, eukaryotic viruses, archaea, and bacteriophages coexisting with the host for potential mutual benefit [44]. In the last two decades, extensive research has been done to explore the nature and therapeutic potential of gut microbiota [45]. Emerging evidence has elucidated the role of gut microbiota in protecting from pathogens, maintaining metabolic, endocrine, and immune functions, and modifying drug action and metabolism (Figure 3) [45]. Gut microbial metabolites include short-chain fatty acids, inosine, and bacteriocins, which have shown a broad spectrum of biological activities in previous studies including anticancer and immunomodulatory functions [27,46,47].



Figure 3. Epstein-Barr Virus (EBV) medicated carcinogenesis and the potential protecting role of gut microbiota through the production of metabolites.

3.1. Probiotics and EBV

Probiotics are living organisms that provide beneficial effects to the host when consumed in sufficient amounts. *Lactobacilli* and *Bifidobacteria* are the most common probiotics [48]. Probiotics have been used as antimicrobials and for the prevention and management of different allergies [48]. For instance, supplementation of *Lactobacilli casei* (strain Shirotais) (6.5 × 10⁹ CFU per day) to endurance athletes for 20 weeks reduced the plasma levels of EBV and CMV (cytomegalovirus) antibodies, presumably by improving the host immunity [49].

3.2. Short-Chain Fatty Acids (SCFAs)

SCFAs are formed principally by fermentation of undigested starch and nonstarch polysaccharides in the large bowel [50]. These are small monocarboxylic acids comprised mainly of acetic (or acetate), propionic (or propionate), and butyric (or butyrate) acids (Figure 4) [50].



Figure 4. Chemical structure of short-chain fatty acids.

SCFAs are metabolised by the colonocytes and the unmetabolized fractions are transported into the portal circulation to be used as an energy source for the hepatocytes, except for acetate which is barely oxidized in the liver [51]. As a result, only a small amount of SCFAs, originating from the colon, reaches the systemic circulation, so their faecal concentration has been used as a proxy for SCFAs production in the colon [45]. Many studies have shown the beneficial role of SCFAs in different diseases, including diabetes, cancer, and hypertension. Table 1 summarises the potential therapeutic use of SCFAs and their proposed mechanisms of action. Several reports demonstrated the protective role of SCFAs against infections primarily attributed either to their direct antivirulence effect or indirect effect on the host immune system [52]. Recently, SCFAs showed antiviral effects against respiratory syncytial virus bronchiolitis and, interestingly, it is also reported that SCAFs can activate the EBV lytic cycle [52,53]. In particular, propionate and butyrate have been found to activate the EBV lytic cycle. They have also been reported as general inhibitors of class I and class II histone deacetylases in different cell types [53].

SCFA	Viral Infection	Type of Study	Description and Outcome of the Study	References	
Acetate	 Respiratory syncytial vi rus (RSV)	In vitro	Pretreatment of the A549 cells with acetate resulted in a reduction of RSV in- fection and increased expressions of retinoic acid-inducible gene-I (<i>RIG-I</i>) and interferon-stimulated gene15 (<i>ISG15</i>).		
		In vivo	The intranasal treatment of acetate in mice reduced RSV infection and viral load and increased expression of <i>RIG-I</i> and Interferon β 1 (IFNB1) in the lung tissue.	viral e lung	
		Ex vivo	Ex vivo treatment with acetate in RSV-positive infants showed reduced viral load and increased expression of <i>ISG15</i> , <i>RIG-I</i> , and MAVS (mitochondrial antiviral-signalling protein).	[54]	
		Clinical	Stools from 17 RSV-positive infants showed that the increased levels of acetate were associated with faster recovery, a shorter duration of fever, and higher oxygen saturation at admission.		
Propionate	Herpes simplex virus 1 (HSV-1)	In vivo	Mice were given sodium propionate supplement in drinking water three weeks before induction of the corneal HSV-1 infection, the animals had fewer ocular lesions when compared to the control group. Lymphoid and corneal tissues showed lower levels of CD4 (T-helper) Th1 and Th17 cells, macrophages, and neutrophils and higher levels of regulatory T cells (Treg) when compared to the control.	[55]	
Butyrate	SARS-CoV-2 (COVID-19)	Clinical	High throughput RNA sequencing of many genes related to SARS-CoV-2 in- fection (4526 upregulated genes and 3167 downregulated genes) in butyrate- treated organoids and showed that butyrate downregulates genes essential for SARS-CoV-2 infection, such as <i>ACE2</i> and its associated genes. Butyrate was also found to upregulate toll-like receptors (TLR) and other antiviral path- ways	[56]	

3.3. Inosine

Inosine, the key intermediate in purine metabolism, is produced via the deamination of adenosine by specific RNA deaminases (Figure 5) [57]. In living organisms, inosine is an abundant nucleic acid base, and it serves as a crucial intermediate in purine metabolism [57]. In a recent in vivo study, the enhanced immunotherapy response by intestinal *Bifidobacterium pseudolongum* was modulated mainly by inosine metabolite and it was dependent on T cell expression of the adenosine A2A receptor [58].

Isoprinosine (IP), known as inosine pranobex or methisoprinol, is a 1:3 complex of inosine and pranobex, respectively (Figure 5) [46]. The latter is a mixture of dimethyl amino isopropanol (dimepranol) and p-acetamidobenzoate (acedoben) (Figure 5) [46]. Since 1971, IP has been widely used against viral infections, including the herpes simplex virus (HSV), HPV, HIV, cytomegalovirus (CMV), influenza, and acute respiratory infections, as well as EBV infections, primarily due to its immunomodulatory effect and safety profile. The antiviral properties of IP, when given after the onset of virus diseases, is due to enhancing the natural immune response of lymphocytes [46].



Inosine pranobex Complex

Figure 5. Formation of inosine by specific RNA deaminase; ADA: adenosine deaminases.

Table 2 summarises the key in vivo studies of isoprinosine against different viral infections. The mechanisms of action of IP as an immunomodulator in viral infections have been proposed to be through increasing the levels of proinflammatory cytokines, such as IL-2 and INF-c in mitogen- or antigen-activated cells, which initiates T-lymphocyte differentiation resulting in induced lymphoproliferative action [59,60]. INF-c, in combination with the direct effect of IP, decreases the production of anti-inflammatory cytokines, including IL-10 suggesting the potential immunomodulatory effect of IP on innate and adaptive immunity. IP was also reported to increase the natural killer (NK) cells population and activity and potentiate phagocytosis and macrophage chemotaxis [61–63].

The in vivo antiviral activity of IP was evaluated on murine gammaherpesvirus 68 (MHV68), which is a natural pathogen of mice, and used as a model for EBV infection. After 2 weeks of treatment, IP resulted in elevated levels of virus-neutralizing antibodies, leukocytes, and neutrophils. However, the antiviral effect was time-limited and disappeared after 120–150 days. The incidence of tumour formation was 7.5% in the MHV-infected group after treatment with IP, compared to 17.5% in the untreated group. This study showed that IP administration should be repeated in chronic EBV infections [64].

Table 2. Summary of key studies investigating the enfcacy of isoprinosine (ir) in different viral infections.				
Infection	Type of Study	Main Findings	References	
Epstein-Barr virus	In vitro	The protective immunomodulating effect of IP was investigated on EBV- transformed B lymphoid cells. IP was found to be an immunomodulating agent with an ability to enhance the response of sensitized peripheral blood mononuclear cells (PBL) to EBV-antigens.	[65]	
Acute respiratory viral Clinical: Randomised, double-blind, Phase 4		Faster improvement in subjects in the IP group versus those in the placebo	[66]	
infections	study (n = 231) with placebo (n = 232)	group.		
SARS-CoV-2 (COVID-19)	Clinical: The study was conducted during June-September 2020 in three nursing homes (in the Czech Republic) with 301 residents, 156 of whom (51.8%) tested positive for the SARS-CoV-2 virus (PCR test).	Demonstrated the positive effect of IP against COVID-19. The case-fatality rate in these three nursing homes at the end of the pandemic's first wave was lower than similar nursing homes in the Czech Republic.	[67]	

Table 2. Summary of key studies investigating the efficacy of isoprinosine (IP) in different viral infections.

3.4. Bacteriocins

Bacteriocins are ribosomally-synthesized small amphiphilic peptides produced by archaea and bacteria and extracellularly transported via an ATPbinding cassette transporter [47]. Several bacteriocins produced by the lactic acid bacteria (LAB) display bactericidal or bacteriostatic effects on similar or closely related bacterial strains [68]. They are fundamentally classified into two classes, class I lantibiotics and class II nonlantibiotics [47]. Lantibiotics, are gene-encoded peptides that are nontoxic to humans and have versatile applications as antibiotics, biopreservatives, or bactericidal agents in cosmetics [47]. They act by forming pores in the cell membrane resulting in a decrease in the intracellular pH and efflux of small metabolites [47]. Class I (lantibiotics) bacteriocins include the single peptides nisin, mersacidin, and lacticin, while class II (non-lantibiotics) include pediocin, lactacin, and lactococcin. Although a precise mechanism of action for every bacteriocin is still unclear, binding to lipid II, the main transporter of peptidoglycans from the cytoplasm to the cell wall, is reported in different studies [47]. Several reports documented the antibacterial properties of bacteriocin. However, very few reports to date have displayed their potential antiviral effect [69]. In particular, previous studies demonstrated the antiviral properties of bacteriocins against different viruses including the oncogenic viruses HSV-1 and HSV-2 [70-73].

4. Natural Products as Antiviral Agents against EBV

Several oncogenic viruses have been identified since the association between viruses and cancer was discovered in 1910 [3]. Even though the vaccines against HBV and HPV have been highly successful, more preventative and therapeutic options are necessary for other oncogenic viruses. Recent research has focused on the recovery of antivirals from medicinal plants and foods, such as curcumin, andrographolide, resveratrol, epigallocatechin gallate (EGCG), and moronic acid (Table 3) [74–76]. However, more research is necessary to discover novel, potent, and safe natural antiviral agents against oncoviruses. Bioactive molecules from various natural sources, such as bryostatins, didemnin, cephalostatin from marine organisms, silymarin, stilbenes, vincristine from plants, rhizoxin, mitomycin, rapamycin from microorganisms, curcumin, allicin, and β -carotene from fruits, vegetables, and herbs have been analysed in vitro for their potential activity against oncogenic viruses [4,77–79]. However, their molecular mechanisms of action have not been investigated adequately, and further studies are crucial to understand their efficacy in living organisms through in vivo studies. Few natural medicine products including berberine, manassantin B, grifolin, and triptolide have been tested both in vivo and in vitro in relation to the EBV antiviral and anticancer mechanisms) [80]. The summary of plants and natural bioactive compounds with antiviral effects against EBV is presented in Table 3. The following section highlights the most interesting findings.

Andrographis paniculata is a medicinal plant commonly used in traditional medicine in Asia. The herb has a bitter taste and is used to treat the common cold, colic pain, liver disorders, and bowel issues [75]. *A. paniculata* was reported with several biological properties such as immunoregulatory, anti-HIV, and antibacterial activities in previous studies [75]. Recent research showed the antiviral effect of the ethanolic extract of *A. paniculata* against EBV [75]. The mechanism of inhibition was reported to occur via blocking the transcription of the immediate–early genes that encode lytic proteins Rta and Zta. *Andrographolide* was the compound of interest in *A. paniculata* which effectively inhibited EBV at 5 µg/mL [75]. Furthermore, the compound was found to be non-toxic to the P3HR1 cells at that concentration, indicating its potential anti-EBV effect.

Polygonum cuspidatum is another popular medicinal plant in Asia, which has been used as a hepatoprotective and cholagogic traditional medicine in China, Japan, and Korea [81]. The ethanolic extract of *P. cuspidatum* was reported to inhibit the transcription of EBV immediate early genes, and the expression of lytic proteins Rta, Zta, and EA-D [82,83]. The extract also inhibited the expression of latent membrane protein one triggering the EBV-positive cells toward apoptosis [82]. In addition, the ethyl acetate subfraction from ethanolic extract displayed a more potent EBV inhibitory effect [84]. The subfraction blocked the transcription of immediate early genes, reduced DNA replication, and hindered the production of virus particles [84]. Furthermore, the primary active components in *P. cuspidatum* were reported to be resveratrol and emodin. The effective concentration of emodin required to inhibit the expression of immediate-early protein by 50% (EC50) obtained from flow cytometry was 4.83 µg/mL (17.87 µM) and its EC50 value to reduce DNA replication was 1.2 µg/mL [84]. These values were substantially lower than those of resveratrol, EGCG, and andrographolide, suggesting that emodin effectively inhibited the EBV lytic cycle and can be a potential lead for drug development for EBV-related diseases [84].

Diterpenoids are common secondary metabolites found in plants such as *Scutellaria barbata* and *Euphorbia milii*. *E. milii* has been used in Chinese medicine for detoxifying purposes for centuries [85]. This group of compounds exhibits a wide range of biological activities including antimicrobial, anti-inflammatory, antitumour, and antiviral effects. Neo-clerodane diterpenoids from *S. barbata* showed inhibition toward the EBV lytic cycle [86]. Similarly, diterpenoids in *E. milii* were reported to have potential antiviral activity. In particular, the acetone extract of *E. milii* inhibited the EBV lytic cycle [87]. Thirteen new entrosane-type diterpenoids (1–13) were isolated from the *E. milii* extract and evaluated against EBV. Among those, one derivative showed the most potent inhibitory activity with an EC₅₀ value of 5.4 μ M compared to the positive control (+)-rutamarin (EC₅₀ = 5.4 μ M) [87].

Berberine is a natural compound found in the rhizome, roots, and bark of medicinal plants such as barberry (Berberis vulgaris) and huanglian (Coptidis rhizome) (Figure 6) [88]. Several studies showed the potential anti-inflammatory and chemo-preventative effects of berberine [88]. It has been illustrated that berberine can inhibit cell proliferation, cell cycle arrest, and apoptosis of EBV-associated NPC cells [89]. Berberine targeted EBNA1 transcription and downregulated its expression, inhibited the p-STAT3 expression, and decreased EBV virions production [90]. The in vivo results of a nontoxic dose of berberine showed a decrease in tumour growth of the EBV-associated NPC [91]. In addition, berberine-induced apoptosis illustrated the activation of XAF1 and GADD45a (Figure 6) [89]. Berberine increased the cellular reactive oxygen species levels and upregulated p53 via activation of JNK and p38-MAPK pathways. Then p53 translocated GADD45 α (growth arrest and DNA damage-inducible alpha) protein into the nucleus and the XAF1 (X-linked inhibitor of apoptosis 1) protein into the cytosol. Furthermore, p53 upregulated PUMA, a proapoptotic protein that rapidly induced apoptosis through a baxand mitochondrial-dependent pathway (Figure 6) [89].

Triptolide produced by the thunder god vine (*Tripterygium wilfordii*) was reported to have anti-inflammatory, immunosuppressive, and anticancer activities [92]. Studies have demonstrated that cell proliferation of EBV-positive B lymphoma cells can be inhibited by triptolide via downregulation of *LMP1* transcriptional expression [93]. Triptolide targeted the *LMP1* promoter in type III infection cells, downregulating the *LMP1* mRNA [93]. In comparison, other studies showed that triptolide inhibits the *EBNA1* expression by increasing the sensitivity of mitochondrial apoptosis in NPC [94].



Figure 6. A diagrammatic representation of the intracellular signalling mechanism during berberine-induced apoptosis in EBV-positive B lymphoma cells (modified from Park et al. [89]). In this process, intracellular reactive oxygen species (ROS) production is stimulated upon berberine treatment, leading to the activation of various molecules, ultimately resulting in caspase-dependant apoptosis of EBV-positive B lymphoma cells [89]. Triptolide was reported to induce apoptosis via caspase-9 dependant cell death mediated through the mitochondrial pathway [95].

Compound	Source	Cell Line/Animal	Mode of Action	References	
Berberine	Berberis vulgaris	HK1 and HONE1 nasopharyngeal carcinoma Suppressed the expression of <i>EBNA1</i> . cells and NOD/SCID mice			
		C666-1, HONE1, HK1 and NP460 nasopha- ryngeal carcinoma cells and nude mice	STAT3 activation was inhibited in the nasopharyngeal car- cinoma cells.	[90]	
		CNE2 nasopharyngeal carcinoma cells and BALB/C-NU male mice	Regulated the expression of key proteins of the MAPK/ERK pathway.	[91]	
		IM-9 multiple myeloma cells	Upregulated XAF1 and GADD45a expression by MAPK and functional p53.	[89]	
Resveratrol	Polygonum cuspidatum	Raji Burkitt lymphoma cells and female mice	Activation of TPA-induced EBV-early antigen was suppressed.	[76]	
		P3HR-1 Burkitt lymphoma cells	Decreased the growth and development of the virus.	[96]	
		Akata and Raji Burkitt lymphoma cells	Decreased protein synthesis and lowered reactive oxygen species (ROS) levels. Redox-sensitive transcription factors AP-1 and NF-kB were suppressed.	[97]	
		Akata and B95-8 Burkitt lymphoma cells	Antiapoptotic proteins Mcl-1 and survivin were downreg- ulated.	[98]	
Apigenin	Punica granatum	NA and HA nasopharyngeal carcinoma and P3HR1 Burkitt lymphoma cells	Suppressed the promoter activities of two viral <i>IE</i> genes and decreased the viral reactivation	[99]	
(+)-Rutamarin	Rutu graveolens	P3HR-1 Burkitt lymphoma cells	Exhibited anti-EBV lytic DNA replication.	[100]	
Wogonin	Scutellaria baicalensis	Raji Burkitt lymphoma cells and four-week- old male BALB/c nude mice.	The expression of NF- <i>k</i> B was downregulated through LMP1/miR155/NF-kB/PU.1 pathway.	[101]	
Epigallocatechin gallate	Camellia sinensis	P3HR-1 Burkitt lymphoma cells	Inhibited the EBV lytic proteins expression.	[102]	
		CNE1-LMP1 nasopharyngeal carcinoma and B95-8 Burkitt lymphoma cells	The MEK/ERK1/2 and PI3K/AKT signalling were suppressed.	[103]	
			Downregulated LMP1.	[104]	
Glycyrrhizin	Glycyrrhiza glabra	HEK (Human embryonic kidney) cells	Inhibited EBV infection. Targeted the first step of the SUMO (small ubiquitin-like modifier)ylation process resulting in limited cell growth and apoptosis.	[105,106]	

Table 3. In vitro and in vivo biological activity of some natural products against Epstein-Barr Virus.

Triptolide	Tripterygium wilfordii	HONE1/Akata, C666-1 nasopharyngeal carci-	Reduced the expression of <i>LMP1</i> in EBV-positive B lym-	[93]
		cells	phocytes.	
		HEK and B95-8 Burkitt lymphoma cells, HeLa	Increased the mitochondrial apoptosis sensitivity in naso-	[94]
		(cervical cancer cells and BALB/c male mice	pharyngeal carcinoma cells.	
		HEK, B95-8 and P3HR-1 Burkitt lymphoma	Targeted and downregulated the translation factors SP1	[107]
		cells	and c-Myc.	
Phytol	Lindernia crustacea	P3HR-1 Burkitt lymphoma cells	RTA expression was inhibited.	[108]
Aloe-emodin	Lindernia crustacea	_	EBV lytic cycle was inhibited.	[108]
Cis/trans-martynoside &	Lindernia crustacea		EBV lytic cycle was affected.	[108]
Emodin	Polygonum cusnidatum	P3HR 1 Burkitt lymphoma colls	FBV lytic cycle was affected	[8/]
Linount	1 orggonum cuspitutum	1 of net Durkit lymphonia cens	Inhibited the transcription of FBV immediate early genes	[04]
			the expression of EBV lytic proteins and reduced EBV	
			DNA replication.	
		NA, HA, HONE-1 and TW01 nasopharyngeal	IA, HONE-1 and TW01 nasopharyngeal EBV reactivation and nasopharyngeal carcinoma recur-	
		carcinoma cells	rence were decreased.	
(+)-Hyperjaponicol B & Hyperjaponicol D	Hypericum japonicum	B95-8 Burkitt lymphoma cells	Suppressed EBV DNA replication.	[110]
Hyperjaponicol H	Hypericum japonicum	B95-8 Burkitt lymphoma cells	Moderately inhibited EBV lytic DNA replication.	[111]
Grifolin	Albatrellus confluens & Bole	-CNE1-LMP1 nasopharyngeal carcinoma cells	Targeted DNMT1 to block aerobic glycolysis.	[112]
	tus pseudocalopus		Blocked DNMT1 localization to restore OXPHOS	
Quercetin	Glycyrrhizia uralensis or G	SNU719 gastric carcinoma cells	Induced cell cycle arrest and strong early apoptosis and ne-	[113]
	glabra		crosis/late apoptosis. Inhibited infection of EBV from lym-	
			phocytes to gastric adenocarcinoma cells.	
Manassantin B	Saururus chinensis	P3HR-1 Burkitt lymphoma cells	Blocked lytic replication and virion production by inhibit-	[114]
			ing mTORC2 activity and blocking the mTORC2-	
			PKC/AKT-signalling pathway.	
Protoapigenone	Thelypteris torresiana	P3HR-1 Burkitt lymphoma cells	ZTA transactivation important for lytic cycle activation	[115]
			was inhibited.	
Curcumin	Curcuma longa	HONE1 and HK1-EBV nasopharyngeal carci-	Induced cell cycle arrest and apoptosis via the mitochon-	[74]
		noma cells	dria- and death receptor-mediated pathways.	

		HeLa cervical cancer cells	Inhibited the transcription level of EBNA1.	[74]
Andrographolide	Andrographis paniculata	P3HR1 Burkitt lymphoma cells	Inhibited the transcription of <i>IE</i> genes that encoded Rta and Zta	[75]
Thirteen <i>ent</i> -rosane-type <i>Euphorbia milii</i> diterpenoids		P3HR-1 Burkitt lymphoma cells	Inhibited lytic replication.	[87]
Twenty-six <i>neo</i> -clerodan diterpenoids	eScutellaria barbata	P3HR-1 Burkitt lymphoma cells	Inhibited lytic replication.	[86]
Moronic acid	Rhus chinensis	HEK, P3HR-1 Burkitt lymphoma cells	Inhibited the expression of Rta, and Zta, and interfered with the function of Rta.	[116]
28 lignans	Saururus chinensis	P3HR-1 Burkitt lymphoma cells	Inhibited lytic replication.	[117]
Polysaccharide	Astragalus membranaceus	Raji Burkitt lymphoma cells	Suppressed the expression of the IE protein, Rta, Zta, and EA-D.	[118]
Sulphated polysaccharides	Microalgae Ankistrodesmus convolutus Synechococcus elongatu and Spirulina platensis	Akata, B95-8, and P3HR- s, Burkitt lymphoma cells s,	Reduced cell-free EBV DNA.	[119]
Angelicin	Psoralea corylifolia	BC-3 and BCBL-1 lymphoma, and B95-8 and Raji Burkitt lymphoma cells	Inhibited lytic replication.	[120]
Lawsone (2-hydroxy-1,4-naphthoqui- none)	<i>Lawsonia inermis</i> (henr leaf)	aIn vivo two-stage mouse skin model	Reduced EBV-early antigen activation Reduced skin carcinogenesis in a mouse model.	[121]
Luteolin (3,4,5,7-tetrahy- droxyflavone)	Fruits and vegetables	NA, HA nasopharyngeal carcinoma, and P3HR-1 Burkitt lymphoma cells	Suppressed the activities of Zta and Rta by deregulating its Sp1 binding.	[122]

16 of 24

Saururus chinensis has been used as an anti-inflammatory, antipyretic agent to treat jaundice and edema in traditional Chinese and Korean medicine. Lignans, isolated from *S. chinensis* and *Litsea verticillate* exhibited an antiviral effect against EBV through inhibition of the lytic cycle along with other biological activity [117,123]. In particular, among 28 lignans isolated from *S. chinensis*, manassantin B showed the most promising inhibition [117].

The sulphated polysaccharides found in microalgae were also reported to have antiviral activity. For instance, the methanol extracts of *Synechococcus elongatus* and *Ankistrodesmus convolutus* were reported with low cytotoxicity and a strong antiviral effect against EBV in Burkitt's lymphoma cells. The antiviral activity was measured by reducing the cell-free EBV DNA [119].

Moronic acid found in *Rhus chinensis* and Brazilian propolis inhibited the expression of Rta, Zta, and an EBV early protein [116]. It was also reported that moronic acid interfered with the function of Rta. In particular, it reduced the ability of Rta to activate a promoter containing a Rta-response element [116]. As the expression of many EBV lytic genes depends on Rta, the treatment of P3HR1 Burkitt's lymphoma cells with moronic acid substantially reduced EBV particles produced by the lytic cycle [116].

Astragalus membranaceus has been used in traditional Chinese medicine for centuries primarily due to its immune-stimulatory, antiviral, antioxidant, and antitumor properties. The plant extract inhibited EBV lytic cycle by suppressing the expression of the immediate–early protein, including Zta, Rta, and EA-D [118]. The authors reported that the antiviral activities of *A. membranaceus* were attributed to its polysaccharides [118].

Henna (*Lawsonia inermis* L.) has been used to treat various ailments, including having antimalarial, antibacterial, and antidiabetic uses for centuries [121]. This plant is found in Africa, southern Asia, and part of Australia. Henna leaf powder and its primary pigment, lawsone (2-hydroxy-1,4-naphthoquinone), both showed significant inhibition (>88%) of EBV-early antigen activation in vitro. Additionally, henna leaf powder and lawsone significantly reduced skin carcinogenesis incidence in a mouse model administered by either oral or topical routes [121].

Cordyceps belong to the genus of Ascomycete fungi. Cordyceps are well known for their medicinal properties [124]. For instance, cordycepin, an adenosine derivative, found in cordyceps, expressed antitumor, antiviral and antifungal properties in previous studies [124]. Since cordycepin has a similar chemical structure to adenosine, it can be intercalated into RNA molecules and can terminate RNA synthesis [124]. Cordycepin can be either isolated from cordyceps or be produced synthetically and was reported to significantly downregulate most EBV genes of those tested [124]. The EBV genome copy number was reduced by up to 55%, in response to 125 μ M cordycepin treatment. Furthermore, cordycepin significantly suppressed EBV transmission from cell to cell in a coculture [124]. Therefore, cordyceps should be further investigated for their activity against gamma herpesvirus infection and resulting cancers.

Manassantin B, a natural lignan extracted from the roots of the Asian lizard's tail plant (*Saururus chinensis*), was found to be efficient in blocking EBV lytic replication and virion production with lower cytotoxicity [114]. The underlying mechanism by which manassantin B targets the viral lytic replication was through suppression of the *BZLF1* gene expression by interrupting the AP-1 signal transduction [114]. It specifically blocked the rapamycin complex 2 (mTORC2)-mediated phosphorylation of AKT Ser/Thr protein kinase at Ser-473 and protein kinase $C\alpha$ (PKC α) at Ser-657. Briefly, Manassantin B inhibited mTORC2 activity by blocking the mTORC2-PKC/AKT-signalling pathway [114].

DNA methylation is an integral part of the EBV lytic cycle in which the viral genome is methylated during the latency period, and it gradually becomes unmethylated in the lytic cycles [125]. In EBV-associated NPC patients, abnormal DNA methylation has been found, which correlates with poor survival outcomes. Targeting this DNMT1, which causes abnormal DNA hypermethylation, is vital for epigenetic cancer therapy [125]. Natural compounds such as Grifolin have demonstrated DNMT1 expression inhibition and mitochondrial translocation [112].

4.1. Antiviral Agents from Food Sources

Recent research has also found effective EBV antiviral agents such as resveratrol and EGCG from regular foods such as vegetables and spices. Resveratrol is a natural phenolic compound found in many plants and fruits. The antiviral activity of resveratrol was reported for the influenza A virus, HIV, polyomavirus, and Herpesviridae family [126]. De Leo et al. demonstrated that resveratrol strongly induced apoptosis of EBV-positive Burkitt's lymphoma cells depending on the viral latency program [97]. In particular, resveratrol greatly suppressed latency I EBV (+) cells, but not latency II and III cells (Figure 2). Additionally, resveratrol inhibited EBV reactivation by suppressing the lytic genes expression [96,97]. The production of virion was also reduced in a dose-dependent manner with the resveratrol treatment.

The popular green tea catechin EGCG is also a potential candidate for antiviral drug development. EGCG inhibited the expression of EBV lytic protein by blocking the transcription of immediate-early genes [102]. Liu et al. reported that EGCG could effectively inhibit the EBV lytic cycle at DNA, transcription, and protein levels [103,104]. Similarly, the dietary flavonoid apigenin, abundant in parsley, chamomile, celery, and vine-spinach, blocked the EBV lytic cycle by suppressing the expression of immediate–early lytic protein Zta, Rta, EA-D, and DNase and significantly reduced the production of EBV virion [99].

In comparison to EGCG and apigenin, luteolin (3,4,5,7-tetrahydroxyflavone) and protoapigenone inhibited EBV by suppressing the activity of immediate–early lytic protein [115,122]. Protoapigenone, a natural derivative of apigenin, inhibited the transactivation function of Zta and had no effect on the ability of Rta [115]. Additionally, protoapigenone was not toxic to the P3HR1 cells at the concentration that inhibit the function of Zta, indicating its therapeutic potential (selectivity index) in preventing EBV lytic proliferation [115]. Luteolin, a popular flavonoid has been reported to have several therapeutic properties including anti-inflammatory, antioxidant, antidiabetic, and antiangiogenesis activities. Luteolin also exhibited a significant inhibitory effect on EBV reactivation by suppressing Zta and Rta activities via deregulating its Sp1 binding, resulting in reduced virion production [122].

Angelicin is an angular furocoumarin naturally found in the seeds of *Psoralea corylifolia*, the roots of *Angelica archangelica*, and the family of Umbelliferae plants. Angelicin has been used to treat various skin diseases together with long-wavelength UV irradiation [120]. It efficiently inhibited lytic replication of murine gammaherpesvirus 68 (MHV-68) in a dose-dependent manner [120].

Ramayanti et al. reported that curcumin and its analogues significantly induced EBV reactivation in nasopharyngeal and gastric carcinomas cells [127]. Moreover, curcuminoids, especially compound EF24 (curcumin with the replacement of the β -diketone by piperidinone), displayed exceptional potential as an EBV lytic activator [127]. Furthermore, curcumin was found to reduce the viability and promote the apoptosis of EBV (+) nasopharyngeal carcinoma cells [127]. It also inhibited the proliferation of tumour cells by targeting the expression of EBV nuclear antigen 1 and exerted an antitumor effect [74].

4.2. Structural Modifications of Natural Products for Enhanced Activity

Natural antiviral agents thus far have not been highly effective in inhibiting EBV and have often displayed cytotoxicity, which limits their application to treat EBV infections. Structural modification of natural compounds and synthesis of new compounds have been conducted in the literature to overcome these limitations. For example, coumarin (+)-rutamarin, found in *Ruta graveolens*, moderately inhibited the EBV lytic cycle (IC₅₀ = 7.0 μ M) and its potency was not adequate for further investigation [100]. Twenty-eight (-)-rutamarin were semisynthesized from (-) chalepin, which is a derivative of (+)-rutamarin [100]. Of these, sixteen compounds were potent against EBV lytic DNA replication. The

most effective antiviral compound had an IC₅₀ value of 0.83 μ M (dropped ninefold compared to unmodified (+)-rutamarin) and a selectivity index value of >120 [100].

Protoflavones are natural flavonoids, well known for antitumor properties [128]. However, their cytotoxicity against normal cells is a major concern [129]. Protoapigenone, a protoflavone, was reported to have an inhibitory effect on the EBV lytic cycle [129]. Twenty-seven compounds were prepared from apigenin, aiming to create protoflavones with less cytotoxic effect [129]. Among these semisynthesis derivatives, protoapigenone 1'-O-isopropyl ether was a promising lead with significant activity against EBV, low cytotoxicity, and a selectivity index of 73 [129].

GAP31 (gelonium anti-HIV protein, 31 kDa) protein, first isolated from the plant *Gelonium multifloru*, has shown great antiviral activities against HIV and HPV 1 infection [130]. The authors also reported the antiviral activities of GAP31 (synthetic) against EBV via targeting *EBNA1*, the only latent gene expressed in most EBV neoplasms [131]. *EBNA1* binds to the replication origin (oriP) to display its biological impact on EBV-driven cell transformation and maintenance [131]. Recombinant GAP31 blocked *EBNA1* dimer formation and consequently impaired EBNA1/oriP binding and transcription [131]. The blocking effect of GAP31 on *EBNA1*/oriP-dependent functions included impairing EBV-driven cell proliferation, transformation, and tumorigenesis [131]. In particular, rGAP31 at 0.32 nM reduced approximately 90% of the lymphoblastoid cell lines proliferation by day 15 in that study [131]. Furthermore, the EBV-driven cell transformation outgrowth was entirely abrogated by 0.64 nM rGAP31 [131]. GAP31 also suppressed EBV-mediated cancer formation and reduced the tumour size in the xenograft mouse model [131].

5. Conclusions and Future Direction

Natural sources with great chemical diversity could be the key to discovering novel, potent, and safe antiviral agents against EBV. However, the research on the therapeutic effect of gut microbial metabolites and natural products against EBV infection and cancers is limited. Natural bioactive compounds in plants, fruits and vegetables, microbes, fungi, and marine organisms have been shown to have significant antiviral and anticancer properties in the current literature. Most of the natural antiviral agents studied against EBV are phenolic compounds that suppressed expression and/or interfered with the functions of the EBV lytic proteins with some compounds inhibiting EBNA1. A few agents acted as nucleoside analogues and terminated RNA synthesis. Compounds such as triptolide, wogonin, and berberine have been tested in animal models for their efficacy against EBVinduced infection. Furthermore, the structural modification of natural compounds can be an effective method to improve their efficacy and overcome the limitations (such as cytotoxicity) of the parent compounds. Although several candidates have shown great potential for future drug development and their molecular mechanisms of action are understood up to some extent, further in vivo, and clinical studies are necessary. In particular, understanding their molecular mechanisms of action through more systematic in vitro and in vivo studies is crucial before conducting clinical trials. Similarly, future research is warranted to investigate the effectiveness and molecular mechanisms of gut microbial metabolites and gut microbiome in general against EBV-induced diseases at both preclinical and clinical levels. The potential role of probiotic foods against EBV could be another exciting area of research.

Author Contributions: R.A.E.: conceptualisation, methodology, investigation, writing—original draft, figures, visualisation. H.T.V.: methodology, investigation, writing—original draft. R.S.: methodology, investigation, writing—original draft. D.C.: writing—review. C.G.L.: writing—review. D.J.B.: conceptualisation, design, methodology, investigation, writing—original draft, figures, review and editing, visualisation, funding, supervision. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding. The APC was funded by Western Sydney University, Australia.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: No new data were created.

Acknowledgments: The research was supported by Western Sydney University, Australia. R.A.E. and is supported by Western Sydney University PhD. Scholarship. R.S. was supported by the Western Sydney University Summer Scholarship. D.J.B is supported by the Research Support Program Fellowship at Western Sydney University. The authors acknowledge the open-access funding support from Western Sydney University. Figures 1, 2 and 4–7 were created with biorender.com.

Conflicts of Interest: As a medical research institute, NICM Health Research Institute receives research grants and donations from foundations, universities, government agencies, individuals, and industries. Sponsors and donors also provide untied funding for work to advance the vision and mission of the institute. The authors declare no conflicts of interest.

References

- Sung, H.; Ferlay, J.; Siegel, R.L.; Laversanne, M.; Soerjomataram, I.; Jemal, A.; Bray, F. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J. Clin.* 2021, 71, 209–249. https://doi.org/10.3322/caac.21660.
- 2. Blackadar, C.B. Historical review of the causes of cancer. World J. Clin. Oncol. 2016, 7, 54–86. https://doi.org/10.5306/wjco.v7.i1.54.
- Moore, P.S.; Chang, Y. Why do viruses cause cancer? Highlights of the first century of human tumour virology. *Nat. Rev. Cancer* 2010, 10, 878–889. https://doi.org/10.1038/nrc2961.
- 4. Fatima, I.; Kanwal, S.; Mahmood, T. Natural Products Mediated Targeting of Virally Infected Cancer. *Dose-Response* **2019**, *17*, 1559325818813227. https://doi.org/10.1177/1559325818813227.
- 5. zur Hausen, H.; de Villiers, E.M. Cancer: "causation" by infections--individual contributions and synergistic networks. *Semin. Oncol.* **2014**, *41*, 860–875.
- Bouvard, V.; Baan, R.; Straif, K.; Grosse, Y.; Secretan, B.; El Ghissassi, F.; Benbrahim-Tallaa, L.; Guha, N.; Freeman, C.; Galichet, L.; et al. A review of human carcinogens—Part B: Biological agents. *Lancet Oncol.* 2009, 10, 321–322. https://doi.org/10.1016/s1470-2045(09)70096-8.
- Butel, J.S. Viral carcinogenesis: Revelation of molecular mechanisms and etiology of human disease. *Carcinog.* 2000, 21, 405–426. https://doi.org/10.1093/carcin/21.3.405.
- Chang, Y.; Moore, P.S.; Weiss, R.A. Human oncogenic viruses: Nature and discovery. *Philos. Trans. R. Soc. B Biol. Sci.* 2017, 372, 20160264. https://doi.org/10.1098/rstb.2016.0264.
- 9. Trichopoulos, D. Infections and human cancer: Cancer surveys, volume 33; microbes and malignancy: Infection as a cause of human cancers. *BMJ* **1999**, *319*, 1207.
- 10. Foulkes, W.D. Inherited Susceptibility to Common Cancers. N. Engl. J. Med. 2008, 359, 2143–2153. https://doi.org/10.1056/nejmra0802968.
- Garber, J.E.; Offit, K. Hereditary Cancer Predisposition Syndromes. J. Clin. Oncol. 2005, 23, 276–292. https://doi.org/10.1200/jco.2005.10.042.
- 12. Greaves, M.; Maley, C.C. Clonal evolution in cancer. Nature 2012, 481, 306-313. https://doi.org/10.1038/nature10762.
- Curtius, K.; Wright, N.A.; Graham, T.A. An evolutionary perspective on field cancerization. *Nat. Rev. Cancer* 2018, 18, 19–32. https://doi.org/10.1038/nrc.2017.102.
- 14. Pitot, H.C. The molecular biology of carcinogenesis. *Cancer* **1993**, 72 (Suppl. 3), 962–970. https://doi.org/10.1002/1097-0142(19930801)72:3+<962::aid-cncr2820721303>3.0.co;2-h.
- 15. Morales-Sánchez, A.; Fuentes-Pananá, E.M. Human Viruses and Cancer. Viruses 2014, 6, 4047–4079. https://doi.org/10.3390/v6104047.
- Khidr, L.; Chen, P.-L. RB, the conductor that orchestrates life, death and differentiation. Oncogene 2006, 25, 5210–5219. https://doi.org/10.1038/sj.onc.1209612.
- Cohen, S.B.; Graham, M.E.; Lovrecz, G.O.; Bache, N.; Robinson, P.J.; Reddel, R.R. Protein Composition of Catalytically Active Human Telomerase from Immortal Cells. *Science* 2007, *315*, 1850–1853. https://doi.org/10.1126/science.1138596.
- Terrin, L.; Dal Col, J.; Rampazzo, E.; Zancai, P.; Pedrotti, M.; Ammirabile, G.; Bergamin, S.; Rizzo, S.; Dolcetti, R.; De Rossi, A. Latent Membrane Protein 1 of Epstein-Barr Virus Activates the hTERT Promoter and Enhances Telomerase Activity in B Lymphocytes. J. Virol. 2008, 82, 10175–10187. https://doi.org/10.1128/jvi.00321-08.
- Lewis, B.P.; Burge, C.B.; Bartel, D.P. Conserved Seed Pairing, Often Flanked by Adenosines, Indicates that Thousands of Human Genes are MicroRNA Targets. *Cell* 2005, 120, 15–20. https://doi.org/10.1016/j.cell.2004.12.035.
- 20. Marquitz, A.R.; Raab-Traub, N. The role of miRNAs and EBV BARTs in NPC. Semin. Cancer Biol. 2012, 22, 166–172. https://doi.org/10.1016/j.semcancer.2011.12.001.

- Chimal-Ramírez, G.K.; Espinoza-Sánchez, N.A.; Fuentes-Pananá, E.M. Protumor Activities of the Immune Response: Insights in the Mechanisms of Immunological Shift, Oncotraining, and Oncopromotion. J. Oncol. 2013, 2013, 1–16. https://doi.org/10.1155/2013/835956.
- 22. Chadburn, A.; Abdul-Nabi, A.M.; Teruya, B.S.; Lo, A.A. Lymphoid Proliferations Associated With Human Immunodeficiency Virus Infection. *Arch. Pathol. Lab. Med.* **2013**, *137*, 360–370. https://doi.org/10.5858/arpa.2012-0095-ra.
- Michaelis, M.; Doerr, H.W.; Cinatl, J. The Story of Human Cytomegalovirus and Cancer: Increasing Evidence and Open Questions. *Neoplasia* 2009, 11, 1–9. https://doi.org/10.1593/neo.81178.
- 24. Andrei, G.; Trompet, E.; Snoeck, R. Novel therapeutics for Epstein–Barr virus. Molecules 2019, 24, 997.
- Zhang, S.; Zhu, Y.; Jin, Y.; Sun, H.; Wang, W.; Zhan, L. Difference between Acyclovir and Ganciclovir in the Treatment of Children with Epstein–Barr Virus-Associated Infectious Mononucleosis. *Evid.-Based Complement. Altern. Med.* 2021, 2021, 1–6. https://doi.org/10.1155/2021/8996934.
- 26. El Sayed, K.A. Natural Products as Antiviral Agents. Stud. Nat. Prod. Chem. 2000, 24, 473–572. https://doi.org/10.1016/s1572-5995(00)80051-4.
- 27. Jaye, K.; Li, C.G.; Chang, D.; Bhuyan, D.J. The role of key gut microbial metabolites in the development and treatment of cancer. *Gut Microbes* **2022**, *14*, 2038865. https://doi.org/10.1080/19490976.2022.2038865.
- 28. Jaye, K.; Li, C.G.; Bhuyan, D.J. The complex interplay of gut microbiota with the five most common cancer types: From carcinogenesis to therapeutics to prognoses. *Crit. Rev. Oncol.* **2021**, *165*, 103429. https://doi.org/10.1016/j.critrevonc.2021.103429.
- Ascherio, A.; Munger, K.L. Epstein–Barr Virus Infection and Multiple Sclerosis: A Review. J. Neuroimmune Pharmacol. 2010, 5, 271–277. https://doi.org/10.1007/s11481-010-9201-3.
- Murata, T. Regulation of Epstein-Barr virus reactivation from latency. *Microbiol. Immunol.* 2014, 58, 307–317. https://doi.org/10.1111/1348-0421.12155.
- Roderburg, C.; Krieg, S.; Krieg, A.; Luedde, T.; Kostev, K.; Loosen, S.H. The Association between Infectious Mononucleosis and Cancer: A Cohort Study of 24,190 Outpatients in Germany. *Cancers* 2022, 14, 5837. https://doi.org/10.3390/cancers14235837.
- 32. Kenney, S.C.; Mertz, J.E. Regulation of the latent-lytic switch in Epstein–Barr virus. Semin. Cancer Biol. 2014, 26, 60–68.
- 33. Münz, C. Latency and lytic replication in Epstein–Barr virus-associated oncogenesis. *Nat. Rev. Genet.* 2019, *17*, 691–700. https://doi.org/10.1038/s41579-019-0249-7.
- 34. Saha, A.; Robertson, E.S. Mechanisms of B-Cell Oncogenesis Induced by Epstein-Barr Virus. J. Virol. 2019, 93, e00238-19. https://doi.org/10.1128/jvi.00238-19.
- Mesri, E.A.; Feitelson, M.A.; Munger, K. Human Viral Oncogenesis: A Cancer Hallmarks Analysis. Cell Host Microbe 2014, 15, 266–282. https://doi.org/10.1016/j.chom.2014.02.011.
- 36. Thompson, M.P.; Kurzrock, R. Epstein-Barr virus and cancer. Clin. Cancer Res. 2004, 10, 803–821.
- 37. Niedobitek, G.; Young, L.; Herbst, H. Epstein-Barr virus infection and the pathogenesis of malignant lymphomas. *Cancer Surv.* **1997**, *30*, 143–162.
- Babcock, G.J.; Hochberg, D.; Thorley-Lawson, D.A. The Expression Pattern of Epstein-Barr Virus Latent Genes In Vivo Is Dependent upon the Differentiation Stage of the Infected B Cell. *Immunity* 2000, 13, 497–506. https://doi.org/10.1016/s1074-7613(00)00049-2.
- Kamranvar, S.A.; Masucci, M.G. Regulation of Telomere Homeostasis during Epstein-Barr virus Infection and Immortalization. Viruses 2017, 9, 217. https://doi.org/10.3390/v9080217.
- Portis, T.; Longnecker, R. Epstein–Barr virus (EBV) LMP2A mediates B-lymphocyte survival through constitutive activation of the Ras/PI3K/Akt pathway. Oncogene 2004, 23, 8619–8628. https://doi.org/10.1038/sj.onc.1207905.
- 41. Sbih-Lammali, F.; Djennaoui, D.; Belaoui, H.; Bouguermouh, A.; Decaussin, G.; Ooka, T., Transcriptional expression of Epstein-Barr virus genes and proto-oncogenes in north African nasopharyngeal carcinoma. J. Med. Virol. **1996**, 49, 7–14.
- 42. zur Hausen, H.; O'Neill, F.J.; Freese, U.K.; Hecker, E., Persisting oncogenic herpesvirus induced by the tumour promotor TPA. *Nature* **1978**, 272, 373–375.
- 43. Soldan, S.S.; Lieberman, P.M., Epstein–Barr virus and multiple sclerosis. Nat. Rev. Microbiol. 2022, 21, 51–64.
- 44. Fan, Y.; Pedersen, O. Gut microbiota in human metabolic health and disease. *Nat. Rev. Microbiol.* 2021, 19, 55–71. https://doi.org/10.1038/s41579-020-0433-9.
- 45. Silva, Y.P.; Bernardi, A.; Frozza, R.L. The Role of Short-Chain Fatty Acids From Gut Microbiota in Gut-Brain Communication. *Front. Endocrinol.* **2020**, *11*, 25. https://doi.org/10.3389/fendo.2020.00025.
- Sliva, J.; Pantzartzi, C.N.; Votava, M. Inosine Pranobex: A Key Player in the Game Against a Wide Range of Viral Infections and Non-Infectious Diseases. *Adv. Ther.* 2019, *36*, 1878–1905. https://doi.org/10.1007/s12325-019-00995-6.
- 47. Cotter, P.; Hill, C.; Ross, R. Bacteriocins: Developing innate immunity for food. *Nat. Rev. Genet.* 2005, *3*, 777–788. https://doi.org/10.1038/nrmicro1273.
- Kesika, P.; Sivamaruthi, B.S.; Thangaleela, S.; Chaiyasut, C. The Antiviral Potential of Probiotics—A Review on Scientific Outcomes. *Appl. Sci.* 2021, 11, 8687. https://doi.org/10.3390/app11188687.
- Gleeson, M.; Bishop, N.C.; Struszczak, L. Effects of Lactobacillus casei Shirota ingestion on common cold infection and herpes virus antibodies in endurance athletes: A placebo-controlled, randomized trial. *Eur. J. Appl. Physiol.* 2016, 116, 1555–1563. https://doi.org/10.1007/s00421-016-3415-x.
- Cummings, J.H.; Pomare, E.W.; Branch, W.J.; Naylor, C.P.; Macfarlane, G.T. Short chain fatty acids in human large intestine, portal, hepatic and venous blood. *Gut* 1987, 28, 1221–1227. https://doi.org/10.1136/gut.28.10.1221.

- 51. Schönfeld, P.; Wojtczak, L. Short-and medium-chain fatty acids in energy metabolism: The cellular perspective. *J. Lipid Res.* **2016**, 57, 943–954.
- 52. Machado, M.G.; Sencio, V.; Trottein, F. Short-Chain Fatty Acids as a Potential Treatment for Infections: A Closer Look at the Lungs. *Infect. Immun.* 2021, *89*, e00188-21. https://doi.org/10.1128/iai.00188-21.
- 53. Gorres, K.L.; Daigle, D.; Mohanram, S.; Miller, G., Activation and repression of Epstein-Barr virus and Kaposi's sarcoma-associated herpesvirus lytic cycles by short-and medium-chain fatty acids. *J. Virol.* **2014**, *88*, 8028-8044.
- Li, D.; Feng, Y.; Tian, M.; Ji, J.; Hu, X.; Chen, F. Gut microbiota-derived inosine from dietary barley leaf supplementation attenuates colitis through PPARγ signaling activation. *Microbiome* 2021, *9*, 83. https://doi.org/10.1186/s40168-021-01028-7.
- Sumbria, D.; Berber, E.; Rouse, B.T. Supplementing the Diet with Sodium Propionate Suppresses the Severity of Viral Immunoinflammatory Lesions. J. Virol. 2021, 95, e02056-20. https://doi.org/10.1128/jvi.02056-20.
- Li, J.; Richards, E.M.; Handberg, E.M.; Pepine, C.J.; Raizada, M.K. Butyrate Regulates COVID-19–Relevant Genes in Gut Epithelial Organoids From Normotensive Rats. *Hypertension* 2021, 77, e13–e16. https://doi.org/10.1161/hypertensionaha.120.16647.
- 57. Srinivasan, S.; Torres, A.; de Pouplana, L.R. Inosine in Biology and Disease. *Genes* 2021, 12, 600. https://doi.org/10.3390/genes12040600.
- Mager, L.F.; Burkhard, R.; Pett, N.; Cooke, N.C.A.; Brown, K.; Ramay, H.; Paik, S.; Stagg, J.; Groves, R.A.; Gallo, M.; et al. Microbiome-derived inosine modulates response to checkpoint inhibitor immunotherapy. *Science* 2020, 369, 1481–1489. https://doi.org/10.1126/science.abc3421.
- Lasek, W.; Janyst, M.; Wolny, R.; Zapała, Ł.; Bocian, K.; Drela, N. Immunomodulatory effects of inosine pranobex on cytokine production by human lymphocytes. *Acta Pharm.* 2015, 65, 171–180. https://doi.org/10.1515/acph-2015-0015.
- Tsang, K.Y.; Fudenberg, H.H.; Pan, J.F.; Gnagy, M.J.; Bristow, C.B. An in vitro study on the effects of isoprinosine on immune responses in cancer patients. *Int. J. Immunopharmacol.* 1983, *5*, 481–490. https://doi.org/10.1016/0192-0561(83)90041-3.
- Ahmed, S.R.; Newman, A.S.; O'Daly, J.; Duffy, S.; Grafton, G.; Brady, C.A.; Curnow, S.J.; Barnes, N.M.; Gordon, J. Inosine Acedoben Dimepranol promotes an early and sustained increase in the natural killer cell component of circulating lymphocytes: A clinical trial supporting anti-viral indications. *Int. Immunopharmacol.* 2017, 42, 108–114. https://doi.org/10.1016/j.intimp.2016.11.023.
- 62. Bekesi, J.; Tsang, P.; Wallace, J.; Roboz, J. Immunorestorative properties of isoprinosine in the treatment of patients at high risk of developing ARC or AIDS. *J. Clin. Lab. Immunol.* **1987**, *24*, 155–161.
- 63. Tsang, K.Y.; Pan, J.F.; Swanger, D.; Fudenberg, H. In vitro restoration of immune responses in aging humans by isoprinosine. *Int. J. Immunopharmacol.* **1985**, *7*, 199–206. https://doi.org/10.1016/0192-0561(85)90027-x.
- Janíčková, O.; Ančicová, L.; Briestenská, K.; Mistríková, J. The effect of Isoprinosine treatment on persistent infection of Balb/c mice infected with murine gammaherpesvirus 68. Acta Virol. 2017, 61, 32–38. https://doi.org/10.4149/av_2017_01_32.
- Sundar, S.K.; Barile, G.; Menezes, J. Isoprinosine enhances the activation of sensitized lymphocytes by Epstein-Barr virus antigens. *Int. J. Immunopharmacol.* 1985, 7, 187–192. https://doi.org/10.1016/0192-0561(85)90025-6.
- 66. Beran, J.; Šalapová, E.; Špajdel, M. Inosine pranobex is safe and effective for the treatment of subjects with confirmed acute respiratory viral infections: Analysis and subgroup analysis from a Phase 4, randomised, placebo-controlled, double-blind study. BMC Infect. Dis. 2016, 16, 1–10. https://doi.org/10.1186/s12879-016-1965-5.
- Beran, J.; Špajdel, M.; Katzerová,, V.; Holoušová,, A.; Malyš,, J.; Rousková,, J.F.; Slíva, J. Inosine pranobex significantly decreased the case-fatality rate among PCR positive elderly with SARS-CoV-2 at three nursing homes in the Czech Republic. *Pathogens* 2020, *9*, 1055.
- 68. Klaenhammer, T.R. Genetics of bacteriocins produced by lactic acid bacteria. FEMS Microbiol. Rev. 1993, 12, 39–85.
- Al Kassaa, I.; Hober, D.; Hamze, M.; Chihib, N.E.; Drider, D. Antiviral Potential of Lactic Acid Bacteria and Their Bacteriocins. Probiotics Antimicrob. Proteins 2014, 6, 177–185. https://doi.org/10.1007/s12602-014-9162-6.
- Todorov, S.D.; Wachsman, M.B.; Knoetze, H.; Meincken, M.; Dicks, L.M. An antibacterial and antiviral peptide produced by Enterococcus mundtii ST4V isolated from soya beans. *Int. J. Antimicrob. Agents* 2005, 25, 508–513. https://doi.org/10.1016/j.ijantimicag.2005.02.005.
- Wachsman, M.B.; Castilla, V.; de Ruiz Holgado, A.P.; de Torres, R.A.; Sesma, F.; Coto, C.E. Enterocin CRL35 inhibits late stages of HSV-1 and HSV-2 replication in vitro. *Antivir. Res.* 2003, 58, 17–24. https://doi.org/10.1016/s0166-3542(02)00099-2.
- 72. Férir, G.; Petrova, M.I.; Andrei, G.; Huskens, D.; Hoorelbeke, B.; Snoeck, R.; Vanderleyden, J.; Balzarini, J.; Bartoschek, S.; Brönstrup, M.; et al. The Lantibiotic Peptide Labyrinthopeptin A1 Demonstrates Broad Anti-HIV and Anti-HSV Activity with Potential for Microbicidal Applications. *PLoS ONE* 2013, *8*, e64010. https://doi.org/10.1371/journal.pone.0064010.
- Torres, N.I.; Noll, K.S.; Xu, S.; Li, J.; Huang, Q.; Sinko, P.J.; Wachsman, M.B.; Chikindas, M.L. Safety, formulation and in vitro antiviral activity of the antimicrobial peptide subtilosin against herpes simplex virus type 1. *Probiotics Antimicrob.* 2013, *5*, 26– 35.
- Liu, L.; Yang, J.; Ji, W.; Wang, C. Curcumin Inhibits Proliferation of Epstein–Barr Virus-Associated Human Nasopharyngeal Carcinoma Cells by Inhibiting EBV Nuclear Antigen 1 Expression. *BioMed Res. Int.* 2019, 2019, 1–10. https://doi.org/10.1155/2019/8592921.
- Lin, T.-P.; Chen, S.-Y.; Duh, P.-D.; Chang, L.-K.; Liu, Y.-N. Inhibition of the Epstein-Barr Virus Lytic Cycle by Andrographolide. Biol. Pharm. Bull. 2008, 31, 2018–2023. https://doi.org/10.1248/bpb.31.2018.

- 76. Kapadia, G.J.; Azuine, M.A.; Tokuda, H.; Takasaki, M.; Mukainaka, T.; Konoshima, T.; Nishino, H. Chemopreventive effect of resveratrol, sesamol, sesame oil and sunflower oil in the Epstein–Barr virus early antigen activation assay and the mouse skin two-stage carcinogenesis. *Pharmacol. Res.* 2002, 45, 499–505.
- 77. Li, T.; Peng, T. Traditional Chinese herbal medicine as a source of molecules with antiviral activity. *Antivir. Res.* **2013**, *97*, 1–9. https://doi.org/10.1016/j.antiviral.2012.10.006.
- Musarra-Pizzo, M.; Pennisi, R.; Ben-Amor, I.; Mandalari, G.; Sciortino, M. Antiviral Activity Exerted by Natural Products against Human Viruses. *Viruses* 2021, 13, 828. https://doi.org/10.3390/v13050828.
- Hassan, S.T.; Masarčíková, R.; Berchová, K. Bioactive natural products with anti-herpes simplex virus properties. J. Pharm. Pharmacol. 2015, 67, 1325–1336.
- Ruchawapol, C.; Yuan, M.; Wang, S.-M.; Fu, W.-W.; Xu, H.-X. Natural Products and Their Derivatives against Human Herpesvirus Infection. *Molecules* 2021, 26, 6290. https://doi.org/10.3390/molecules26206290.
- Zhang, H.; Li, C.; Kwok, S.-T.; Zhang, Q.-W.; Chan, S.-W. A Review of the Pharmacological Effects of the Dried Root of *Polygonum cuspidatum*(Hu Zhang) and Its Constituents. *Evid.-Based Complement. Altern. Med.* 2013, 2013, 1–13. https://doi.org/10.1155/2013/208349.
- 82. Yiu, C.-Y.; Chen, S.-Y.; Chen, Y.-P.; Lin, T.-P. Inhibition of the ethanolic extract from Polygonum cuspidatum root on the functions of Epstein-Barr Virus latent membrane protein 1. J. Food Drug Anal. 2013, 21, 20–26.
- Yiu, C.-Y.; Chen, S.-Y.; Huang, C.-W.; Yeh, D.-B.; Lin, T.-P. Inhibitory effects of Polygonum cuspidatum on the Epstein-Barr virus lytic cycle. J. Food Drug Anal. 2011, 19, 3. https://doi.org/10.38212/2224-6614.2234.
- Yiu, C.-Y.; Chen, S.-Y.; Yang, T.-H.; Chang, C.-J.; Yeh, D.-B.; Chen, Y.-J.; Lin, T.-P. Inhibition of Epstein-Barr Virus Lytic Cycle by an Ethyl Acetate Subfraction Separated from Polygonum cuspidatum Root and Its Major Component, Emodin. *Molecules* 2014, 19, 1258–1272. https://doi.org/10.3390/molecules19011258.
- 85. Kemboi, D.; Peter, X.; Langat, M.; Tembu, J. A review of the ethnomedicinal uses, biological activities, and triterpenoids of Euphorbia species. *Molecules* **2020**, *25*, 4019.
- Wu, T.; Wang, Q.; Jiang, C.; Morris-Natschke, S.L.; Cui, H.; Wang, Y.; Yan, Y.; Xu, J.; Lee, K.-H.; Gu, Q. neo-Clerodane Diterpenoids from *Scutellaria barbata* with Activity against Epstein–Barr Virus Lytic Replication. J. Nat. Prod. 2015, 78, 500–509. https://doi.org/10.1021/np500988m.
- Liu, S.-N.; Hu, J.; Tan, S.H.; Wang, Q.; Xu, J.; Wang, Y.; Yuan, Y.; Gu, Q. ent-Rosane diterpenoids from Euphorbia milii showing an Epstein–Barr virus lytic replication assay. *RSC Adv.* 2017, 7, 46938–46947. https://doi.org/10.1039/c7ra08877a.
- Wang, C.; Wang, H.; Zhang, Y.; Guo, W.; Long, C.; Wang, J.; Liu, L.; Sun, X. Berberine inhibits the proliferation of human nasopharyngeal carcinoma cells via an Epstein-Barr virus nuclear antigen 1-dependent mechanism. *Oncol. Rep.* 2017, 37, 2109– 2120. https://doi.org/10.3892/or.2017.5489.
- Park, G.B.; Park, S.H.; Kim, D.; Kim, Y.S.; Yoon, S.H.; Hur, D.Y. Berberine induces mitochondrial apoptosis of EBV-transformed B cells through p53-mediated regulation of XAF1 and GADD45α. *Int. J. Oncol.* 2016, 49, 411–421. https://doi.org/10.3892/ijo.2016.3502.
- Tsang, C.M.; Cheung, Y.C.; Lui, V.W.-Y.; Yip, Y.L.; Zhang, G.; Lin, V.W.; Cheung, K.C.-P.; Feng, Y.; Tsao, S.W. Berberine suppresses tumorigenicity and growth of nasopharyngeal carcinoma cells by inhibiting STAT3 activation induced by tumor associated fibroblasts. *BMC Cancer* 2013, *13*, 619–619. https://doi.org/10.1186/1471-2407-13-619.
- Zhou, F.; Hu, J.; Dai, N.; Song, L.; Lin, T.; Liu, J.; Li, K.; Peng, Z.; He, Y.; Liao, D.-F. Berberine and ginsenoside Rg3 act synergistically via the MAPK/ERK pathway in nasopharyngeal carcinoma cells. *J. Funct. Foods* 2020, 66, 103802. https://doi.org/10.1016/j.jff.2020.103802.
- Wong, K.-F.; Yuan, Y.; Luk, J. Tripterygium wilfordii bioactive compounds as anticancer and anti-inflammatory agents. *Clin. Exp. Pharmacol. Physiol.* 2012, 39, 311–320. https://doi.org/10.1111/j.1440-1681.2011.05586.x.
- Zhou, H.; Guo, W.; Long, C.; Wang, H.; Wang, J.; Sun, X. Triptolide inhibits proliferation of Epstein–Barr virus-positive B lymphocytes by down-regulating expression of a viral protein LMP1. *Biochem. Biophys. Res. Commun.* 2015, 456, 815–820. https://doi.org/10.1016/j.bbrc.2014.12.023.
- Zhou, H.; Liu, Y.; Wang, C.; Liu, L.; Wang, H.; Zhang, Y.; Long, C.; Sun, X. Triptolide inhibits Epstein-Barr nuclear antigen 1 expression by increasing sensitivity of mitochondria apoptosis of nasopharyngeal carcinoma cells. *J. Exp. Clin. Cancer Res.* 2018, 37, 192. https://doi.org/10.1186/s13046-018-0865-5.
- Carter, B.Z.; Mak, D.H.; Schober, W.D.; McQueen, T.; Harris, D.; Estrov, Z.; Evans, R.L.; Andreeff, M. Triptolide induces caspasedependent cell death mediated via the mitochondrial pathway in leukemic cells. *Blood* 2006, 108, 630–637. https://doi.org/10.1182/blood-2005-09-3898.
- 96. Yiu, C.-Y.; Chen, S.-Y.; Chang, L.-K.; Chiu, Y.-F.; Lin, T.-P. Inhibitory Effects of Resveratrol on the Epstein-Barr Virus Lytic Cycle. *Molecules* **2010**, *15*, 7115–7124. https://doi.org/10.3390/molecules15107115.
- De Leo, A.; Arena, G.; Lacanna, E.; Oliviero, G.; Colavita, F.; Mattia, E. Resveratrol inhibits Epstein Barr Virus lytic cycle in Burkitt's lymphoma cells by affecting multiple molecular targets. *Antivir. Res.* 2012, *96*, 196–202. https://doi.org/10.1016/j.antiviral.2012.09.003.
- Espinoza, J.L.; Takami, A.; Trung, L.Q.; Kato, S.; Nakao, S. Resveratrol Prevents EBV Transformation and Inhibits the Outgrowth of EBV-Immortalized Human B Cells. *PLoS ONE* 2012, 7, e51306. https://doi.org/10.1371/journal.pone.0051306.
- 99. Wu, C.-C.; Fang, C.-Y.; Cheng, Y.-J.; Hsu, H.-Y.; Chou, S.-P.; Huang, S.-Y.; Tsai, C.-H.; Chen, J.-Y. Inhibition of Epstein-Barr virus reactivation by the flavonoid apigenin. *J. Biomed. Sci.* 2017, 24, 1–13. https://doi.org/10.1186/s12929-016-0313-9.

- 100. Lin, Y.; Wang, Q.; Gu, Q.; Zhang, H.; Jiang, C.; Hu, J.; Wang, Y.; Yan, Y.; Xu, J. Semisynthesis of (-)-Rutamarin Derivatives and Their Inhibitory Activity on Epstein–Barr Virus Lytic Replication. J. Nat. Prod. 2017, 80, 53–60. https://doi.org/10.1021/acs.jnatprod.6b00415.
- 101. Wu, X.; Liu, P.; Zhang, H.; Li, Y.; Salmani, J.M.M.; Wang, F.; Yang, K.; Fu, R.; Chen, Z.; Chen, B. Wogonin as a targeted therapeutic agent for EBV (+) lymphoma cells involved in LMP1/NF-κB/miR-155/PU. 1 pathway. *BMC Cancer* **2017**, *17*, 1–11.
- Chang, L.-K.; Wei, T.-T.; Chiu, Y.-F.; Tung, C.-P.; Chuang, J.-Y.; Hung, S.-K.; Li, C.; Liu, S.-T. Inhibition of Epstein–Barr virus lytic cycle by (–)-epigallocatechin gallate. *Biochem. Biophys. Res. Commun.* 2003, 301, 1062–1068.
- 103. Liu, S.; Li, H.; Chen, L.; Yang, L.; Li, L.; Tao, Y.; Li, W.; Li, Z.; Liu, H.; Tang, M. (-)-Epigallocatechin-3-gallate inhibition of Epstein–Barr virus spontaneous lytic infection involves ERK1/2 and PI3-K/Akt signaling in EBV-positive cells. *Carcinogenesis* 2013, 34, 627–637.
- Liu, S.; Li, H.; Tang, M.; Cao, Y. (-)-Epigallocatechin-3-gallate inhibition of Epstein-Barr virus spontaneous lytic infection involves downregulation of latent membrane protein 1. *Exp. Ther. Med.* 2018, 15, 1105–1112. https://doi.org/10.3892/etm.2017.5495.
- 105. Lin, J.-C.; Cherng, J.-M.; Hung, M.-S.; Baltina, L.A.; Baltina, L.; Kondratenko, R. Inhibitory effects of some derivatives of glycyrrhizic acid against Epstein-Barr virus infection: Structure–activity relationships. *Antivir. Res.* 2008, 79, 6–11. https://doi.org/10.1016/j.antiviral.2008.01.160.
- 106. Bentz, G.L.; Lowrey, A.J.; Horne, D.C.; Nguyen, V.; Satterfield, A.R.; Ross, T.D.; Harrod, A.E.; Uchakina, O.N.; McKallip, R.J. Using glycyrrhizic acid to target sumoylation processes during Epstein-Barr virus latency. *PLoS ONE* 2019, 14, e0217578. https://doi.org/10.1371/journal.pone.0217578.
- 107. Long, C.; Xu, Q.-B.; Ding, L.; Yang, L.; Ji, W.; Gao, F.; Ji, Y. Triptolide inhibits human telomerase reverse transcriptase by down-regulating translation factors SP1 and c-Myc in Epstein-Barr virus-positive B lymphocytes. Oncol. Lett. 2021, 21, 1–1. https://doi.org/10.3892/ol.2021.12541.
- 108. Tsai, Y.-C.; Hohmann, J.; El-Shazly, M.; Chang, L.-K.; Dankó, B.; Kúsz, N.; Hsieh, C.-T.; Hunyadi, A.; Chang, F.-R. Bioactive constituents of Lindernia crustacea and its anti-EBV effect via Rta expression inhibition in the viral lytic cycle. *J. Ethnopharmacol.* 2020, 250, 112493. https://doi.org/10.1016/j.jep.2019.112493.
- Wu, C.-C.; Chen, M.-S.; Cheng, Y.-J.; Ko, Y.-C.; Lin, S.-F.; Chiu, I.-M.; Chen, J.-Y. Emodin Inhibits EBV Reactivation and Represses NPC Tumorigenesis. *Cancers* 2019, 11, 1795. https://doi.org/10.3390/cancers11111795.
- Hu, L.; Zhang, Y.; Zhu, H.; Liu, J.; Li, H.; Li, X.-N.; Sun, W.; Zeng, J.; Xue, Y.; Zhang, Y. Filicinic Acid Based Meroterpenoids with Anti-Epstein–Barr Virus Activities from *Hypericum japonicum*. Org. Lett. 2016, 18, 2272–2275. https://doi.org/10.1021/acs.orglett.6b00906.
- Wu, R.; Le, Z.; Wang, Z.; Tian, S.; Xue, Y.; Chen, Y.; Hu, L.; Zhang, Y. Hyperjaponol H, A New Bioactive Filicinic Acid-Based Meroterpenoid from Hypericum japonicum Thunb. ex Murray. *Molecules* 2018, 23, 683. https://doi.org/10.3390/molecules23030683.
- 112. Luo, X.; Hong, L.; Cheng, C.; Li, N.; Zhao, X.; Shi, F.; Liu, J.; Fan, J.; Zhou, J.; Bode, A.M.; et al. DNMT1 mediates metabolic reprogramming induced by Epstein–Barr virus latent membrane protein 1 and reversed by grifolin in nasopharyngeal carcinoma. *Cell Death Dis.* **2018**, *9*, 6. https://doi.org/10.1038/s41419-018-0662-2.
- Lee, M.; Son, M.; Ryu, E.; Shin, Y.S.; Kim, J.G.; Kang, B.W.; Sung, G.-H.; Cho, H.; Kang, H. Quercetin-induced apoptosis prevents EBV infection. Oncotarget 2015, 6, 12603–12624. https://doi.org/10.18632/oncotarget.3687.
- 114. Wang, Q.; Zhu, N.; Hu, J.; Wang, Y.; Xu, J.; Gu, Q.; Lieberman, P.M.; Yuan, Y. The mTOR inhibitor manassantin B reveals a crucial role of mTORC2 signaling in Epstein-Barr virus reactivation. *J. Biol. Chem.* **2020**, 295, 7431–7441. https://doi.org/10.1074/jbc.ra120.012645.
- 115. Tung, C.-P.; Chang, F.-R.; Wu, Y.-C.; Chuang, D.-W.; Hunyadi, A.; Liu, S.-T. Inhibition of the Epstein–Barr virus lytic cycle by protoapigenone. J. Gen. Virol. 2011, 92 Pt 8, 1760–1768. https://doi.org/10.1099/vir.0.031609-0.
- Chang, F.-R.; Hsieh, Y.-C.; Chang, Y.-F.; Lee, K.-H.; Wu, Y.-C.; Chang, L.-K. Inhibition of the Epstein–Barr virus lytic cycle by moronic acid. *Antivir. Res.* 2010, 85, 490–495. https://doi.org/10.1016/j.antiviral.2009.12.002.
- 117. Cui, H.; Xu, B.; Wu, T.; Xu, J.; Yuan, Y.; Gu, Q. Potential Antiviral Lignans from the Roots of *Saururus chinensis* with Activity against Epstein–Barr Virus Lytic Replication. *J. Nat. Prod.* **2014**, *77*, 100–110. https://doi.org/10.1021/np400757k.
- 118. Guo, Q.; Sun, X.; Zhang, Z.; Zhang, L.; Yao, G.; Li, F.; Yang, X.; Song, L.; Jiang, G. The effect of Astragalus polysaccharide on the Epstein-Barr virus lytic cycle.. *Acta Virol.* 2014, *58*, 76–83. https://doi.org/10.4149/av_2014_01_76.
- 119. Kok, Y.-Y.; Chu, W.-L.; Phang, S.-M.; Mohamed, S.M.; Naidu, R.; Lai, P.-J.; Ling, S.-N.; Mak, J.-W.; Lim, P.K.-C.; Balraj, P.; et al. Inhibitory activities of microalgal extracts against Epstein-Barr virus DNA release from lymphoblastoid cells. J. Zhejiang Univ. Sci. B 2011, 12, 335–345. https://doi.org/10.1631/jzus.b1000336.
- 120. Cho, H.-J.; Jeong, S.-G.; Park, J.-E.; Han, J.-A.; Kang, H.-R.; Lee, D.; Song, M.J. Antiviral activity of angelicin against gammaherpesviruses. *Antivir. Res.* 2013, 100, 75–83. https://doi.org/10.1016/j.antiviral.2013.07.009.
- 121. Kapadia, G.J.; Rao, G.S.; Sridhar, R.; Ichiishi, E.; Takasaki, M.; Suzuki, N.; Konoshima, T.; Iida, A.; Tokuda, H. Chemoprevention of skin cancer: Effect of *Lawsonia inermis* L.(Henna) leaf powder and its pigment artifact, lawsone in the Epstein-Barr virus early antigen activation assay and in two-stage mouse skin carcinogenesis models. *Anticancer Agents Med. Chem.* 2013, 13, 1500–1507.
- 122. Wu, C.-C.; Fang, C.-Y.; Hsu, H.-Y.; Chen, Y.-J.; Chou, S.-P.; Huang, S.-Y.; Cheng, Y.-J.; Lin, S.-F.; Chang, Y.; Tsai, C.-H.; et al. Luteolin inhibits Epstein-Barr virus lytic reactivation by repressing the promoter activities of immediate-early genes. *Antivir. Res.* 2016, *132*, 99–110. https://doi.org/10.1016/j.antiviral.2016.05.007.

- 123. Wang, D.; Tan, G.T.; Van Hung, N.; Cuong, N.M.; Pezzuto, J.M.; Fong, H.H.; Soejarto, D.D.; Zhang, H. *Litsea* Species as Potential Antiviral Plant Sources. *Am. J. Chin. Med.* **2016**, *44*, 275–290. https://doi.org/10.1142/s0192415x16500166.
- 124. Ryu, E.; Son, M.; Lee, M.; Lee, K.; Cho, J.Y.; Cho, S.; Lee, S.K.; Lee, Y.M.; Cho, H.; Sung, G.-H.; et al. Cordycepin is a novel chemical suppressor of Epstein-Barr virus replication. *Oncoscience* **2014**, *1*, 866–881. https://doi.org/10.18632/oncoscience.110.
- 125. Cao, Y.; Xie, L.; Shi, F.; Tang, M.; Li, Y.; Hu, J.; Zhao, L.; Zhao, L.; Yu, X.; Luo, X. Targeting the signaling in Epstein–Barr virusassociated diseases: Mechanism, regulation, and clinical study. *Signal Transduct. Target Ther.* **2021**, *6*, 1–33.
- 126. De Leo, A.; Arena, G.; Stecca, C.; Raciti, M.; Mattia, E. Resveratrol Inhibits Proliferation and Survival of Epstein Barr Virus– Infected Burkitt's Lymphoma Cells Depending on Viral Latency Program. *Mol. Cancer Res.* 2011, 9, 1346–1355.
- 127. Ramayanti, O.; Brinkkemper, M.; Verkuijlen, S.A.; Ritmaleni, L.; Go, M.L.; Middeldorp, J.M. Curcuminoids as EBV Lytic Activators for Adjuvant Treatment in EBV-Positive Carcinomas. *Cancers* **2018**, *10*, 89. https://doi.org/10.3390/cancers10040089.
- 128. Hunyadi, A.; Martins, A.; Danko, B.; Chang, F.-R.; Wu, Y.C. Protoflavones: A class of unusual flavonoids as promising novel anticancer agents. *Phytochem. Rev.* 2014, *13*, 69–77. https://doi.org/10.1007/s11101-013-9288-2.
- 129. Vágvölgyi, M.; Girst, G.; Kúsz, N.; Ötvös, S.B.; Fülöp, F.; Hohmann, J.; Servais, J.-Y.; Seguin-Devaux, C.; Chang, F.-R.; Chen, M.S.; et al. Less Cytotoxic Protoflavones as Antiviral Agents: Protoapigenone 1'-O-isopropyl ether Shows Improved Selectivity Against the Epstein–Barr Virus Lytic Cycle. Int. J. Mol. Sci. 2019, 20, 6269. https://doi.org/10.3390/ijms20246269.
- 130. Stirpe, F.; Olsnes, S.; Pihl, A. Gelonin, a new inhibitor of protein synthesis, nontoxic to intact cells. Isolation, characterization, and preparation of cytotoxic complexes with concanavalin A. *J. Biol. Chem.* **1980**, 255, 6947–6953.
- 131. Shen, C.-L.; Huang, W.-H.; Hsu, H.-J.; Yang, J.-H.; Peng, C.-W. GAP31 from an ancient medicinal plant exhibits anti-viral activity through targeting to Epstein-Barr virus nuclear antigen 1. *Antivir. Res.* 2019, *164*, 123–130. https://doi.org/10.1016/j.antivi-ral.2019.02.015.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.