International Journal of
Molecular Sciences

MDPI

Correction

Correction: Micheli et al. Beneficial Effect of H_2S-Releasing Molecules in an In Vitro Model of Sarcopenia: Relevance of Glucoraphanin. *Int. J. Mol. Sci.* 2022, *23*, 5955

Laura Micheli [1,†], Emma Mitidieri [2,†], Carlotta Turnaturi [2], Domenico Vanacore [2], Clara Ciampi [1], Elena Lucarini [1], Giuseppe Cirino [2], Carla Ghelardini [1], Raffaella Sorrentino [3], Lorenzo Di Cesare Mannelli [1,*] and Roberta d'Emmanuele di Villa Bianca [2]

1 Department of Neuroscience, Psychology, Drug Research and Child Health-Neurofarba—Section of Pharmacology and Toxicology, University of Florence, 50139 Florence, Italy; laura.micheli@unifi.it (L.M.); clara.ciampi@stud.unifi.it (C.C.); elena.lucarini@unifi.it (E.L.); carla.ghelardini@unifi.it (C.G.)
2 Department of Pharmacy, School of Medicine and Surgery, University of Naples Federico II, 80131 Naples, Italy; emma.mitidieri@unina.it (E.M.); carlotta.turnaturi@unina.it (C.T.); domenico.vanacore@unina.it (D.V.); cirino@unina.it (G.C.); demmanue@unina.it (R.d.d.V.B.)
3 Department of Molecular Medicine and Medical Biotechnology, School of Medicine, University of Naples Federico II, 80131 Naples, Italy; raffaella.sorrentino@unina.it
* Correspondence: lorenzo.mannelli@unifi.it
† These authors contributed equally to this work.

In the original publication [1], there was a mistake in Figure 5 as published. During the assembly of the representative panel, the condition "dexamethasone + glucoraphanin" of carbonylated proteins and the house-keeping protein β-Actin were unintentionally represented by wrong images. The corrected Figure 5 appears below.




Figure 5. Protein carbonylation in C2C12 myotubes. C2C12 myotubes were treated for 48 h with 1 μM DEX, both alone and in the presence of 30 μM glucoraphanin, 150 μM L-cysteine and 150 μM

check for updates

Received: 16 April 2025
Accepted: 21 April 2025
Published: 8 May 2025

Citation: Micheli, L.; Mitidieri, E.; Turnaturi, C.; Vanacore, D.; Ciampi, C.; Lucarini, E.; Cirino, G.; Ghelardini, C.; Sorrentino, R.; Di Cesare Mannelli, L.; et al. Correction: Micheli et al. Beneficial Effect of H_2S-Releasing Molecules in an In Vitro Model of Sarcopenia: Relevance of Glucoraphanin. *Int. J. Mol. Sci.* 2022, *23*, 5955. *Int. J. Mol. Sci.* **2025**, *26*, 4476. https://doi.org/10.3390/ijms26104476

Copyright: © 2025 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).

mercaptopyruvate. Western blot analysis was carried out on cell homogenates using a specific antibody against DNPH. Figures show the densitometric analysis (**top**) and the illustrative immunoblot (**bottom**). Each sample was normalized to β-Actin expression. Values are expressed % of integrated density and calculated as mean ± S.E.M. of 3 different experiments. ** $p < 0.01$ versus control; ^ $p < 0.05$ and ^^ $p < 0.01$ versus DEX.

The authors state that the scientific conclusions are unaffected. This correction was approved by the academic editor. The original publication has also been updated.

Reference

1. Micheli, L.; Mitidieri, E.; Turnaturi, C.; Vanacore, D.; Ciampi, C.; Lucarini, E.; Cirino, G.; Ghelardini, C.; Sorrentino, R.; Di Cesare Mannelli, L.; et al. Beneficial Effect of H_2S-Releasing Molecules in an In Vitro Model of Sarcopenia: Relevance of Glucoraphanin. *Int. J. Mol. Sci.* **2022**, *23*, 5955. [CrossRef] [PubMed]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.