

## Short Note

# $N'^1,N'^4$ -bis(2-(7,8-dimethyl-2,4-dioxo-3,4-dihydrobenzo[g]pteridin-10(2H)-yl)ethylidene)succinohydrazide

Taishu Kawada <sup>1</sup>, Katsuhito Kino <sup>2,\*</sup>, Yoshihiko Matsuzawa <sup>2</sup>, Masayuki Morikawa <sup>3</sup>, Yasuko Okamoto <sup>3</sup>, Takanobu Kobayashi <sup>1</sup>  and Yoshiyuki Tanaka <sup>3</sup> 

<sup>1</sup> Kagawa School of Pharmaceutical Sciences, Tokushima Bunri University, 1314-1 Shido, Sanuki-shi, Kagawa 769-2193, Japan

<sup>2</sup> Faculty of Science and Engineering, Tokushima Bunri University, 1314-1 Shido, Sanuki-shi, Kagawa 769-2193, Japan

<sup>3</sup> Faculty of Pharmaceutical Sciences, Tokushima Bunri University, 180 Nishihama-Boji, Yamashiro-cho, Tokushima 770-8514, Japan

\* Correspondence: kkino@kph.bunri-u.ac.jp; Tel.: +81-87-899-7290

**Abstract:** The title compound,  $N'^1,N'^4$ -bis(2-(7,8-dimethyl-2,4-dioxo-3,4-dihydrobenzo[g]pteridin-10(2H)-yl)ethylidene)succinohydrazide (**1**), was obtained by the reaction of formylmethylflavin and succinic acid dihydrazide. The product **1** was characterized by <sup>1</sup>H-NMR, <sup>13</sup>C-NMR, HRMS and UV.

**Keywords:** flavin; bisflavin; hydrazide; Schiff base



**Citation:** Kawada, T.; Kino, K.; Matsuzawa, Y.; Morikawa, M.; Okamoto, Y.; Kobayashi, T.; Tanaka, Y.  $N'^1,N'^4$ -bis(2-(7,8-dimethyl-2,4-dioxo-3,4-dihydrobenzo[g]pteridin-10(2H)-yl)ethylidene)succinohydrazide. *Molbank* **2022**, *2022*, M1436. <https://doi.org/10.3390/M1436>

Academic Editor: Luke R. Odell

Received: 29 July 2022

Accepted: 26 August 2022

Published: 30 August 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).


## 1. Introduction

Flavins, chemical compounds containing an isoalloxazine ring in their molecule, are part of the complex redox centers of proton pumps and act as photoreceptors in phototropism [1]. Flavins are a large group of compounds involved in biological processes such as photosynthesis or phototropism [1]. They are found in food products, and they are present in enzymes and photoreceptors [1]. They are active chromophores in the processes of oxidation and reduction that take place in cells. It is well known that flavin acts as a cofactor for oxidoreductases, and this property is exploited in several biological and chemical oxidation reactions [1]. For example, it was previously reported that DNA in the presence of riboflavin is damaged under UV-A irradiation [2–6], with the ability to cause mutations associated with this [7–9]. To apply the oxidizing capacity of flavin for various purposes, formylmethylflavin (FMF) is used as a raw material for further derivatization [10–19].

Previous studies have reported bisflavin compounds having two flavins [20], and these molecules have a new catalytic activity or molecular recognition [21–24]. We report here the facile synthesis of a novel bisflavin. The title compound,  $N'^1,N'^4$ -bis(2-(7,8-dimethyl-2,4-dioxo-3,4-dihydrobenzo[g]pteridin-10(2H)-yl)ethylidene)succinohydrazide (**1**), was synthesized and characterized as follows.

## 2. Results and Discussion

Formylmethylflavin (FMF) was dissolved in acetic acid, a 50% (v/v) aqueous solution. Succinic acid dihydrazide was added to this solution, and the mixture was stirred for 30 min at room temperature (Figure 1), and the precipitation was formed. This precipitation, product **1**, was characterized as the desired product by NMR (<sup>1</sup>H-NMR and <sup>13</sup>C-NMR), HRMS, and UV analysis (see Supplementary Files). The structure of **1** is shown in Figure 1.



**Figure 1.** Synthesis of *N',N'*-bis(2-(7,8-dimethyl-2,4-dioxo-3,4-dihydrobenzo[*g*]pteridin-10(2*H*)-yl)ethylidene)succinohydrazide (**1**).

The precipitate was dissolved in AcOH, and silica gel TLC showed a large spot ( $R_f = 0.40$ ) and a small one ( $R_f = 0.80$ ). Only the small spot was extracted from the silica gel with solvent, and the TLC analysis of the extracted sample showed both spots having  $R_f = 0.40$  and  $0.80$ , respectively. When the solvent was changed from AcOH to TFA, the spot with an  $R_f = 0.80$  was larger than that with an  $R_f = 0.40$ . Therefore, the sample with the  $R_f = 0.80$  spot may have contained a protonated bisflavin.

Bisflavin **1** was poorly soluble in many solvents and barely soluble in DMSO. The  $^1\text{H}$ -NMR measurement of this compound was determined by supersaturation concentration. However, because the supersaturation state could not be maintained for the  $^{13}\text{C}$ -NMR measurements over a long time, a mixture of AcOH- $d_4$  and TFA- $d$  was used. The peak assignments were made based on a previous report [16] and predictions using ChemOffice. In addition, the  $^1\text{H}$ -NMR measurements were confirmed by determining the exchangeable protons via the addition of  $\text{D}_2\text{O}$  and by COSY analysis. We also believe that the complexity of the spectrum is due to structural polymorphism. For the  $^{13}\text{C}$ -NMR measurements, DEPT was also performed. Peak assignments are shown in the Materials and Methods section.

It has been previously reported that bisflavin makes  $\pi$ - $\pi$  interactions, which change its redox state [24,25]. Therefore, bisflavin may have a different photoreactivity than riboflavin, which will be examined in future works. In addition, bisflavin has been investigated as a potential inhibitor of amyloid- $\beta$  [26], and the synthesized product **1** might act as some kind of inhibitor. Finally, the simple synthetic method reported here will lead to the rapid synthesis of other bisflavins.

### 3. Materials and Methods

FMF was synthesized via known methods [13]. FMF (142 mg, 0.5 mmol) was dissolved in 9 M acetic acid (20 mL, Wako Pure Chemical Industries, Ltd., Osaka, Japan). Succinic acid dihydrazide (36 mg, 0.25 mmol, Kanto Chemical Co., Inc., Tokyo, Japan) was added to this solution and the mixture was stirred for 30 min at room temperature. The resulting precipitate was filtered under a vacuum and washed with water and MeOH (Wako Pure Chemical Industries, Ltd., Osaka, Japan). The precipitate yielded 101 mg (60%, based on succinic acid dihydrazide) of a light-yellow solid. The reactions were monitored by chromatography on TLC Silica Gel 60 glass plates (Merck KGaA, Darmstadt, Germany) using BuOH:AcOH:H<sub>2</sub>O = 4:1:1 (BuOH; Kishida chemical Co., Ltd., Osaka, Japan) as an eluent. FMF had an  $R_f$  value of 0.82. The precipitate contained a product showing two  $R_f$  values of 0.40 and 0.80.

The melting/decomposition point was determined with “OptiMelt Automated Melting Point System” (MPA100; Stanford Research Systems, Sunnyvale, CA, USA).  $^1\text{H}$ -NMR and  $^{13}\text{C}$ -NMR spectra were recorded on an Avance 400 (Bruker BioSpin, Rheinstetten, Germany) using DMSO- $d_6$  (Acros Organics, Geel, Belgium) and AcOH- $d_4$ :TFA- $d$  = 10:1 (Cambridge isotope laboratories, Inc., Tewksbury, Massachusetts, USA) as the solvent. Mass spectra were recorded on a mass spectrometer “SYNAPT G2-Si HDMS” (Waters, Milford, MA, USA). UV absorption spectra (220 to 900 nm) were recorded on an Ultrospec

3100 pro (GE Healthcare Japan Corporation, Tokyo, Japan) in DMSO (Wako Pure Chemical Industries, Ltd.).

Melting point: >233 °C (decomposition)

UV (DMSO):  $\lambda_{\max}$  440 nm (log  $\epsilon$  4.29), 345 nm (4.10), 270 nm (4.70)

HRESI-Q-TOFMS  $m/z$ : 677.2333 [M-H]<sup>−</sup> (calculated for C<sub>32</sub>H<sub>29</sub>N<sub>12</sub>O<sub>6</sub>, 677.2333)

<sup>1</sup>H-NMR (400 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  11.38 (s, 1H), 11.36 (s, 1H), 11.35 (s, 1H), 11.20 (s, 1H), 11.18 (s, 1H), 11.17 (s, 1H), 11.15 (s, 1H) (11.38–11.15; succinohydrazide 1,4-CONH and flavin 3-NH), 7.95 (s, 1H), 7.92 (s, 1H), 7.91 (s, 1H), 7.90 (s, 1H), 7.86 (s, 1H), 7.81 (s, 1H), 7.77 (s, 1H), 7.74 (s, 1H), 7.72 (s, 1H) (7.95–7.72; flavin H-6,9), 7.73 (t, 1H,  $J$  = 2.8 Hz), 7.37 (t, 1H,  $J$  = 3.6 Hz), 7.35 (t, 1H,  $J$  = 1.5 Hz) (7.73–7.35; flavin N<sup>10</sup>-CH<sub>2</sub>-CHN), 5.44 (d, 2H,  $J$  = 1.5 Hz), 5.41 (d, 2H,  $J$  = 3.6 Hz), 5.37 (d, 2H,  $J$  = 2.8 Hz) (5.44–5.37; flavin N<sup>10</sup>-CH<sub>2</sub>-CHN), 2.48–2.18 ppm (16H, flavin 7,8-CH<sub>3</sub> and CO-CH<sub>2</sub>-CH<sub>2</sub>-CO, overlap with the signal of DMSO)

<sup>13</sup>C-NMR (100 MHz, AcOH-*d*<sub>4</sub>:TFA-*d* = 10:1):  $\delta$  172.5 (succinohydrazide 1-CONH), 156.7 (flavin 2, 4-CO), 156.6 (flavin 2, 4-CO), 145.5 (flavin C<sup>4a</sup>, 5a, 7, 8, 9a, 10a), 144.8 (flavin C<sup>4a</sup>, 5a, 7, 8, 9a, 10a), 141.0 (flavin C<sup>4a</sup>, 5a, 7, 8, 9a, 10a), 140.8 (flavin C<sup>4a</sup>, 5a, 7, 8, 9a, 10a), 132.0 (flavin 6,9-CH), 129.2 (flavin C<sup>4a</sup>, 5a, 7, 8, 9a, 10a), 128.0 (flavin C<sup>4a</sup>, 5a, 7, 8, 9a, 10a), 82.1 (flavin N<sup>10</sup>-CH<sub>2</sub>-CH), 57.9 (flavin N<sup>10</sup>-CH<sub>2</sub>-CH), 53.6 (flavin 7,8-CH<sub>3</sub>), 27.0 ppm (CO-CH<sub>2</sub>-CH<sub>2</sub>-CO)

Silica gel TLC (BuOH:AcOH:H<sub>2</sub>O = 4:1:1) showed spots (UV, 254 nm) with an  $R_f$  = 0.40 and 0.80.

**Supplementary Materials:** The following supporting information can be downloaded online, Supplementary File S1: mol file; Supplementary File S2: <sup>1</sup>H-NMR; Supplementary File S3: <sup>13</sup>C-NMR; Supplementary File S4: HRMS; Supplementary File S5: UV-VIS.

**Author Contributions:** Conceptualization, K.K.; validation, T.K. (Taishu Kawada), K.K., Y.M., M.M. and Y.O.; formal analysis, T.K. (Taishu Kawada), K.K., Y.M., M.M. and Y.O.; investigation, T.K. (Taishu Kawada), K.K., Y.M., M.M., Y.O., T.K. (Takanobu Kobayashi) and Y.T.; resources, K.K., Y.O. and Y.T.; data curation, T.K. (Taishu Kawada), K.K., M.M. and Y.O.; writing—original draft preparation, T.K. (Taishu Kawada) and K.K.; writing—review and editing, T.K. (Taishu Kawada), K.K., M.M., Y.O., T.K. (Takanobu Kobayashi) and Y.T.; visualization, T.K. (Taishu Kawada), M.M. and Y.O.; supervision, K.K.; project administration, K.K.; funding acquisition, K.K. All authors have read and agreed to the published version of the manuscript.

**Funding:** This work was supported by research grants from the 114Bank Foundation.

**Data Availability Statement:** Not applicable.

**Acknowledgments:** Thank you for the administrative support of Tokushima Bunri University.

**Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

## References

1. Silva, E.; Edwards, A.M. (Eds.) *Flavins: Photochemistry and Photobiology*; RSC Publishing: Cambridge, UK, 2006.
2. Ravanat, J.-L.; Saint-Pierre, C.; Cadet, J. One-electron oxidation of the guanine moiety of 2'-deoxyguanosine: Influence of 8-oxo-7,8-dihydro-2'-deoxyguanosine. *J. Am. Chem. Soc.* **2003**, *125*, 2030–2031. [[CrossRef](#)] [[PubMed](#)]
3. Morikawa, M.; Kino, K.; Oyoshi, T.; Suzuki, M.; Kobayashi, T.; Miyazawa, H. Product analysis of photooxidation in isolated quadruplex DNA; 8-oxo-7,8-dihydroguanine and its oxidation product at 3'-G are formed instead of 2,5-diamino-4 H-imidazol-4-one. *RSC Adv.* **2013**, *3*, 25694–25697. [[CrossRef](#)]
4. Ikeda, H.; Saito, I. 8-Methoxydeoxyguanosine as an effective precursor of 2-aminoimidazolone, a major guanine oxidation product in one-electron oxidation of DNA. *J. Am. Chem. Soc.* **1999**, *121*, 10836–10837. [[CrossRef](#)]
5. Ming, X.; Matter, B.; Song, M.; Veliath, E.; Shanley, R.; Jones, R.; Tretyakova, N. Mapping structurally defined guanine oxidation products along DNA duplexes: Influence of local sequence context and endogenous cytosine methylation. *J. Am. Chem. Soc.* **2014**, *136*, 4223–4235. [[CrossRef](#)]
6. Luo, W.; Muller, J.G.; Burrows, C.J. The pH-dependent role of superoxide in riboflavin-catalyzed photooxidation of 8-oxo-7,8-dihydroguanosine. *Org. Lett.* **2001**, *3*, 2801–2804. [[CrossRef](#)] [[PubMed](#)]

7. Neeley, W.L.; Delaney, J.C.; Henderson, P.T.; Essigmann, J.M. In vivo bypass efficiencies and mutational signatures of the guanine oxidation products 2-aminoimidazolone and 5-guanidino-4-nitroimidazole. *J. Biol. Chem.* **2004**, *279*, 43568–43573. [[CrossRef](#)] [[PubMed](#)]
8. Suzuki, M.; Kino, K.; Kawada, T.; Morikawa, M.; Kobayashi, T.; Miyazawa, H. Analysis of nucleotide insertion opposite 2,2,4-triamino-5(2H)-oxazolone by eukaryotic B- and Y-Family DNA polymerases. *Chem. Res. Toxicol.* **2015**, *28*, 1307–1316. [[CrossRef](#)]
9. Stathis, D.; Lischke, U.; Koch, S.C.; Deiml, C.A.; Carell, T. Discovery and mutagenicity of a guanidinoformimine lesion as a new intermediate of the oxidative deoxyguanosine degradation pathway. *J. Am. Chem. Soc.* **2012**, *134*, 4925–4930. [[CrossRef](#)]
10. Fall, H.H.; Petering, H.G. Metabolite Inhibitors. I. 6,7-Dimethyl-9-formylmethylisoalloxazine, 6,7-dimethyl-9-(2'-hydroxyethyl)-isoalloxazine and derivatives. *J. Am. Chem. Soc.* **1956**, *78*, 377–380. [[CrossRef](#)]
11. Sanjust, E.; Cocco, D.; Curreli, N.; Rescigno, A.; Sollai, F.; Bannister, J.V. Flavin-grafted poly(vinyl alcohol): Preparation and properties. *J. Appl. Polym. Sci.* **2002**, *85*, 2471–2477. [[CrossRef](#)]
12. Singh, R.; Geetanjali; Chauhan, S.M.S. Electron transfer in natural and unnatural flavoporphyrins. *Bioorg. Chem.* **2004**, *32*, 140–169. [[CrossRef](#)] [[PubMed](#)]
13. Kino, K.; Miyazawa, H.; Sugiyama, H. User-friendly synthesis and photoirradiation of a flavin-linked oligomer. *Genes Environ.* **2007**, *29*, 23–28. [[CrossRef](#)]
14. Svoboda, J.; König, B.; Sadeghian, K.; Schützb, M. 2'-Oxoethyl flavin revisited. *Z. Für Nat. B.* **2008**, *63*, 47–54. [[CrossRef](#)]
15. Morikawa, M.; Kino, K.; Senda, T.; Suzuki, M.; Kobayashi, T.; Miyazawa, H. Formation of a flavin-linked peptide. *Molecules* **2014**, *19*, 9552–9561. [[CrossRef](#)]
16. Morikawa, M.; Kino, K.; Asada, E.; Katagiri, K.; Mori-Yasumoto, K.; Suzuki, M.; Kobayashi, T.; Miyazawa, H. N'-[2-(7,8-Dimethyl-2,4-dioxo-3,4-dihydrobenzo[g]pteridin-10(2H)-yl)ethylidene]-4-nitrobenzohydrazide. *Molbank* **2014**, *2014*, M836. [[CrossRef](#)]
17. Chen, D.; Huang, Y.; Jiang, H.; Yassen, W.; Guo, D.; Su, Y.; Xue, B.; Jin, X.; Zhu, X. Fabrication of activity-reporting glucose oxidase nanocapsules with oxygen-independent fluorescence variation. *ACS Appl. Mater. Interfaces* **2018**, *10*, 26005–26015. [[CrossRef](#)]
18. Vicens, Q.; Mondragón, E.; Reyes, F.E.; Coish, P.; Aristoff, P.; Berman, J.; Kaur, H.; Kells, K.W.; Wickens, P.; Wilson, J.; et al. Structure–activity relationship of flavin analogues that target the flavin mononucleotide riboswitch. *ACS Chem. Biol.* **2018**, *13*, 2908–2919. [[CrossRef](#)]
19. Tagami, T.; Arakawa, Y.; Minagawa, K.; Imada, Y. Efficient use of photons in photoredox/enamine dual catalysis with a peptide-bridged flavin–amine hybrid. *Org. Lett.* **2019**, *21*, 6978–6982. [[CrossRef](#)]
20. Leonard, N.J.; Lambert, R.F. Synthetic spectroscopic models related to coenzymes and base pairs. VI. Synthesis of ( $\omega$ -arylalkyl)-6,7-dimethylisoalloxazines, spectroscopic model compounds related to flavin-adenine dinucleotide. *J. Org. Chem.* **1969**, *34*, 3240–3248. [[CrossRef](#)]
21. Walter, A.; Storch, G. Synthetic C6-functionalized aminoflavin catalysts enable aerobic bromination of oxidation-prone substrates. *Angew. Chem. Int. Ed.* **2020**, *59*, 22505–22509. [[CrossRef](#)]
22. Imada, Y.; Ohno, T.; Naota, T. Synthesis and electrochemical behavior of clothespin-shaped bisflavin compounds. *Tetrahedron Lett.* **2008**, *49*, 2523–2526. [[CrossRef](#)]
23. Murahashi, S.; Ono, S.; Imada, Y. Asymmetric Baeyer-Villiger reaction with hydrogen peroxide catalyzed by a novel planar-chiral bisflavin. *Angew. Chem. Int. Ed.* **2002**, *41*, 2366–2368. [[CrossRef](#)]
24. Yano, Y.; Ohya, E.; Kuwabara, Y. Association and reactivity of amphiphilic flavins. 8-Alkyl mercaptolavins and the corresponding bis-flavins in aqueous solution. *Chem. Lett.* **1984**, *13*, 1009–1012. [[CrossRef](#)]
25. Zipplies, M.F.; Staab, H.A.  $\pi \dots \pi$ -interactions of flavins.-II. [3.3](3,10)Isoalloxazinophane and quinoxaline-like flavin interactions. *Tetrahedron Lett.* **1984**, *25*, 1035–1038. [[CrossRef](#)]
26. Barthel, A.; Trieschmann, L.; Ströhl, D.; Kluge, R.; Böhm, G.; Csuk, R. Synthesis of dimeric quinazolin-2-one, 1,4-benzodiazepin-2-one, and isoalloxazine compounds as inhibitors of amyloid peptides association. *Arch. Pharm. Chem. Life Sci.* **2009**, *342*, 445–452. [[CrossRef](#)] [[PubMed](#)]