

Communication

Synthesis of Catena-bis(μ -bromo)-(O-methyl-N-phenylthiocarbamate)-dicopper(I) and Its Reactivity towards PAr_3 (Ar = Ph, *p*-Tol)

Wafa Arar^{1,2}, Lydie Viau^{2,*} , Isabelle Jourdain^{2,*} , Michael Knorr^{2,*} , Carsten Strohmann³ , Rebecca Scheel³ and Azaiez Ben Akacha¹

¹ Selective Organic-Heterocyclic Synthesis-Biological Activity Evaluation (LR17ESO01), Université de Tunis-El Manar, Tunis 2092, Tunisia; wafaarar22@gmail.com (W.A.); azaiez.benakacha@fst.utm.tn (A.B.A.)

² Institut UTINAM UMR 6213 CNRS, Université de Franche-Comté, 16, Route de Gray, 25030 Besançon, France; lydie.viau@univ-fcomte.fr

³ Anorganische Chemie, Technische Universität Dortmund, Otto-Hahn Straße 6, 44227 Dortmund, Germany; carsten.strohmman@tu-dortmund.de (C.S.); rebecca.scheel@tu-dortmund.de (R.S.)

* Correspondence: isabelle.jourdain@univ-fcomte.fr (I.J.); michael.knorr@univ-fcomte.fr (M.K.); Tel.: +33-3-81-66-62-70 (M.K.)

Abstract: In order to investigate the coordination chemistry of *O*-alkyl *N*-aryl thiocarbamate ligands toward coinage metals, CuBr was reacted with one equivalent of MeOC(=S)N(H)Ph **L** in MeCN solution to afford the 1D-polymeric title compound $[\{\text{Cu}(\mu_2\text{-Br})_2\text{Cu}\}\{\mu_2\text{-MeOC(=S)N(H)Ph}\}_2]_n$ **CP1**. Compound **1** was characterized by IR spectroscopy and an elemental analysis. The formation of a polymeric 1D ribbon built upon μ_2 -bridging bromido and thione ligands via the C=S bond was ascertained by a single-crystal X-ray diffraction study performed at 100 K. In the presence of PAr_3 (Ar = Ph, *p*-Tol), the polymer chain was broken to yield the mononuclear complexes $[(\text{Ar}_3\text{P})_2\text{Cu}\{\text{MeOC(=S)N(H)Ph}\}\text{Br}]$ **C1** and **C2**.

Keywords: copper bromide; thiocarbamate; thione; crystal structure; hydrogen bonding; aryl phosphine



Citation: Arar, W.; Viau, L.; Jourdain, I.; Knorr, M.; Strohmman, C.; Scheel, R.; Ben Akacha, A. Synthesis of Catena-bis(μ -bromo)-(O-methyl-N-phenylthiocarbamate)-dicopper(I) and Its Reactivity towards PAr_3 (Ar = Ph, *p*-Tol). *Molbank* **2023**, *2023*, M1655. <https://doi.org/10.3390/M1655>

Academic Editor: Rodrigo Abonia

Received: 25 April 2023

Revised: 25 May 2023

Accepted: 29 May 2023

Published: 31 May 2023




Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Thiocarbamates, also called thiocarbamides, with the general formula ROC(=S)N(H)R' play not only a role in organic chemistry as building blocks for further transformations but also in applications in various domains, such as biological and pharmacological activities or catalysis [1–5]. Furthermore, they have attracted the attention of coordination chemists, since the soft C=S thione function (according to Pearson's HSAB principle) [6] readily coordinates to a wide range of transition metal complexes, forming complexes with Cu(I), Ag(I), Au(I), Hg(II), Ru(II), Rh(III), etc. [7–11]. Furthermore, they may act as potential ditopic ligands capable of coordinating to a metal center both via their soft C=S function or through the harder HNR_2 group as demonstrated by the crystal structure determination of the square-planar bis(O-methylphenylthiocarbamato)(triphenylphosphine)-palladium [11]. We have recently demonstrated that *O*-methyl-*N*-phenyl thiocarbamate **L** can even be coordinated to *p*-block metal salts, such as BiI_3 [12]. Some representative examples of crystallographically characterized mono- and dinuclear complexes ligated by *O*-methyl-*N*-phenyl thiocarbamate are presented in Scheme 1 [7,8,12].

With the aim to investigate putative arene-C–H $\cdots\pi$ (quasi-chelate ring) interactions in copper(I) crystal structures, Tiekink et al. reported in 2014 on the synthesis of the complex $[(\text{Ph}_3\text{P})_2\text{Cu}\{\text{MeOC(=S)N(H)Ph}\}\text{Cl}]$ by the reaction of CuCl with **L** in the presence of two equivalents of PPh_3 [7], shown as complex **A** in Scheme 1. In a continuation of our investigations on the coordination chemistry of thione-type ligands on diverse metal centers [13–16], our group extended the preparation of this type of Cu(I) complex


to CuBr and CuI, systematically analyzed by crystal structure determinations a wide series of complexes $[(\text{Ar}_3\text{P})_2\text{Cu}\{\text{ROC}(=\text{S})\text{N}(\text{H})\text{Ph}\}\text{X}]$ ($\text{Ar} = \text{Ph}$, $p\text{-C}_6\text{H}_4\text{OMe}$; $\text{X} = \text{Br}$, I), and investigated the biological activity of these compounds featuring intramolecular N-H...Hal bonding, giving rise to six-membered cycles [17]. We also reported that the formation of $[(\text{Ar}_3\text{P})_2\text{Cu}\{\text{ROC}(=\text{S})\text{N}(\text{H})\text{Ph}\}\text{I}]$ is preceded in the initial phase of the reaction by the formation of mono-dimensional coordination polymers of type $[\{\text{Cu}(\mu_2\text{-I})_2\text{Cu}\}(\mu_2\text{-L})_2]_n$ [17]. We now investigate in more detail the outcome of the reaction of CuBr with **L**, describing herein the synthesis and crystal structure of the resulting coordination polymer $[\{\text{Cu}(\mu_2\text{-Br})_2\text{Cu}\}\{\mu_2\text{-MeOC}(=\text{S})\text{N}(\text{H})\text{Ph}\}_2]_n$ **CP1** and reporting on its reactivity towards PAr_3 .



Scheme 1. Examples of some *O*-methyl *N*-arylthiocarbamate complexes.

2. Results and Discussion

The hitherto unknown coordination polymer $[\{\text{Cu}(\mu_2\text{-Br})_2\text{Cu}\}\{\mu_2\text{-MeOC}(=\text{S})\text{N}(\text{H})\text{Ph}\}_2]_n$ **CP1** was obtained by the addition of one equivalent of *O*-methyl *N*-phenyl thiocarbamate **L** to a solution of CuBr in MeCN (Scheme 2). After stirring for 20 min, the precipitation of a colorless microcrystalline material occurred, which upon heating for 5 min almost redissolved. Upon cooling, colorless crystals started to grow. According to the elemental analysis of this air-stable material, one **L** molecule is attached per CuBr motif. This polymeric material, whose characterization is described below, reacts straightforwardly upon the addition of two equivalents of PPh_3 per CuBr unit to form by the splitting of the $\mu_2\text{-Br}$ bridges of **CP1** the mononuclear tetrahedral complex $[(\text{Ph}_3\text{P})_2\text{Cu}\{\text{MeOC}(=\text{S})\text{N}(\text{H})\text{Ph}\}\text{Br}]$ **C1** (Scheme 2). Since this complex has been recently fully characterized, we refer for its spectroscopic data to ref. [17]. In an analogous manner, **CP1** reacts with tris(*p*-tolyl)phosphine PTol_3 to yield the derivative $[(\text{Tol}_3\text{P})_2\text{Cu}\{\text{MeOC}(=\text{S})\text{N}(\text{H})\text{Ph}\}\text{Br}]$ **C2** with over an 85% yield as an air-stable crystalline solid. Its ^{31}P NMR spectrum in CDCl_3 shows a single broadened resonance at $\delta -6.34$ with a $\Delta\delta$ of 0.92 ppm with respect to free PTol_3 [18]. The ^1H NMR spectrum displays a strong singlet at $\delta 2.33$ ppm, attributed to the six equivalent methyl groups of the *p*-tolyl rings. The characteristic signals of the thiocarbamate moiety at $\delta 4.06$ and 10.54 ppm are assigned to $-\text{OCH}$ and the NH group while the aromatic protons are observed as a multiplet between $\delta 7.03$ and 7.39 ppm (Figure S2). In the proton-decoupled ^{13}C NMR spectrum, a single resonance at $\delta 21.4$ ppm is seen for the six magnetically equivalent Me substituents. The C-atom of the ligated thione function resonates at $\delta 187.3$ ppm (Figure S3). The ATR-IR spectrum features a broadened N-H vibration at 3160 cm^{-1} , indicative of an intramolecular N-H...Br interaction, as was crystallographically evidenced for **C1** (Figure S4). Upon the treatment of **CP1** with an excess of PPh_3 , the dissociation of the coordinated thione ligand occurs, yielding finally, among other non-identified products, the cluster compounds $[(\text{PPh}_3)_2\text{Cu}(\mu_2\text{-Br})_2\text{CuPPh}_3]$ and $[\text{Cu}_4\text{Br}_4(\text{PPh}_3)_4]$, which are already known in the literature, as evidenced by X-ray diffraction [19].



Scheme 2. Synthesis of the title compound **CP1** and its reactivity with PAr_3 yielding **C1** and **C2**.

Since the compound **CP1** is not soluble enough to be characterized by NMR spectroscopy, we examined the material crystallizing in the triclinic space group $P\bar{1}$ by an X-ray diffraction study performed at 100 K. As shown in Figure 1, a mono-dimensional ribbon has formed, which is constituted of rhomboid-shaped dinuclear $\{\text{Cu}(\mu_2\text{-Br})_2\text{Cu}\}$ units that are interconnected through the bridging $\text{C}=\text{S}$ groups of the **L** molecules. Each Cu(I) center of **CP1** has thus a tetrahedral coordination sphere (neglecting the $\text{Cu} \cdots \text{Cu}$ interaction). The Cu–Cu contact within the $\{\text{Cu}(\mu_2\text{-Br})_2\text{Cu}\}$ rhomboid of 2.8339(10) Å falls within the limit of the Van der Waals radii of two Cu atoms (2.8 Å) and can be considered as weakly bonding. However, without computation, it remains speculative to discuss the contribution of metallophilic interactions, which can occur both in molecular compounds or coordination polymers incorporating neighboring coinage metal centers [20–22].

Similar values have been reported for the 1D CPs $[\{\text{Cu}(\mu_2\text{-Br})_2\text{Cu}\}\{\mu_2\text{-PhC(=S)NH}_2\}]_n$ and $[\{\text{Cu}(\mu_2\text{-Br})_2\text{Cu}\}(\mu\text{-PhS(CH}_2)_3\text{SPh})_2]_n$, in which the metal centers within the rhomboid clusters are separated by 2.857(3) and 2.7851(12) Å, respectively [23,24]. For the molecular compounds $[\{\text{Cu}(\mu_2\text{-Br})_2\}(1\text{-oxa-4,7-dithiacyclononane})_2]$ (2.852(1) Å), $[\{\text{Cu}(\mu_2\text{-Br})_2\}(\text{dibenzo}[e,m][1,4,8,11]\text{dioxadithia-cyclotetradecine-S,S'})_2]$ (2.919(7) Å), and the supramolecular compound tetrakis(4,5-bis(methylsulfanyl)-1,3-dithiole-2-thione)-bis(μ_2 -bromo)dicopper (2.893(1) Å), the $\text{Cu} \cdots \text{Cu}$ interaction lies in a similar range [16,25,26]. A further example is the thione-bridged dinuclear complex $[\text{Cu}_2\text{Br}_2(\mu\text{-S-mimzSH})_2(\eta^1\text{-S-mimzSH})_2]$ (2.877(6) Å), obtained by the reaction of CuBr with 1-methyl-1,3-imidazoline-2-thione (mimzSH) in a 1:2 molar ratio [27].

This is in contrast with the recently described iodo analog $[\{\text{Cu}(\mu_2\text{-I})_2\text{Cu}\}\{\mu_2\text{-MeOC(=S)N(H)Ph}\}_2]_n$ **CP2**, for which a much shorter Cu–Cu separation of only 2.6563(7) Å was evidenced [17]. The $\text{C}=\text{S}$ bond is slightly elongated with respect to that of non-ligated MeOC(=S)N(H)Ph (1.711(5) vs. 1.6708(11) Å) [28].




Figure 1. Crystal structure of the 1D ribbon of $[\{\text{Cu}(\mu_2\text{-Br})_2\text{Cu}\}(\mu_2\text{-L})_2]_n$ **CP1** running along the *a* axis. Selected bond lengths (Å) and angles (deg) of **CP1**: Cu \cdots Cu¹ 2.8649(10), Cu–Cu² 2.8339(10), Br–Cu 2.4620(7), Cu–S 2.2966(12), Cu–S² 2.4157(12), S–C1 1.711(5), O–C1 1.313(5), O–C2 1.453(5), N–C1 1.335(5), N–C3 1.421(6); Cu²–Cu–Cu¹ 171.28(5), S–Cu–Br 117.68(4), S–Cu–Br¹ 107.64(4), S–Cu–S 106.11(4), Br–Cu–Br¹ 109.12(2), Br–Cu–Cu¹ 54.83(2), Br–Cu–Cu² 130.17(3), Br¹–Cu–Cu² 120.12(3), S–Cu–Cu¹ 131.40(4), S–Cu–Cu² 51.13(3), S–Cu–Cu¹ 131.30(4), N–C1–S 121.7(3), C8–C3–N 115.3(4), and C7–C8–C3 120.4(3). Symmetry transformation used to generate equivalent atoms: ¹ 2-*x*, 1-*y*, 2-*z*; ² 1-*x*, 1-*y*, 2-*z*.

Noteworthy is also the occurrence of the secondary N-H \cdots Br interaction of 2.62(6) Å (*d*N \cdots Br 3.440(4) Å, indicated by the red dashed lines in Figure 1. This interaction is noticeably shorter and in consequence stronger than the weak intramolecular N-H \cdots I contacts of 2.748 Å and 2.803 Å encountered for **CP2** [17]. A similar systematic shortening of the intramolecular N-H \cdots X bonding occurring in $[(\text{Ph}_3\text{P})_2\text{Cu}\{\text{MeOC}(=\text{S})\text{N}(\text{H})\text{Ph}\}\text{X}]$ (X = I, Br, Cl) and giving rise to six-member cycles has also been evidenced recently by us (see Scheme 1) [17]. Note that in the latter case, this intramolecular hydrogen bonding is even stronger for the molecular bromido complex **C1** compared to **CP1** (2.51(2) vs. 2.62(6)). Furthermore, a loose intramolecular S \cdots H contact of 2.672 Å occurs between the sulfur atom and the H atom of the NH group.

In the packing, the ribbons of **CP1** are aligned in a co-parallel arrangement and are running along the *a* axis (Figure 2). There are also weak supramolecular interchain contacts between Br and the H atom of C7 of the aryl cycle (C7-H7 \cdots Br 2.919 Å).

To check the phase purity of this material, a grounded sample has been analyzed by powder X-ray diffraction (PXRD). The experimental PXRD pattern matches well with the simulated ones, confirming the homogeneity of the phases. A comparison of the PXRD patterns of **CP1** is illustrated in Figure 3.




Figure 2. Perspective view of the packing, showing several ribbons of CP1 running along the *a* axis in a parallel arrangement.




Figure 3. Comparison of the experimental PXRD pattern of CP1 with the simulated one (blue).

3. Experimental Section

Synthesis of CP1—We added 1 eq. of *O*-methyl *N*-phenylthiocarbamate **L** (334 mg, 2 mmol) to a solution of CuBr (286 mg, 2 mmol) in 12 mL of acetonitrile. A colorless precipitation was immediately noticed. The mixture was stirred 20 min at room temperature and then shortly heated till almost all of the precipitates had redissolved. After 1 d, pale yellowish crystals of CP1 were formed and then collected by filtration. Yield: 79%. Anal. Calc. for C₈H₉BrCuNOS (M.W = 310.67 g·mol^{−1}). C, 33.09; H, 4.01; N, 4.29; S 9.81%. Found: C, 32.95; H, 4.00; N, 4.15; S 9.75. IR-ATR: 1026 (C–O), 1215 (C=S), 1440 ν(C–N), 3199 ν(N–H···Br), 3276 ν(N–H) cm^{−1}.

Synthesis of C1—We added two equivalents of PPh₃ (52 mg, 0.2 mmol) to a suspension of CP1 (31 mg, 0.1 mmol) in 4 mL of MeCN, prepared as described above. The mixture

was then heated for 20 min to 60 °C till all of the product was dissolved. Upon cooling, **C1** crystallized. The characterization data are identical to those described previously [18].

Synthesis of C2—We added two equivalents of PTol₃ (608 mg, 2 mmol) in several portions to an in situ prepared suspension of **CP1** (1 mmol). The mixture was then heated for 20 min to 60 °C till all product was dissolved. Upon cooling, **C2** crystallized in form of colorless plates and was isolated with 82% yield. Anal. Calc. for C₅₀H₅₁CuBrNOP₂S (M.W. = 919.43 g·mol^{−1}): C, 65.32; H, 5.59; N, 1.52; S, 3.49%. Found C, 65.20; H, 5.53; N, 1.59; S, 3.42%. ¹H NMR: δ 2.33 (s, 18H, CH₃), 4.08 (s, 3H, OCH₃), 7.04 (d, 12H, PC₆H₄, ³J_{HH} = 7.9 Hz), 7.15–7.39 (m, 17H, Ph and PC₆H₄, ³J_{HH} = 7.9, ³J_{PH} = 9.7 Hz), 10.54 (s, br, 1H, NH). ¹³C{¹H} NMR: δ 21.4 (CH₃Ar), 58.4 (CH₃O), 121.9 (CH, Ph, *Cmeta*), 125.3 (CH, Ph, *Cpara*), 128.9 (CH, Ph, *Cortho*), 129.2 (d, CH, *m*-PC₆H₄, ³J_{CP} = 9.2 Hz), 130.4 (d, C_q, *Cipso*PC₆H₄, ¹J_{CP} = 27.7 Hz), 133.9 (d, CH, *o*-PC₆H₄, ²J_{CP} = 14.9 Hz), 137.2 (s, C_q, *Cipso*), 139.4 (C_q, *p*-PC₆H₄), 187.3 (C_q, C=S). ³¹P{¹H}: δ −6.34. IR-ATR: 1093 ν(C–O), 1222 ν(C=S), 1444 ν(C–N), 3160, 3110 ν(N–H) cm^{−1}.

Crystal data for C₈H₉BrCuNOS were as follows: M = 310.67 g·mol^{−1}, pale yellow crystals, crystal size 0.362 × 0.102 × 0.047 mm³, Triclinic, space group *P* $\bar{1}$, *a* = 5.6823(2) Å, *b* = 7.6782(4) Å, *c* = 11.4019(5) Å, α = 97.689(2)°, β = 91.877(2)°, γ = 91.143(2)°, *V* = 492.58(4) Å³, *Z* = 2, *D*_{calc} = 2.095 g/cm³, *T* = 100 K, *R*₁ = 0.0475, *R*_w = 0.0976 for 5163 reflections with *I* ≥ 2σ(*I*) and 2995 independent reflections. Largest diff. peak/hole e/Å^{−3} 1.04/−1.34. Data were collected on Bruker D8 Venture four-circle diffractometer equipped with a PHOTON II CPAD detector by Bruker AXS GmbH, using graphite-monochromated MoK_α radiation *l* = 0.71073 Å and were deposited at the Cambridge Crystallographic Data Centre as CCDC 2252606 (Supplementary Materials). The data can be obtained free of charge from the Cambridge Crystallographic Data Centre via <http://www.ccdc.cam.ac.uk/getstructures>, accessed on 3 February 2023. The structure was solved by direct methods and refined by full-matrix least-squares calculation against *F*² (SHELXL, 2015) [29–31].

4. Conclusions

We have crystallographically evidenced that CuBr reacts like CuI with a stoichiometric amount of *O*-methyl-*N*-phenylthiocarbamate to yield a halide-bridged mono-dimensional coordination polymer. We are currently extending this facile preparation of polymeric materials to other *O*-alkyl *N*-arylthiocarbamates, such as MeOC(=S)N(H)Cy, and are investigating the splitting of the polymer chain by other P and As donor ligands.

Supplementary Materials: The following supporting information can be downloaded online: CIF file, Check-CIF report, and IR and NMR spectra.

Author Contributions: W.A. and M.K. prepared the compounds; C.S., R.S. and L.V. collected the X-ray data and solved the structure; I.J., L.V. and M.K. designed the study and analyzed the data and M.K. wrote the paper. A.B.A., I.J. and M.K. contributed with the conceptualization. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: The X-ray data were deposited at the CCDC as stated in the paper.

Acknowledgments: We thank Stéphanie Befly for recording the IR and NMR spectra.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Németh, A.G.; Keserű, G.M.; Ábrányi-Balogh, P. A novel three-component reaction between isocyanides, alcohols or thiols and elemental sulfur: A mild, catalyst-free approach towards *O*-thiocarbamates and dithiocarbamates. *Beilstein J. Org. Chem.* **2019**, *15*, 1523–1533. [PubMed]
2. Zhang, J.; Zang, Q.; Yang, F.; Zhang, H.; Sun, J.Z.; Tang, B.Z. Sulfur Conversion to Multifunctional Poly(*O*-thiocarbamate)s through Multicomponent Polymerizations of Sulfur, Diols, and Diisocyanides. *J. Am. Chem. Soc.* **2021**, *143*, 3944–3950. [CrossRef]

3. Shadap, L.; Tyagi, J.L.; Poluri, K.M.; Novikov, S.; Timothy Lo, C.W.; Mozharivskyj, Y.; Kollipara, M.R. Insights to the strained thiocarbamate derivative complexes of platinum group metals induced by azide as a co-ligand: Characterization and biological studies. *J. Organomet. Chem.* **2020**, *920*, 121345. [\[CrossRef\]](#)
4. Krátky, M.; Volková, M.; Novotná, E.; Trejtnar, F.; Stolaříková, J.; Vinšová, J. Synthesis and biological activity of new salicylanilide N,N-disubstituted carbamates and thiocarbamates. *Bioorg. Med. Chem.* **2014**, *22*, 4073–4082. [\[CrossRef\]](#) [\[PubMed\]](#)
5. Tan, C.K.; Le, C.; Yeung, Y.Y. Enantioselective bromolactonization of cis-1,2-disubstituted olefinic acids using an amino-thiocarbamate catalyst. *Chem. Commun.* **2012**, *48*, 5793–5795. [\[CrossRef\]](#)
6. Pearson, R.G. Recent advances in the concept of hard and soft acids and bases. *J. Chem. Edu.* **1987**, *64*, 561–567. [\[CrossRef\]](#)
7. Yeo, C.I.; Halim, S.N.A.; Weng Ng, S.; Tan, S.L.; Zukerman-Schpector, J.; Ferreira, M.A.B.; Tiekink, E.R.T. Investigation of putative arene-C–H... π (quasi-chelate ring) interactions in copper(I) crystal structures. *Chem. Commun.* **2014**, *50*, 5984–5986.
8. Yeo, C.I.; Teow, S.Y.; Liew, L.Y.; Chew, J.; Tiekink, E.R.T. Crystal structure of chlorido-(O-methyl phenylcarbamothioamide- κ S)-bis(triphenylphosphane- κ P)silver(I), C₄₄H₃₉AgClNOP₂S. *Z. Kristallogr. NCS* **2020**, *235*, 1473–1475. [\[CrossRef\]](#)
9. Hall, V.J.; Siasios, G.; Tiekink, E.R.T. Trioganophosphinegold(I)carbonimidothioates. *Aust. J. Chem.* **1993**, *46*, 561–570. [\[CrossRef\]](#)
10. Bandoli, G.; Clemente, D.A.; Sindellari, L.; Tondello, E. Preparation, properties, and crystal structure of dichlorobis-(O-ethyl thiocarbamate)mercury(II). *J. Chem. Soc. Dalton Trans.* **1975**, 449–452. [\[CrossRef\]](#)
11. Furlani, C.; Tarantelli, T.; Gastaldi, L.; Porta, P. Complexing behaviour of thiocarbamic esters: Crystal and molecular structure of bis-(O-methylphenylthiocarbamate) (triphenylphosphine)-palladium(II). *J. Chem. Soc. A* **1971**, 3778–3783. [\[CrossRef\]](#)
12. Arar, W.; Khatyr, A.; Knorr, M.; Strohmman, C.; Schmidt, A. Bis(μ -iodo)-tetrakis(O-methyl N-phenylthiocarbamate)-tetraiodo-dibismuth. *Molbank* **2022**, *2022*, M1381. [\[CrossRef\]](#)
13. Arar, W.; Khatyr, A.; Knorr, M.; Brieger, L.; Krupp, A.; Strohmman, C.; Efrat, M.L.; Ben Akacha, A. Synthesis, crystal Structures and hirshfeld analyses of phosphonothioamidates (EtO)2P(=O)C(=S)N(H)R (R = Cy, Bz) and their coordination on CuI and HgX₂ (X = Br, I). *Phosphorus Sulfur. Silicon Relat. Elem.* **2021**, *196*, 845–858. [\[CrossRef\]](#)
14. Hameau, A.; Guyon, F.; Knorr, M.; Enescu, M.; Strohmman, C. Self-Assembly of Dithiolene-based Coordination Polymers of Mercury(II): Dithioether versus Thiocarbonyl Bonding. *Monatsh. Chem.* **2006**, *137*, 545–555. [\[CrossRef\]](#)
15. Guyon, F.; Hameau, A.; Khatyr, A.; Knorr, M.; Amrouche, H.; Fortin, D.; Harvey, P.D.; Strohmman, C.; Ndiaye, A.L.; Huch, V.; et al. Syntheses, Structures, and Photophysical Properties of Mono- and Dinuclear Sulfur-Rich Gold(I) Complexes. *Inorg. Chem.* **2008**, *47*, 7483–7492. [\[CrossRef\]](#)
16. Hameau, A.; Guyon, F.; Khatyr, A.; Knorr, M.; Strohmman, C. 4,5-Bis(methylthio)-1,3-dithiole-2-thione, a versatile sulphur-rich building block for the self-assembly of Cu(I) and Ag(I) coordination polymers: Dithioether versus thiocarbonyl bonding. *Inorg. Chim. Acta* **2012**, *388*, 60–70. [\[CrossRef\]](#)
17. Arar, W.; Ben Ali, R.; El May, M.V.; Khatyr, A.; Jourdain, I.; Knorr, M.; Brieger, L.; Scheel, R.; Strohmman, C.; Chaker, A.; et al. Synthesis, crystal structures and biological activities of halogeno-(O-alkylphenylcarbamothioate)bis(triarylphosphine)copper (I) complexes. *J. Mol. Struct.* **2023**, *1284*, 135370. [\[CrossRef\]](#)
18. Chou, W.N.; Pomerantz, M. N-phenyl-P,P,P-triarylphospho- λ 5-azenes, triarylphosphines, and triarylphosphine oxides. Substituent effects on nitrogen-15, phosphorus-31, and carbon-13 NMR spectra. *J. Org. Chem.* **1991**, *56*, 2762–2769. [\[CrossRef\]](#)
19. Hisao, N.; Morio, H.; Yoshihiko; Masahiroj, K.; Tsumoto, O. The Crystal Structure of Di- μ -Bromo-tris(triphenylphosphine) dicopper(II). *Bull. Soc. Jpn.* **1981**, *54*, 1247–1248. [\[CrossRef\]](#)
20. Moussa, J.; Chamoreau, L.M.; Gullo, M.P.; Esposti, A.D.; Barbieri, A.; Amouri, H. Induced phosphorescence from Pt \rightarrow Ag and Ag(I) \cdots Ag(I) metallophilic interactions in benzenedithiolatodiimine-Pt₂/Ag₂ clusters: A combined experimental and theoretical investigation. *Dalton Trans.* **2016**, *45*, 2906–2913. [\[CrossRef\]](#)
21. Benard, M.; Bodensieck, U.; Braunstein, P.; Knorr, M.; Strampfer, M.; Strohmman, C. Conformation Control in Polymetallic Mesocycles by Metal–Metal Bonding: The First Example of an Hg–Cu Interaction. *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 2758–2761.
22. Harisomayajula, N.S.; Makovetskyi, S.; Tsai, Y.C. Cuprophilic Interactions in and between Molecular Entities. *Chem. Eur. J.* **2019**, *25*, 8936–8954. [\[CrossRef\]](#) [\[PubMed\]](#)
23. Troyano, J.; Zapata, E.; Perles, J.; Amo-Ochoa, P.; Fernández-Moreira, V.; Martínez, J.I.; Zamora, F.; Delgado, S. Multifunctional Copper(I) Coordination Polymers with Aromatic Mono- and Ditopic Thioamides. *Inorg. Chem.* **2019**, *58*, 3290–3301. [\[CrossRef\]](#) [\[PubMed\]](#)
24. Knorr, M.; Guyon, F.; Khatyr, A.; Strohmman, C.; Allain, M.; Aly, S.M.; Lapprand, A.; Fortin, D.; Harvey, P.D. Construction of (CuX)_{2n} Cluster-Containing (X = Br, I; n = 1, 2) Coordination Polymers Assembled by Dithioethers ArS(CH₂)_mSAr (Ar = Ph, p-Tol; m = 3, 5): Effect of the Spacer Length, Aryl Group, and Metal-to-Ligand Ratio on the Dimensionality, Cluster Nuclearity, and the Luminescence Properties of the Metal–Organic Frameworks. *Inorg. Chem.* **2012**, *18*, 9917–9934.
25. Lee, S.Y.; Park, S.; Lee, S.S. Copper(I) Silver(I), and Palladium(II) Complexes of a Thioxamacrocyclic Displaying Unusual Topologies. *Inorg. Chem.* **2009**, *48*, 11335–11341. [\[CrossRef\]](#)
26. Lucas, C.R.; Liang, W.; Miller, D.O.; Bridson, J.N. Metal Complexes of 1-Oxa-4,7-dithiacyclononane. *Inorg. Chem.* **1997**, *36*, 4508–4513. [\[CrossRef\]](#)
27. Lobana, T.S.; Sharma, R.; Sharma, R.; Butcher, R.J. Metal Derivatives of Heterocyclic Thioamides: Synthesis and Crystal Structures of Copper Complexes with 1-Methyl-1,3-imidazoline-2-thione and 1,3-Imidazoline-2-thione. *Z. Anorg. Allg. Chem.* **2008**, *634*, 1785–1790. [\[CrossRef\]](#)
28. Ho, S.Y.; Lai, C.S.; Tiekink, E.R.T. O-Methyl N-phenylthiocarbamate. *Acta Cryst.* **2003**, *E59*, o1155–o1156.

29. Sheldrick, G. SHELXT- Integrated space-group and crystal-structure determination. *Acta Crystallogr. A* **2015**, *71*, 3–8. [[CrossRef](#)]
30. Sheldrick, G. Crystal structure refinement with SHELXL. *Acta Crystallogr. C* **2015**, *71*, 3–8. [[CrossRef](#)]
31. Dolomanov, O.V.; Bourhis, L.J.; Gildea, R.J.; Howard, J.A.; Puschmann, H. OLEX2: A complete structure solution, refinement and analysis program. *J. Appl. Cryst.* **2009**, *42*, 339–341. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.