

Article Effects of Invasive Plant Diversity on Soil Microbial Communities

Xiaoyan Wang [†], Xue Wang [†], Wei Wang ¹, Jiang Wang ^{*} and Feihai Yu ^{*}

School of Life Science, Taizhou University, Taizhou 318000, China

* Correspondence: wangjiang@tzc.edu.cn (J.W.); feihaiyu@126.com (F.Y.)

+ These authors contributed equally to this work.

Abstract: Native plant communities can be invaded by different numbers of alien plant species or by the same number of alien plant species with different levels of evenness. However, little is known about how alien invasive plant species richness and evenness affect soil microbial communities. We constructed native herbaceous plant communities invaded by exotic plants with different richness (1, 2, 4 and 8 species) and evenness (high and low) and analyzed soil physico-chemical properties and the diversity and composition of soil fungal and bacterial communities by high-throughput Illumina sequencing. Overall, the species richness and evenness of invasive plants had no significant effect on bacterial and fungal alpha diversity (OTUs, Shannon, Simpson, Chao1 and ACE) or the soil physico-chemical properties. However, invasive species richness had a significant impact on the relative abundance of the most dominant fungi, Ascomycota and *Bipolaris*, and the dominant bacteria, Actinobacteriota, which increased with increasing invasive species richness. The relative abundance of some specific invasive plants in the community. This study sheds new light on the effects of plant co-invasion on soil microbial communities, which may help us understand the underlying mechanisms of multiple alien plant invasion processes from the perspective of soil microorganisms.

Keywords: bacterial community; co-invasion; diversity effect; fungal community; invasive species evenness; invasive species richness

1. Introduction

Invasions by alien plants are a world-wide problem that has received great attention [1–4]. Alien plant invasions may not only displace native plant species and damage local flora and biodiversity but may also negatively influence many other ecosystem functions and services [5–9] in relation to the regional context [10]. Thus, there is a great need to develop measures to efficiently manage and control alien plant invasions [11,12]. To this end, we need to first assess the impacts of alien plant invasions on ecosystem functions such as soil microbial communities [13–16]. Soil microbial communities can alleviate the effects of invasive plants on local plant communities by degrading allelopathic substances [17], promoting further invasion via symbiosis with arbuscular mycorrhizal fungi or inhibiting further invasion via the accumulation of pathogens [18,19].

Previous studies have shown that alien plant invasions can have profound effects on soil microbial communities [14,20,21]. For instance, alien plant invasions can affect the soil properties [22–25], species richness and composition of soil microbial communities [26,27], thus destroying the long-term balance between native plants and soil microorganisms [15,16,27]. Some studies have shown that plant invasions can improve soil microbial diversity by providing more energy sources from secondary metabolites or a greater quantity and quality of litter [26,28–30]. Other studies have found that plant invasions reduce soil microbial diversity (e.g., AM fungi) [31–33] or have no significant effect [33,34].

One overlooked phenomenon in invasion ecology is that a native plant community can be invaded by different numbers of alien plant species [35]. Thus, while many studies have



Citation: Wang, X.; Wang, X.; Wang, W.; Wang, J.; Yu, F. Effects of Invasive Plant Diversity on Soil Microbial Communities. *Diversity* **2022**, *14*, 992. https://doi.org/10.3390/d14110992

Academic Editors: Bruce Osborne and Panayiotis Dimitrakopoulos

Received: 13 October 2022 Accepted: 14 November 2022 Published: 17 November 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). examined the impacts of alien plant invasions on soil microbial communities, the impacts of alien invasive species richness have never been considered. Recent studies found that increasing invasive plant species richness significantly increased the productivity of alien invasive plants and decreased that of native plant communities [35–38]. This suggests that increasing alien plant species can promote the invasion success of alien plants into native plant communities [35]. Since previous studies have shown that changes in both alien plant productivity and native plant productivity can impact soil microbial communities [39–42], we hypothesize that alien plant richness can influence soil microbial communities.

Another overlooked phenomenon is that a native plant community can be invaded by the same number of alien plant species with different levels of evenness [35]. As an important aspect of species diversity, evenness is often used to explain inconsistencies in species richness effects [43–46]. In several cases, evenness explained more biodiversity effects than species richness [47,48]. For instance, it has been shown that species richness had impacts on bacterial communities through the modulation of plant evenness in native grassland communities [46,49]. To our best knowledge, little is known about the effect of invasive plant evenness on soil microbial communities. We speculate that a high level of invasive plant evenness can result in high microbial diversity due to the different contributions of all invasive plants to soil micro-environment changes. By contrast, when the evenness of alien invasive plants is low, only one or two alien species are dominant, so alien plant diversity has similar effects to a monoculture of the dominant species [49,50].

We constructed native herbaceous plant communities invaded by alien plants with different richness (1, 2, 4 and 8 species) and evenness (high and low) to explore the impacts of invasive plant species diversity on soil fungal and bacterial communities. Specifically, we addressed the following questions: (1) Do alien invasive plant richness and evenness affect the diversity and composition of soil fungal communities? (2) Do alien invasive plant richness affect the diversity and composition of soil fungal communities? (3) Do alien invasive plant richness and evenness affect the abundance of some specific microbial taxa?

2. Materials and Methods

2.1. Native and Invasive Species Preparation

The plant species pool used to construct experimental plant communities consisted of 18 native plant species and 10 alien invasive plant species that commonly grow in grasslands around Taizhou city (121.43° E, 28.68° N), Zhejiang Province, China [35]. Eight of the native species were perennials (*Achyranthes bidentata* Blume, *Persicaria filiformis* (Thunb.) Nakai, *Aster trinervius* subsp. *ageratoides* (Turczaninow) Grierson, *Bellis perennis* L., *Carpesium abrotanoides* L., *Patrinia scabiosifolia* Link, *Penthorum chinense* Pursh and *Plantago asiatica* L.), and ten were annual/biennial. The ten invasive species include four perennials (*Alternanthera philoxeroides* (Mart.) Griseb., *Mirabilis jalapa* L., *Solidago canadensis* L., and *Talinum paniculatum* (Jacq.) Gaertn.) and six annuals (*Ageratum conyzoides* L., *Symphyotrichum subulatum* (Michx.) G.L.Nesom, *Bidens frondosa* L., *Bidens pilosa* L., *Celosia argentea* L., and *Sesbania cannabina* (Retz.) Poir.). Detailed information on these species was described by Wang et al. (2022) [35].

Seeds (except *Alternanthera philoxeroides*) of all native and invasive species were collected from field sites around Taizhou city and stored at 4 °C till use. In April 2020, these seeds were sown into plastic containers (30 cm long \times 30 cm wide \times 30 cm high) filled with a mixture of river sand and peat. The plastic containers were placed in a greenhouse at Taizhou University, Zhejiang Province, China. For *A. philoxeroides*, ramets (asexual individuals) were propagated from stem cuttings. One month later, seedlings with similar sizes were used to construct plant communities. Height differences within the same species were limited to 2 cm.

2.2. Experimental Design

On 27 May 2020, 96 native plant communities were constructed in 96 containers (each 120 cm in diameter \times 74 cm in height) with 18 native plant species. In each container, each native species was initially planted with three seedlings. The containers were filled with a mixture of soil (0.62 \pm 0.17 g kg⁻¹ total N, 0.13 \pm 0.03 g kg⁻¹ total P, mean \pm SE, n = 5), peat and sand at a volume ratio of 2:1:1. Slow release fertilizer (N:P:K = 14:14:14, Osmocote exact standard 3–4 M) was added to the soil at a dose of 3 g L^{-1} . After one week, individuals of invasive plants with four levels of species richness (1, 2, 4 and 8 species) and two levels of evenness (high and low) were introduced to 90 of the 96 native plant communities (Table A1). For the 1-species treatment, individuals of each of the 10 invasive species were planted in three containers, making 30 communities. For each of the 2-species, 4-species and 8-species treatments, 10 different species mixtures were constructed by randomly selecting them from the species pool in such a way that each species had an equal chance of being selected. Each of these mixtures was replicated twice: one was used for high evenness (2-species, 12:12; 4-species, 6:6:6:6; 8-species, 3:3:3:3:3:3:3) and the other for low evenness (2-species, 22:2; 4-species, 18:3:2:1; 8-species, 13:3:2:2:1:1:1:1). For communities with invasive plants, the number of individuals of invasive species in each community was equal (i.e., 24 individuals). In the remaining six native plant communities, invasive species were not introduced. For a more detailed description of the experiment, see Wang et al. (2022) [35].

2.3. Soil Sample Collection and Physico-Chemical Analysis

On 11 October 2020, we took five soil samples (0–10 cm deep) under each of the 90 communities invaded by alien plant species [51,52]. The five soil samples from each container were thoroughly homogenized to form a composite sample. Then, a sub-sample of 10 mL was preserved at -80 °C until DNA extraction for soil microbial analysis. Another sub-sample was air-dried for analysis of soil physico-chemical properties. Soil pH was measured in a 1:2.5 soil:water (w:w) mixture using a digital pH meter (Mettler Toledo FE20, Shanghai, China). Soil organic matter was determined by the Walkley-Black acid digestion method. Soil N-NH4⁺, N-NO3⁻ (extracted from a 1:5 suspension of soil in 2 M KCl solution) and total N (digested with concentrated H₂SO₄ and HClO₄) were measured using a continuous flow analyzer (Auto Analyzer 3, Bran and Luebbe, Norderstedt, Germany). Total P (digested with concentrated H₂SO₄ and HClO₄) and available P (extracted with 0.5 M NaHCO₃) were determined by an inductively coupled plasma emission spectrometer (Optima 2100DV, PerkinElemer, Waltham, MA, USA).

2.4. Microbial Sample Collection and Sequencing

Soil DNA was extracted using a PowerSoil[™] DNA Isolation Kit (MoBio, Carlsbad, CA, USA) following the manufacturer's instructions and then sent to Novogene Co., Ltd., China for amplification, library preparation and sequencing. The V4 region of the bacterial 16S rDNA gene was amplified using primers 515F (GTGCCAGCMGCCGCGGTAA) and 806R (GGACTACNNGGGTATCTAAT) [53]. The fungal ITS region rDNA was amplified using primers ITS5-1737F (GGAAGTAAAAGTCGTAACAAGG) and ITS2-2043R (GCTGCGTTCTTCATCGATGC) [54]. The sequencing was done on an Illumina Novaseq 6000 platform, with 250 bp paired-end reads generated. The raw sequencing data have been deposited into the NCBI Sequence Read Archive (SRA) database (Accession Number: PRJNA870235 and PRJNA861831).

2.5. Bioinformatic Analysis

Barcode extraction, paired read assembly and quality filtering of the raw data were processed using the Software QIIME v.1.9.1 [55]. Chimeric sequences were identified and removed using UCHIME [56]. Operational taxonomic units (OTUs) were identified at the 97% similarity level with the UPARSE v.7.1b [57]. The taxonomy of each OTU of bacteria and fungi was assigned by the Uclust method using SILVA release 132 (https://www.

arb-silva.de/documentation/release-132, accessed on 5 July 2021) and UNITE database (http://qiime.org/scripts/assign_taxonomy.html, accessed on 5 July 2021), respectively.

2.6. Statistical Analysis

Alpha diversity (Shannon, Simpson, Chao1 and ACE indices) and beta diversity were calculated using the QIIME software (Version 1.9.1) [55]. The effects of invasive plant species richness (log₂ scale), evenness and their interaction on alpha diversity index, soil physicochemical properties and the relative abundance of the most abundant phyla and genera of soil fungi and bacteria were tested by linear mixed models. In these models, invasive species richness was treated as a fixed continuous term, evenness as a fixed categorical term and species composition as a random term [35]. The treatment with a single invasive species was considered a low evenness treatment because monocultures were on the regression lines of uneven rather than even treatments [35,58]. We performed Principal Coordinates Analysis (PCoA) and Adonis to test whether species composition of soil microbial communities differed between different treatments based on Weighted Unifrac distance. Linear regressions were used to assess the relationships of invasive species richness with the alpha diversity index, and relative abundances of phyla and genera of fungi and bacteria. Residuals of all variables were checked for homoscedasticity and normality. All statistical analyses were conducted using R 4.1.2 (https://www.r-project.org/, accessed on 10 June 2022).

3. Results

3.1. Effects of Invasive Species Diversity on Alpha Diversity of Soil Microbial Communities

Alien invasive species richness, evenness or their interaction did not significantly affect alpha diversity (OTUs, Shannon, Simpson, Chao1 and ACE) of either soil fungal communities (Table 1A, Figure 1A–E) or soil bacterial communities (Table 1B, Figure 1F–J).



Figure 1. Effects of invasive species richness and evenness on the OTUs, Shannon, Simpson, Chao1 and ACE of soil fungal communities (**A**–**E**) and soil bacterial communities (**F**–**J**). Raw data points are shown. See Table 1A,B for the results of the linear mixed models.

Table 1. Summary of linear mixed models testing the effects of invasive species richness, evenness and their interaction on alpha diversity of soil fungal communities (**A**), soil bacterial communities (**B**) and soil physico-chemical properties (**C**). Species composition was included as a random term. Degrees of freedom are (1, 38), (1, 48) and (1, 48) for the effects of richness, evenness and their interaction, respectively.

	Richn	ess (R)	Evenn	ess (E)	R' E			
Variable	F	Р	F	Р	F	Р		
		(A) Soil fu	ngal commun	ities				
OTUs	0.515	0.477	0.337	0.564	0.298	0.588		
Shannon	0.555	0.461	0.007	0.933	0.292	0.592		
Simpson	0.286	0.596	0.033	0.857	0.781	0.381		
Chao1	0.284	0.597	0.170	0.682	0.357	0.553		
ACE	0.341	0.563	0.128	0.722	0.316	0.577		
		(B) Soil bact	terial commu	nities				
OTUs	0.585	0.449	1.344	0.252	0.008	0.931		
Shannon	0.568	0.456	1.856	0.180	0.768	0.385		
Simpson	2.155	0.150	2.204	0.144	1.433	0.237		
Chao1	1.668	0.204	2.688	0.108	0.587	0.447		
ACE	1.232	0.274	2.529	0.118	0.150	0.701		
	(C) Soil physic	o-chemical pi	roperties				
pН	0.012	0.913	0.004	0.951	0.448	0.506		
Organic matter	0.019	0.891	0.724	0.399	1.317	0.257		
Total N	1.508	0.227	1.257	0.268	2.303	0.136		
NH4_N	0.021	0.886	1.835	0.182	0.843	0.363		
NO ₃ _N	0.005	0.944	2.427	0.126	2.755	0.104		
Total P	0.475	0.495	0.277	0.602	0.037	0.849		
Available P	0.058	0.811	2.977	0.091	0.801	0.375		
Available K	0.023	0.882	0.884	0.352	3.272	0.077		

3.2. Effects of Invasive Species Diversity on Soil Physico-Chemical Properties

Alien invasive species richness, evenness or their interaction had no significant effect on any of the soil physico-chemical properties (soil pH, organic matter, total N, NH₄_N, NO₃_N, total P, available P and available K; Table 1C, Table 2).

Table 2. Soil physico-chemical properties under communities with different levels of invasive plant species richness and evenness (mean \pm SE). See Table 1C for the results of the linear mixed models.

Richness	Evenness	рН	Organic matter (g/kg)	Total N (mg/g)	NH ₄ _N (mg/kg)	NO ₃ _N (mg/kg)	Total P (mg/g)	Available P (mg/kg)	Available K (mg/kg)
1	Low	7.27 ± 0.04	285.05 ± 17.48	0.77 ± 0.07	10.45 ± 1.20	33.76 ± 3.24	3.45 ± 0.06	122.05 ± 7.49	163.27 ± 8.90
2	Low	7.35 ± 0.05	303.00 ± 37.82	0.78 ± 0.08	11.90 ± 2.29	37.29 ± 6.68	3.43 ± 0.14	125.64 ± 17.72	182.28 ± 31.35
	High	7.34 ± 0.06	328.04 ± 27.28	0.58 ± 0.06	8.74 ± 1.33	35.72 ± 7.62	3.48 ± 0.11	122.13 ± 15.25	189.38 ± 24.27
4	Low	7.34 ± 0.05	275.48 ± 37.52	0.67 ± 0.09	9.96 ± 1.69	36.20 ± 6.91	3.32 ± 0.10	132.27 ± 12.46	180.98 ± 27.74
	High	7.26 ± 0.09	303.12 ± 18.88	0.65 ± 0.06	8.21 ± 0.61	31.32 ± 3.71	3.47 ± 0.12	105.33 ± 11.21	161.78 ± 13.50
8	Low	7.27 ± 0.04	296.57 ± 27.73	0.66 ± 0.05	11.12 ± 2.22	43.31 ± 6.00	3.39 ± 0.20	139.36 ± 13.88	191.89 ± 18.65
	High	7.29 ± 0.05	278.40 ± 11.35	0.70 ± 0.05	11.11 ± 1.48	26.67 ± 4.40	3.37 ± 0.20	115.02 ± 16.12	150.62 ± 16.54

3.3. Effects of Invasive Species Diversity on Soil Microbial Community Composition

Soil fungal communities in the 1-, 2- and 4-species treatments were separated from those in the 8-species treatment along the first axis of PCoA (Figure 2A). Results of the Adonis test also confirmed that species composition of soil fungal communities differed between the lower (1-, 2- and 4-species) and the higher (8-species) species richness treatments (Table 3A). For soil bacterial communities, the 1- and 8-species treatments were distinguished from the 2- and 4-species treatments along the first axis of PCoA (Figure 2B), and the results of the Adonis test also confirmed that the differences were significant (Table 3A). However, the Adonis test showed that neither the species compositions of soil fungal

communities nor soil bacterial communities differ significantly between the high and the low evenness treatment within each level of species richness (Table 3B).



Figure 2. Results of PCoA for soil fungal (**A**) and bacterial (**B**) communities. See Table 3A for the results of Adonis test. The bold boxes indicate the central location of the sample points under different richness treatments.

Table 3. Adonis test for differences in community composition of fungi and bacteria between different richness and evenness treatments of invasive species based on Bray-Curtis distance.

		df	F	R^2	Р
	(A) Comparison among different richness	s of alien in	vasive plan	t species	
Fungi	1 0		-	-	
	1 vs. 2 species	1 (48)	0.704	0.014	0.872
	1 vs. 4 species	1 (48)	1.135	0.023	0.291
	1 vs. 8 species	1 (48)	3.048	0.060	0.001
	2 vs. 4 species	1 (38)	0.747	0.019	0.807
	2 vs. 8 species	1 (38)	1.803	0.045	0.030
	4 vs. 8 species	1 (38)	1.967	0.049	0.016
Bacteria					
	1 vs. 2 species	1 (48)	1.333	0.027	0.113
	1 vs. 4 species	1 (48)	1.767	0.036	0.017
	1 vs. 8 species	1 (48)	1.149	0.023	0.242
	2 vs. 4 species	1 (38)	0.763	0.020	0.812
	2 vs. 8 species	1 (38)	1.595	0.040	0.026
	4 vs. 8 species	1 (38)	1.791	0.045	0.021
	(B) Comparison among different evennes	s of alien in	vasive plar	nt species	
Fungi			-	-	
-	Low vs. high evenness of the 2 species	1 (18)	0.910	0.048	0.517
	Low vs. high evenness of the 4 species	1 (18)	0.637	0.034	0.929
	Low vs. high evenness of the 8 species	1 (18)	0.737	0.039	0.844
Bacteria					
	Low vs. high evenness of the 2 species	1 (18)	0.694	0.037	0.930
	Low vs. high evenness of the 4 species	1 (18)	0.814	0.043	0.593
	Low vs. high evenness of the 8 species	1 (18)	0.958	0.051	0.501

3.4. Effects of Invasive Species Diversity on Relative Abundances of Soil Microbial Taxa

Regarding soil fungal taxa, Ascomycota, Basidiomycota and Rozellomycota were the three most abundant phyla (Figure 3A). Among them, invasive plant species richness had a significant effect on Ascomycota and Rozellomycota (Table 4); with increasing invasive plant species richness, the relative abundance of Ascomycota increased significantly, while that of Rozellomycota decreased (Figures 3A and 4A). Among the 20 most abundant

fungal genera, invasive plant species richness had a significant effect on *Bipolaris*, unidentified_*Rozellomycota_sp* and *Clitopilus* (Table 4A); with increasing invasive plant species richness, the relative abundance of *Bipolaris*, the most abundant genus, increased significantly, while that of unidentified_*Rozellomycota_sp* and *Clitopilus* decreased (Figures 3C and 4A).



Figure 3. Relative abundance of the 10 most abundant phyla of soil fungi (**A**), the 20 most abundant phyla of soil bacteria (**B**) and the 20 most abundant genera of soil fungi (**C**) and bacteria (**D**) under the communities with different invasive plant species richness and evenness.

Table 4. Summary of linear mixed models testing the effects of invasive species richness, evenness and their interaction on the relative abundance of (**A**) soil fungal and (**B**) soil bacterial taxa. Values are in bold when p < 0.05. Species composition was included as a random term. Degrees of freedom are (1, 38), (1, 48) and (1, 48) for the effects of richness, evenness and their interaction, respectively.

Variable	Richn	ess (R)	Evenn	ess (E)	$\mathbf{R} imes \mathbf{E}$			
	F	Р	F	Р	F	Р		
(A) Fungi Phylum								
Ascomycota	4.896	0.033	0.119	0.732	0.894	0.349		
Rozellomycota	7.365	0.010	0.093	0.762	0.180	0.673		
Genus								
Bipolaris	5.203	0.028	0.022	0.883	0.275	0.602		
unidentified_Rozellomycota_sp	7.835	0.008	0.043	0.836	0.011	0.918		
Clitopilus	6.076	0.018	0.606	0.440	0.222	0.639		

Variable	Richn	ess (R)	Evenn	ess (E)	$\mathbf{R} imes \mathbf{E}$			
	F	Р	F	Р	F	Р		
(B) Bacteria								
Phylum								
Actinobacteriota	4.920	0.033	1.838	0.182	0.844	0.363		
Bdellovibrionota	0.252	0.619	0.210	0.649	7.782	0.008		
Armatimonadota	1.065	0.309	4.358	0.042	0.018	0.894		
Genus								
Sphingomonas	1.465	0.234	0.067	0.796	4.576	0.038		
Buchnera	0.314	0.579	0.918	0.343	7.560	0.008		
Streptomyces	8.822	0.005	6.024	0.018	0.126	0.724		



Figure 4. Relationships of the relative abundance of the fungal phylum of Ascomycota and Rozellomycota, the fungal genus of *Bipolaris*, unidentified_*Rozellomycota_sp* and *Clitopilus* (**A**), the bacterial phylum of Actinobacteriota and the bacterial genus of *Streptomyces* (**B**) with invasive plant richness. Raw data points are shown. See Table 4 for the results of the linear mixed models.

Regarding soil bacterial taxa, Proteobacteria, unidentified_Bacteria and Actinobacteriota were the three most abundant phyla (Figure 3B). Invasive plant richness has a significant influence on Actinobacteriota, which increased significantly with the increase of invasive plant species richness (Table 4 and Figures 3B and 4B). Invasive plant evenness had a significant effect on Armatimonadota, and invasive plant richness and evenness had a significant interactive effect on Bdellovibrionota (Table 4B, Figure 3B). Among the 20 most abundant genera of bacteria, invasive plant species richness and evenness had a significant effect on *Streptomyces*, which increased significantly with the increase of invasive species richness (Table 4B, Figures 3D and 4B). In addition, the interaction of invasive plant richness and evenness had a significant effect on *Sphingomonas* and *Buchnera* (Table 4B, Figure 3D).

Table 4. Cont.

4. Discussion

A large body of studies has investigated the effects of alien plant invasions on soil microbial communities [14–16,20,21], but few have considered the effects of invasive species diversity (richness and evenness). While we found no significant effect of alien invasive plant species richness and evenness on the alpha diversity of soil fungal and bacterial communities, there was a significant effect of invasive species richness on the community composition of both soil fungi and bacteria and on the relative abundances of some specific microbial taxa.

4.1. Effects of Invasive Species Diversity on Alpha Diversity of Soil Microbial Communities

Although, in the same experimental setup, increasing invasive species richness was found to significantly change the species compositions of the plant communities by increasing the productivity of alien invasive plant species and decreasing that of native plant species [35], it had no impact on alpha diversity of either soil fungal or bacterial communities. Previous studies testing the role of native plant richness showed that alpha diversity of soil fungal communities can have a positive [16,59,60], negative [61,62] or no relationship [63–67] with native plant species richness. The positive relationship could be because of more diverse microhabitats and/or resources created by the higher richness of plant species [16,63,68]. For example, soil fungal richness increased with native plant richness through the increase of soil total carbon [16]. The negative relationship results from the greater impact of context (e.g., plant biomass, soil pH and plant species identity) [61,62]. For example, soil resources and plant species had a stronger effect on soil biota than plant species richness [69].

The neutral effect of invasive plant species richness on soil microbial diversity in this study may be related to the antagonistic effects among different invasive plants on soil microorganisms [20]. This can be explained from two aspects: different invasive plants have opposite effects on the same microbial group, and the same invasive plant communities have opposite effects on different microbial groups. In this study, the first aspect can be confirmed by the opposite effects of the relative abundance of different invasive plants on specific microbial groups (Table A2). For example, the relative abundance of the invasive plants *Alternanthera philoxeroides, Sesbania cannabina* and *Talinum paniculatum* was significantly negatively correlated with the relative abundance of *Bipolaris*, the genus with the highest relative abundance of fungi, while the relative abundance of *Bidens frondosa* showed the opposite trend (Table A2).

For the second aspect, when multiple invasive plants co-exist, some species favor some particular microbial taxa and inhibit others, with the result that there is no significant change, as reported before [23]. In this study, this explanation can be evidenced by the opposite trend of the effect of invasive species diversity on the relative abundance of some fungal taxa (i.e., Ascomycota vs. Rozellomycota and *Bipolaris* vs. unidentified_*Rozellomycota*_sp and *Clitopilus;* Figure 4A). The same situation is present in soil bacterial communities, i.e., increasing invasive species richness increased the relative abundance of Actinobacteriota and decreased that of *Streptomyces* (Figure 4B).

Soil is an important environmental factor for microbial life. Many studies on the effects of plant richness on microbial life have found that this relationship is ultimately explained not only by plant species richness but also by soil physico-chemical properties, such as soil pH [70,71]. In this study, although soil microbial diversity was significantly negatively correlated with some soil physico-chemical properties (Table A3), invasive species richness and evenness had no significant effect on soil physico-chemical properties. Therefore, invasive species diversity may not change soil microbial diversity via altering soil physico-chemical properties.

4.2. Effects of Invasive Species Diversity on Soil Microbial Taxa and Community Composition

Specific taxa of microorganisms play an important role in maintaining microbial community structure and affecting ecosystem functions [72,73] (In this study, invasive species richness and evenness have significant effects on some specific taxa of fungi and bacteria (Table 4; Figure 4). Our results revealed that high invasive plant richness promoted the most abundant fungal phylum of Ascomycota and the most abundant fungal genus of *Bipolaris* and suppressed the fungal phylum of Rozellomycota and fungal genus of unidentified_*Rozellomycota_sp* and *Clitopilus*.

Ascomycota is one of the most common phyla among soil fungi and is the most abundant phyla in the soil of many invasive sites [16,29]. Similarly, Ascomycota possessed more than 50% relative abundance in each treatment of this study and increased significantly with increasing invasive species richness. As Ascomycota contained a wide diversity of fungal taxa, ranging from pathogens to mutualists, its ecological function depends on the comprehensive effect of each member. The increasing tendency of relative abundance of Ascomycota with invasive species richness in this study may be partly explained by its member *Bipolaris*, the most abundant genus of fungi, which also increased significantly with increasing invasive species richness. The genus of *Bipolaris* includes more than 20 species of pathogens that can infect a broad range of grasses [74–76], including invasive plants (e.g., *Microstegium*) [77]. Our results suggest that increasing invasive species richness may facilitate pathogen emergence and amplification, and these effects may inhibit native plant growth and promote further invasions. On the other hand, the accumulation of pathogens may also inhibit the invasion of alien plants [77].

For soil bacterial communities, Actinobacteriota, one of the largest taxonomic units of bacteria, increased significantly with increasing invasive species richness. As plant growthpromoting rhizobacteria, Actinobacteriota constitute important drivers of rhizosphere nutrient cycling and play a role as biocontrol agents against a range of pathogenic fungi and promote plant growth by phosphate solubilization, secondary metabolite production and antimicrobial synthesis [78–80]. Our results agreed with the findings of a previous study, which showed that increasing native crop diversity increased the relative abundance of Actinobacteriota [81]. In this study, the increased relative abundance of Actinobacteriota in more diverse invasive plant communities has the same trend with its member *Streptomyces*, which is the most abundant and arguably the most important actinomycetes, and is a good source of bioactive compounds, antibiotics, and extracellular enzymes [79,80,82]. Although studies have shown that plant invasion (e.g., Ageratina adenophora) increases the relative abundance of *Streptomyces* [83], the relative abundances of Actinobacteriota and Streptomyces were all negatively related to the relative abundance of the invasive plants Sesbania cannabina and Talinum paniculatum (Table A2). Therefore, invasive species richness may affect this microbe (Actinobacteriota and Streptomyces) through species-species interactions and species-environment interactions, and this effect may be beneficial to the growth and competitiveness of invasive plants themselves [83].

5. Conclusions

We conclude that alien invasive plant species diversity had no effect on the alpha diversity of both soil fungal and bacterial communities, likely because the dominant microbial groups showed opposite responses to alien invasive species diversity. One caveat is that the soil samples were collected only after one growing season of invasion, so the impact of invasive species diversity on the alpha diversity of soil microbial communities may not be big enough. Thus, future studies could test the longer-term effect of alien invasive plant species diversity on the alpha diversity of soil microbes. Given the important effects of soil microbial communities on plant growth, especially pathogens, it is possible that soil microorganisms may have important effects on native plants or invasive plants to alter community composition. Therefore, it is worth further researching if invasive plant diversity can affect the plant community composition through the influence of soil pathogens.

Author Contributions: Conceptualization, X.W. (Xiaoyan Wang), X.W. (Xue Wang), J.W. and F.Y.; methodology, X.W. (Xiaoyan Wang), X.W. (Xue Wang) and W.W.; formal analysis, X.W. (Xiaoyan Wang) and W.W.; writing—original draft preparation, X.W. (Xiaoyan Wang) and J.W.; writing—review and editing, X.W. (Xue Wang), J.W. and F.Y.; funding acquisition, X.W. (Xiaoyan Wang) and J.W. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Basic Public Welfare Research Projects of Zhejiang Province (No. LGN19C150004); Natural Science Foundation of Zhejiang Province (Nos. LY22C030001, LTY22C030004); National Natural Science Foundation of China (No. 31870504); 2021 Taizhou College Students Science and Technology Innovation Project (No. 2).

Institutional Review Board Statement: There are no humans or vertebrates involved in the current study.

Informed Consent Statement: No human subjects were involved or recruited in the current study.

Data Availability Statement: Data will be available upon request.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A

Table A1. Species composition ratio in different richness levels and evenness levels.

					1-sp	ecies								ц	ich or					2-spe	ecies			т															
Invasive species Bidens nilosa	1 24	2	3	4	5	6	7	8	9	10	1 12	2	3	н 4	5	6 12	7	8	9	10	1 22	2	3	4	20w ev 5	6 2	7	8	9	10									
Bidens frondosa Ageratum conyzoides		24	24								12			12				12	12		2			22		-		22	22										
Sesbania cannabina Talinum paniculatum				24	24									12	12	12				12				2	2	22				2									
Celosia argentea Solidago canadensis						24	24					12			12		12	12				2		-	22		22	2		-									
Mirabilis jalapa Alternanthera								24				12								12		22					_	_		22									
philoxeroides Aster subulatus									24	24			12 12				12		12				22 2				2		22										
		High	ovon	nose			4-8	specie	5				т		onnos	26							ц	Jigh o	vonno				8-sp	ecies			T.		onno	20			
1	n	2	4	E	6	7	0	0	10	1	2	2	4 L		6	-7	0	0	10	1	2	2	4	ngn e	6	55 7	0	0	10	1	2	2	1	500 60	6	7	0	0	10
6	6	3 6	4	5	6	1	8	9	10	18	2 3	3 2	4	5	6	/	8	9	10	1	2	3	4	3	3	3	8 3	3	3	1	2	3	4 3	5 2	1	13	8 1	9	2
6				6	6	6				3				3	18	1				3	3			3	3	3	3	3	3	13	1			1	2	1	1	2	3
	((6		6	6		(6		2	2	18		1	2		10	3	2	3	3	2	3	3	3	3	3	3	2	13	3	1	1	1	1	2	2	1
6	6	6	6				6	6	6	2	2	3	2				2	18	18	3	3	3	3	3	3	3	3	3	3 3	3 2	1	1	13	2	15	2	3 2	3	2
				6	6		6	6						1	3	18	18	2		3	3	3	3	3	3	3	3	-		2	1	2	2	13	3	1	1	-	
		6		6	6	6	6	6	6			18		2	2	3	3 1	3	2	3	3	3	3	3	3	3	з	3	3	1	1	2	1	1	1	1	13	13	1
6				6		6	0	0	6	1				18			1	5	1	3	3	3	3		5	3	3	3	3	1	2	1	1		1	3	1	1	13
	6	6	6					6			18	1	3					1		3	3	3	3	3	3			3	3	1	2	13	2	3	2			1	1

The numbers in the third row represent the number of experiment replicates. For 1-species, there are three replicates for each species, which was not shown in the table. Other numbers represent the seedling number of different species planted in each community.

	Bidens pilosa	Bidens frondosa	Ageratum conyzoides	Sesbania cannabina	Talinum paniculatum	Celosia argentea	Solidago canadensis	Mirabilis jalapa	Alternanthera philoxeroides	Aster subulatus
Fungi										
Phylum										
Áscomycota	-0.375 *	0.265	-0.328	-0.465 **	-0.248	0.052	-0.241	0.337	-0.474 **	-0.340
Basidiomycota	0.195	0.124	0.080	0.531 **	0.179	-0.114	0.202	-0.219	0.367 *	-0.182
Rozellomycota	0.265	-0.225	0.541 **	0.391 *	0.066	0.065	0.099	0.186	0.481 **	0.174
Chytridiomycota	-0.178	-0.281	0.314	-0.166	-0.007	0.243	-0.106	-0.152	0.229	0.044
Mortierellomycota	-0.151	-0.037	0.293	0.352	-0.102	-0.278	0.331	-0.256	0.370 *	0.598 **
Glomeromycota	0.126	0.206	0.145	-0.220	0.237	-0.312	0.066	0.382 *	-0.173	-0.045
Mucoromycota	-0.001	-0.078	-0.126	0.492 **	0.174	0.069	0.102	-0.100	0.018	0.148
Monoblepharomycota	0.255	-0.122	0.052	-0.189	0.549 **	0.063	-0.005	-0.077	-0.100	-0.079
Genus										
Binolaris	-0.309	0.419 *	-0.352	-0.389 *	-0.426 *	-0.350	-0.139	0 197	-0.429 *	-0.282
Alternaria	0.026	-0.366 *	-0.225	-0.090	0.094	0.380 *	-0.043	0 161	-0.281	0.068
Monogranhella	-0.181	-0.242	0.514 **	-0.155	0.174	0 297	-0.045	-0.083	-0.201	-0.155
monogrupheni	0.101	0.212	0.011	0.100	0.17 1	0.277	0.010	0.000	0.210	0.100
unidentified_Tremellodendropsidales_sp	0.038	-0.007	-0.162	0.323	0.101	-0.076	0.128	-0.340	0.398 *	-0.196
Blumeria	0.028	-0.182	-0.142	0.388 *	0.270	0.153	-0.106	0.516 **	0.417 *	0.160
unidentified	-0.026	-0.058	-0.148	0.288	0.075	-0.089	0.156	-0.304	0.366 *	-0.191
Cladosporium	-0.020	-0.317	-0.230	0.051	0.006	0.101	-0.116	0.280	-0.187	-0.078
unidentified_Rozellomycota_sp	0.279	-0.237	0.588 **	0.445 *	0.048	0.097	0.056	0.298	0.476 **	0.124
Clitopilus	0.668 **	0.195	0.293	0.570 **	0.157	-0.035	0.187	0.113	0.256	0.137
Albifimbria	-0.124	-0.164	-0.114	-0.153	0.037	0.266	0.090	-0.096	-0.123	0.346
Trichoderma	-0.050	0.093	0.440 *	0.112	0.329	-0.164	-0.034	-0.178	0.137	0.121
unidentified_Dothideomycetes_sp	0.335	-0.038	0.049	-0.051	0.289	-0.089	-0.181	0.104	-0.058	-0.167
Bacteria										
Phylum										
Proteobacteria	-0.013	0.207	-0.139	0.197	0.267	-0.062	0.225	0.058	0.060	-0.498 **
unidentified_Bacteria	0.242	-0.147	0.221	0.229	-0.186	-0.024	0.027	0.062	0.488 **	-0.113
Actinobacteriota	-0.431 *	0.061	-0.180	-0.478 **	-0.358 *	-0.117	-0.271	0.152	-0.277	0.169
Verrucomicrobiota	-0.001	-0.100	0.212	-0.209	0.142	0.056	-0.115	-0.295	-0.357 *	-0.055
Firmicutes	0.006	-0.019	-0.008	-0.185	-0.133	-0.055	-0.176	-0.064	-0.182	0.506 **
Thermoplasmatota	0.149	-0.269	0.394 *	0.068	0.013	0.294	-0.203	-0.066	0.159	0.527 **
Acidobacteriota	0.132	-0.400 *	-0.139	0.109	0.055	0.158	0.233	-0.186	-0.060	0.172
Myxococcota	-0.029	-0.169	-0.056	0.490 **	0.244	0.421 *	-0.021	-0.028	-0.165	-0.228
Gemmatimonadota	-0.005	-0.192	0.349	-0.353	-0.219	-0.022	0.055	-0.190	0.106	0.065
Gemmatimonadetes	-0.024	0.363 *	-0.019	-0.438 *	-0.241	-0.270	0.339	-0.178	0.007	0.076
Latescibacterota	0.228	-0.286	-0.155	0.332	0.406 *	0.218	0.152	-0.081	-0.017	0.023
Bdellovibrionota	0.045	-0.092	0.249	0.221	0.188	0.191	0.020	-0.038	-0.047	-0.377 *
Nitrospirota	-0.106	-0.036	0.042	-0.123	-0.063	-0.065	0.381 *	-0.287	0.005	-0.383 *
RCP2-54	-0.005	-0.025	0.121	0.333	-0.034	0.107	0.118	0.172	0.374 *	-0.164
Desulfobacterota	0.072	0.097	0.146	-0.221	-0.056	-0.025	-0.092	-0.039	-0.138	0.723 **
Genus										
Acidibacter	-0.086	-0.271	0.227	0.010	0.386 *	0.113	0.037	-0.116	-0.328	-0.067
IS-44	0.341	-0.424 *	-0.105	0.376 *	0.354	0.116	-0.180	0.125	0.035	0.094
Sphingomonas	-0.284	0.313	-0.165	-0.302	-0.239	-0.341	0.373 *	-0.099	0.102	-0.296
Ópitutus	0.089	0.071	0.320	-0.309	0.048	-0.014	-0.237	-0.189	-0.325	0.171
Lactobacillus	-0.059	-0.068	-0.010	-0.033	-0.065	-0.097	-0.045	-0.130	-0.111	0.469 **
Ohtaekwangia	0.146	-0.278	-0.272	0.430 *	0.084	0.142	-0.021	0.037	0.088	-0.062

Table A2. Pearson correlation analysis between top 20 predominant phyla and genus and percentage of initial plant number for 10 invasive plants.

Table A2.	Cont.
-----------	-------

	Bidens pilosa	Bidens frondosa	Ageratum conyzoides	Sesbania cannabina	Talinum paniculatum	Celosia argentea	Solidago canadensis	Mirabilis jalapa	Alternanthera philoxeroides	Aster subulatus
SWB02	0.184	-0.178	-0.121	0.337	0.066	0.214	0.158	-0.025	-0.057	0.063
Terrimonas	0.192	-0.019	0.368 *	0.007	0.190	-0.172	0.100	0.235	0.044	0.165
Aeromicrobium	-0.237	0.331	-0.242	-0.237	-0.240	-0.144	-0.343	-0.199	-0.283	-0.202
Subgroup_10	-0.053	-0.186	0.018	-0.188	-0.071	-0.131	0.252	-0.076	-0.010	0.445 *
Lacunispĥaera	-0.109	-0.113	0.225	-0.065	0.194	0.078	-0.061	-0.307	-0.241	-0.380 *
Dongia	-0.073	-0.065	-0.052	0.075	0.244	-0.118	0.229	-0.044	0.216	-0.342
Allorhizobium-Neorhizobium- Pararhizobium-Rhizobium	-0.324	0.401 *	-0.002	0.080	-0.343	-0.288	-0.062	0.154	0.239	-0.358 *
Streptomyces	-0.268	-0.017	-0.089	-0.524 **	-0.363 *	-0.199	-0.410 *	-0.160	-0.372 *	-0.070
Azoarcus	0.383 *	0.007	-0.210	0.247	0.359 *	0.344	-0.224	0.077	0.288	-0.307
YC-ZSS-LKJ147	0.059	-0.290	0.315	-0.314	-0.232	-0.010	-0.034	-0.252	0.039	-0.112

Only significant or critical significant results are shown here. Values in the table are correlation coefficients, with bold font indicating a significant correlation and bold and italic font indicating critical significance. Asterisk indicates a significant correlation. "*" represents $0.01 \le p < 0.05$; "**" represents p < 0.01.

Table A2	Poarcon correlation ar	alveic botwoon coi	1 nh	veico ch	omical	proportion and	funco	l and h	actorial	divorcity	indicoc
Table A5.	i carson correlation al	ary sis between so	որո	y SICO-CII	ennear	properties and	Tunga	i anu ba	acteriar	arversity	munces.

	Fungi					Bacteria				
	OTUs	Shannon	Simpson	Chao1	ACE	OTUs	Shannon	Simpson	Chao1	ACE
pН	0.069	0.092	0.101	0.070	0.065	-0.009	0.060	0.044	0.030	-0.005
Organic matter	0.004	0.066	0.009	-0.013	-0.004	0.054	-0.154	-0.191	0.044	0.063
Total N	0.012	0.002	-0.012	-0.009	-0.010	-0.008	-0.063	-0.046	-0.011	0.028
NH4_N	-0.080	-0.043	-0.066	-0.082	-0.079	0.006	0.073	0.052	-0.047	-0.052
NO ₃ _N	-0.275 **	-0.106	-0.033	-0.239 *	-0.241 *	-0.067	0.042	0.065	-0.140	-0.124
Total P	-0.252 *	-0.191	-0.123	-0.265 **	-0.240 *	0.039	0.074	0.073	0.029	0.051
Available P	-0.225 *	-0.207 *	-0.182	-0.188	-0.178	-0.225 *	-0.068	0.017	-0.261 *	-0.250 *
Available K	-0.039	-0.097	-0.139	-0.025	-0.015	-0.158	-0.034	-0.020	-0.208 *	-0.203 *

Values in the table are correlation coefficients, with bold font indicating a significant correlation and bold and italic font indicating critical significance. Asterisk indicates a significant correlation. "*" represents $0.01 \le p < 0.05$; "**" represents p < 0.01.

References

- 1. van Kleunen, M.; Dawson, W.; Essl, F.; Pergl, J.; Winter, M.; Weber, E.; Kreft, H.; Weigelt, P.; Kartesz, J.; Nishino, M.; et al. Global exchange and accumulation of non-native plants. *Nature* **2015**, *525*, 100–103. [CrossRef] [PubMed]
- Lambertini, M.; Leape, J.; Marton-Lefèvre, J.; Mittermeier, R.A.; Rose, M.; Robinson, J.G.; Stuart, S.N.; Waldman, B.; Genovesi, P. Invasives: A major conservation threat. *Science* 2011, 333, 404–405. [CrossRef] [PubMed]
- Atwater, D.Z.; Callaway, R.M. Extended consequences of selection by exotic invaders on natives. *New Phytol.* 2022, 233, 588–591.
 [CrossRef] [PubMed]
- Wang, W.; Shi, J.; Xie, Q.; Jiang, Y.; Yu, N.; Wang, E. Nutrient exchange and regulation in arbuscular mycorrhizal symbiosis. *Mol. Plant* 2017, 10, 1147–1158. [CrossRef] [PubMed]
- 5. Pysek, P.; Hulme, P.E.; Simberloff, D.; Bacher, S.; Blackburn, T.M.; Carlton, J.T.; Dawson, W.; Essl, F.; Foxcroft, L.C.; Genovesi, P.; et al. Scientists' warning on invasive alien species. *Biol. Rev.* 2020, *95*, 1511–1534. [CrossRef]
- Simberloff, D.; Martin, J.L.; Genovesi, P.; Maris, V.; Wardle, D.A.; Aronson, J.; Courchamp, F.; Galil, B.; Garcia-Berthou, E.; Pascal, M.; et al. Impacts of biological invasions: What's what and the way forward. *Trends Ecol. Evol.* 2013, 28, 58–66. [CrossRef]
- Vila, M.; Espinar, J.L.; Hejda, M.; Hulme, P.E.; Jarosik, V.; Maron, J.L.; Pergl, J.; Schaffner, U.; Sun, Y.; Pysek, P. Ecological impacts of invasive alien plants: A meta-analysis of their effects on species, communities and ecosystems. *Ecol. Lett.* 2011, 14, 702–708. [CrossRef]
- 8. Xu, H.; Liu, Q.; Wang, S.; Yang, G.; Xue, S. A global meta-analysis of the impacts of exotic plant species invasion on plant diversity and soil properties. *Sci. Total Environ.* **2022**, *810*, 152286. [CrossRef]
- Pisani, D.; Pazienza, P.; Perrino, E.V.; Caporale, D.; De Lucia, C. The economic valuation of ecosystem services of biodiversity components in protected areas: A review for a framework of analysis for the Gargano National Park. *Sustainability* 2021, 13, 11726. [CrossRef]
- Stinca, A.; Musarella, C.M.; Rosati, L.; Laface, V.L.A.; Licht, W.; Fanfarillo, E.; Wagensommer, R.P.; Galasso, G.; Fascetti, S.; Esposito, A.; et al. Italian vascular flora: New findings, updates and exploration of floristic similarities between regions. *Diversity* 2021, 13, 600. [CrossRef]
- 11. Byun, C.; Lee, E.J. Ecological application of biotic resistance to control the invasion of an invasive plant, *Ageratina altissima*. *Ecol. Evol.* **2017**, *7*, 2181–2192. [CrossRef] [PubMed]
- 12. Hoffberg, S.L.; Mauricio, R.; Hall, R.J. Control or re-treat? Model-based guidelines for managing established plant invasions. *Biol. Invasions* **2018**, *20*, 1387–1402. [CrossRef]
- Zhang, Z.; Liu, Y.; Brunel, C.; van Kleunen, M. Soil-microorganism-mediated invasional meltdown in plants. *Nat. Ecol. Evol.* 2020, 4, 1612–1621. [CrossRef] [PubMed]
- 14. Zhang, G.L.; Bai, J.H.; Tebbe, C.C.; Huang, L.B.; Jia, J.; Wang, W.; Wang, X.; Yu, L.; Zhao, Q.Q. Plant invasion reconstructs soil microbial assembly and functionality in coastal salt marshes. *Mol. Ecol.* **2022**, *31*, 4478–4494. [CrossRef] [PubMed]
- 15. Torres, N.; Herrera, I.; Fajardo, L.; Bustamante, R.O. Meta-analysis of the impact of plant invasions on soil microbial communities. BMC Evol. Biol. 2021, 21, 172. [CrossRef] [PubMed]
- 16. Shen, C.; Wang, J.; He, J.Z. Plant diversity enhances soil fungal diversity and microbial resistance to plant invasion. *Appl. Environ. Microbiol.* **2021**, *87*, e00251-00221. [CrossRef]
- 17. Li, Y.-P.; Feng, Y.-L.; Chen, Y.-J.; Tian, Y.-H. Soil microbes alleviate allelopathy of invasive plants. *Sci. Bull.* **2015**, *60*, 1083–1091. [CrossRef]
- 18. Zhang, F.-J.; Li, Q.; Chen, F.-X.; Xu, H.-Y.; Inderjit; Wan, F.-H. Arbuscular mycorrhizal fungi facilitate growth and competitive ability of an exotic species Flaveria bidentis. *Soil Biol. Biochem.* **2017**, *115*, 275–284. [CrossRef]
- 19. van der Putten, W.H.; Klironomos, J.N.; Wardle, D.A. Microbial ecology of biological invasions. ISME J. 2007, 1, 28–37. [CrossRef]
- Stefanowicz, A.M.; Stanek, M.; Nobis, M.; Zubek, S. Species-specific effects of plant invasions on activity, biomass, and composition of soil microbial communities. *Biol. Fertil. Soils* 2016, 52, 841–852. [CrossRef]
- 21. Jacobs, K.; Conradie, T.; Jacobs, S. Microbial communities in the Fynbos region of South Africa: What happens during woody alien plant invasions. *Diversity* 2020, *12*, 254. [CrossRef]
- 22. Kuebbing, S.E.; Classen, A.T.; Simberloff, D. Two co-occurring invasive woody shrubs alter soil properties and promote subdominant invasive species. *J. Appl. Ecol.* **2014**, *51*, 124–133. [CrossRef]
- 23. Gibbons, S.M.; Lekberg, Y.; Mummey, D.L.; Sangwan, N.; Ramsey, P.W.; Gilbert, J.A. Invasive plants rapidly reshape soil properties in a grassland ecosystem. *mSystems* **2017**, *2*, e00178-00116. [CrossRef] [PubMed]
- 24. Qu, T.; Du, X.; Peng, Y.; Guo, W.; Zhao, C.; Losapio, G. Invasive species allelopathy decreases plant growth and soil microbial activity. *PLoS ONE* **2021**, *16*, e0246685. [CrossRef] [PubMed]
- 25. Ehrenfeld, J.G. Effects of exotic plant invasions on soil nutrient cycling processes. Ecosystems 2003, 6, 503–523. [CrossRef]
- 26. Gaggini, L.; Rusterholz, H.-P.; Baur, B. The invasive plant Impatiens glandulifera affects soil fungal diversity and the bacterial community in forests. *Appl. Soil Ecol.* **2018**, *124*, 335–343. [CrossRef]
- 27. Bell, J.K.; Siciliano, S.D.; Lamb, E.G. A survey of invasive plants on grassland soil microbial communities and ecosystem services. *Sci. Data* 2020, *7*, 86. [CrossRef] [PubMed]
- Zhang, P.; Li, B.; Wu, J.; Hu, S. Invasive plants differentially affect soil biota through litter and rhizosphere pathways: A meta-analysis. *Ecol. Lett.* 2019, 22, 200–210. [CrossRef]

- 29. Wang, C.; Zhou, J.; Liu, J.; Jiang, K.; Xiao, H.; Du, D. Responses of the soil fungal communities to the co-invasion of two invasive species with different cover classes. *Plant Biol.* **2018**, *20*, 151–159. [CrossRef]
- 30. Liao, C.; Peng, R.; Luo, Y.; Zhou, X.; Wu, X.; Fang, C.; Chen, J.; Li, B. Altered ecosystem carbon and nitrogen cycles by plant invasion: A meta-analysis. *New Phytol.* **2008**, *177*, 706–714. [CrossRef]
- Vogelsang, K.M.; Bever, J.D. Mycorrhizal densities decline in association with nonnative plants and contribute to plant invasion. *Ecology* 2009, 90, 399–407. [CrossRef] [PubMed]
- 32. Shah, M.A.; Beaulieu, M.-E.; Reshi, Z.A.; Qureshi, S.; Khasa, D.P. A cross-city molecular biogeographic investigation of arbuscular mycorrhizas in *Conyza canadensis* rhizosphere across native and non-native regions. *Ecol. Process.* **2015**, *4*, 7. [CrossRef]
- Řezáčová, V.; Řezáč, M.; Gryndler, M.; Hršelová, H.; Gryndlerová, H.; Michalová, T. Plant invasion alters community structure and decreases diversity of arbuscular mycorrhizal fungal communities. *Appl. Soil Ecol.* 2021, 167, 104039. [CrossRef]
- Custer, G.F.; van Diepen, L.T.A. Plant invasion has limited impact on soil microbial α-diversity: A meta-analysis. *Diversity* 2020, 12, 112. [CrossRef]
- Wang, X.; Wang, J.; Hu, B.; Zheng, W.-L.; Li, M.; Shen, Z.-X.; Yu, F.-H.; Schmid, B.; Li, M.-H. Richness, not evenness, of invasive plant species promotes invasion success into native plant communities via selection effects. *Oikos* 2022, 2022, e08966. [CrossRef]
- 36. Wilsey, B.J.; Teaschner, T.B.; Daneshgar, P.P.; Isbell, F.I.; Polley, H.W. Biodiversity maintenance mechanisms differ between native and novel exotic-dominated communities. *Ecol. Lett.* **2009**, *12*, 432–442. [CrossRef]
- Kuebbing, S.E.; Souza, L.; Sanders, N.J. Effects of co-occurring non-native invasive plant species on old-field succession. *For. Ecol. Manag.* 2014, 324, 196–204. [CrossRef]
- Kuebbing, S.E.; Classen, A.T.; Sanders, N.J.; Simberloff, D. Above- and below-ground effects of plant diversity depend on species origin: An experimental test with multiple invaders. *New Phytol.* 2015, 208, 727–735. [CrossRef]
- Zak, D.R.; Tilman, D.; Parmenter, R.R.; Rice, C.W.; Fisher, F.M.; Vose, J.; Milchunas, D.; Martin, C.W. Plant production and soil microorganisms in late-successional ecosystems: A continental-scale study. *Ecology* 1994, 75, 2333–2347. [CrossRef]
- Waldrop, M.P.; Holloway, J.M.; Smith, D.B.; Goldhaber, M.B.; Drenovsky, R.E.; Scow, K.M.; Dick, R.; Howard, D.; Wylie, B.; Grace, J.B. The interacting roles of climate, soils, and plant production on soil microbial communities at a continental scale. *Ecology* 2017, 98, 1957–1967. [CrossRef]
- 41. McTee, M.R.; Lekberg, Y.; Mummey, D.; Rummel, A.; Ramsey, P.W. Do invasive plants structure microbial communities to accelerate decomposition in intermountain grasslands? *Ecol. Evol.* **2017**, *7*, 11227–11235. [CrossRef] [PubMed]
- 42. Broughton, L.C.; Gross, K.L. Patterns of diversity in plant and soil microbial communities along a productivity gradient in a Michigan old-field. *Oecologia* **2000**, *125*, 420–427. [CrossRef] [PubMed]
- 43. Wilsey, B.J.; Chalcraft, D.R.; Bowles, C.M.; Willig, M.R. Relationships among indices suggest that richness is an incomplete surrogate for grassland biodiversity. *Ecology* **2005**, *86*, 1178–1184. [CrossRef]
- 44. Wang, X.-Y.; Ge, Y.; Gao, S.; Chen, T.; Wang, J.; Yu, F.-H. Evenness alters the positive effect of species richness on community drought resistance via changing complementarity. *Ecol. Indic.* **2021**, *133*, 108464. [CrossRef]
- 45. Wang, J.; Yu, F.-H. Effects of functional diversity and functional dominance on complementary light use depend on evenness. *J. Veg. Sci.* 2018, 29, 726–736. [CrossRef]
- 46. Sun, Y.-Q.; Wang, J.; Shen, C.; He, J.-Z.; Ge, Y. Plant evenness modulates the effect of plant richness on soil bacterial diversity. *Sci. Total Environ.* **2019**, *662*, 8–14. [CrossRef]
- 47. Zhang, Y.; Chen, H.Y.H.; Reich, P.B. Forest productivity increases with evenness, species richness and trait variation: A global meta-analysis. *J. Ecol.* **2012**, *100*, 742–749. [CrossRef]
- Filstrup, C.T.; King, K.B.S.; McCullough, I.M. Evenness effects mask richness effects on ecosystem functioning at macro-scales in lakes. *Ecol. Lett.* 2019, 22, 2120–2129. [CrossRef]
- 49. Lamb, E.G.; Kennedy, N.; Siciliano, S.D. Effects of plant species richness and evenness on soil microbial community diversity and function. *Plant Soil* **2011**, *338*, 483–495. [CrossRef]
- Kirwan, L.; LÜScher, A.; SebastiÀ, M.T.; Finn, J.A.; Collins, R.P.; Porqueddu, C.; Helgadottir, A.; Baadshaug, O.H.; Brophy, C.; Coran, C.; et al. Evenness drives consistent diversity effects in intensive grassland systems across 28 European sites. *J. Ecol.* 2007, 95, 530–539. [CrossRef]
- Sadiqi, S.; Hamza, M.; Ali, F.; Alam, S.; Shakeela, Q.; Ahmed, S.; Ayaz, A.; Ali, S.; Saqib, S.; Ullah, F.; et al. Molecular characterization of bacterial isolates from soil samples and evaluation of their antibacterial potential against MDRS. *Molecules* 2022, 27, 6281. [CrossRef]
- 52. Porazinska, D.L.; Farrer, E.C.; Spasojevic, M.J.; Bueno de Mesquita, C.P.; Sartwell, S.A.; Smith, J.G.; White, C.T.; King, A.J.; Suding, K.N.; Schmidt, S.K. Plant diversity and density predict belowground diversity and function in an early successional alpine ecosystem. *Ecology* **2018**, *99*, 1942–1952. [CrossRef] [PubMed]
- 53. Bergmann, G.T.; Bates, S.T.; Eilers, K.G.; Lauber, C.L.; Caporaso, J.G.; Walters, W.A.; Knight, R.; Fierer, N. The under-recognized dominance of Verrucomicrobia in soil bacterial communities. *Soil Biol. Biochem.* **2011**, *43*, 1450–1455. [CrossRef] [PubMed]
- 54. Wei, W.; Yang, M.; Liu, Y.; Huang, H.; Ye, C.; Zheng, J.; Guo, C.; Hao, M.; He, X.; Zhu, S. Fertilizer N application rate impacts plant-soil feedback in a sanqi production system. *Sci. Total Environ.* **2018**, *633*, 796–807. [CrossRef]
- Caporaso, J.G.; Kuczynski, J.; Stombaugh, J.; Bittinger, K.; Bushman, F.D.; Costello, E.K.; Fierer, N.; Peña, A.G.; Goodrich, J.K.; Gordon, J.I.; et al. QIIME allows analysis of high-throughput community sequencing data. *Nat. Methods* 2010, 7, 335–336. [CrossRef] [PubMed]

- 56. Edgar, R.C.; Haas, B.J.; Clemente, J.C.; Quince, C.; Knight, R. UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* **2011**, *27*, 2194–2200. [CrossRef]
- 57. Edgar, R.C. UPARSE: Highly accurate OTU sequences from microbial amplicon reads. Nat. Methods 2013, 10, 996–998. [CrossRef]
- Schmitz, M.; Flynn, D.F.B.; Mwangi, P.N.; Schmid, R.; Scherer-Lorenzen, M.; Weisser, W.W.; Schmid, B. Consistent effects of biodiversity on ecosystem functioning under varying density and evenness. *Folia Geobot.* 2013, 48, 335–353. [CrossRef]
- 59. Peay, K.G.; Baraloto, C.; Fine, P.V.A. Strong coupling of plant and fungal community structure across western Amazonian rainforests. *ISME J.* 2013, 7, 1852–1861. [CrossRef]
- Dassen, S.; Cortois, R.; Martens, H.; de Hollander, M.; Kowalchuk, G.A.; van der Putten, W.H.; de Deyn, G.B. Differential responses of soil bacteria, fungi, archaea and protists to plant species richness and plant functional group identity. *Mol. Ecol.* 2017, 26, 4085–4098. [CrossRef]
- 61. Yang, N.; Zhang, Y.; Li, J.; Li, X.; Ruan, H.; Bhople, P.; Keiblinger, K.; Mao, L.; Liu, D. Interaction among soil nutrients, plant diversity and hypogeal fungal trophic guild modifies root-associated fungal diversity in coniferous forests of Chinese Southern Himalayas. *Plant Soil* **2022**. [CrossRef]
- Tedersoo, L.; Bahram, M.; Cajthaml, T.; Põlme, S.; Hiiesalu, I.; Anslan, S.; Harend, H.; Buegger, F.; Pritsch, K.; Koricheva, J.; et al. Tree diversity and species identity effects on soil fungi, protists and animals are context dependent. *ISME J.* 2016, 10, 346–362. [CrossRef] [PubMed]
- Waldrop, M.P.; Zak, D.R.; Blackwood, C.B.; Curtis, C.D.; Tilman, D. Resource availability controls fungal diversity across a plant diversity gradient. *Ecol. Lett.* 2006, 9, 1127–1135. [CrossRef] [PubMed]
- Prober, S.M.; Leff, J.W.; Bates, S.T.; Borer, E.T.; Firn, J.; Harpole, W.S.; Lind, E.M.; Seabloom, E.W.; Adler, P.B.; Bakker, J.D.; et al. Plant diversity predicts beta but not alpha diversity of soil microbes across grasslands worldwide. *Ecol. Lett.* 2015, *18*, 85–95. [CrossRef] [PubMed]
- 65. Nguyen, N.H.; Williams, L.J.; Vincent, J.B.; Stefanski, A.; Cavender-Bares, J.; Messier, C.; Paquette, A.; Gravel, D.; Reich, P.B.; Kennedy, P.G. Ectomycorrhizal fungal diversity and saprotrophic fungal diversity are linked to different tree community attributes in a field-based tree experiment. *Mol. Ecol.* **2016**, *25*, 4032–4046. [CrossRef]
- 66. Fierer, N.; McCain, C.M.; Meir, P.; Zimmermann, M.; Rapp, J.M.; Silman, M.R.; Knight, R. Microbes do not follow the elevational diversity patterns of plants and animals. *Ecology* **2011**, *92*, 797–804. [CrossRef]
- 67. Chen, Y.; Xu, T.; Fu, W.; Hu, Y.; Hu, H.; You, L.; Chen, B. Soil organic carbon and total nitrogen predict large-scale distribution of soil fungal communities in temperate and alpine shrub ecosystems. *Eur. J. Soil Biol.* **2021**, *102*, 103270. [CrossRef]
- Hooper, D.U.; Bignell, D.E.; Brown, V.K.; Brussard, L.; Dangerfield, J.M.; Wall, D.H.; Wardle, D.A.; Coleman, D.C.; Giller, K.E.; Lavelle, P.; et al. Interactions between aboveground and belowground biodiversity in terrestrial ecosystems: Patterns, mechanisms, and feedbacks. *BioScience* 2000, 50, 1049–1061. [CrossRef]
- Tedersoo, L.; Bahram, M.; Põlme, S.; Kõljalg, U.; Yorou, N.S.; Wijesundera, R.; Villarreal Ruiz, L.; Vasco-Palacios, A.M.; Thu, P.Q.; Suija, A.; et al. Fungal biogeography. Global diversity and geography of soil fungi. *Science* 2014, 346, 1256688. [CrossRef]
- Fierer, N. Embracing the unknown: Disentangling the complexities of the soil microbiome. *Nat. Rev. Microbiol.* 2017, 15, 579–590.
 [CrossRef]
- 71. de Vries, F.T.; Manning, P.; Tallowin, J.R.B.; Mortimer, S.R.; Pilgrim, E.S.; Harrison, K.A.; Hobbs, P.J.; Quirk, H.; Shipley, B.; Cornelissen, J.H.C.; et al. Abiotic drivers and plant traits explain landscape-scale patterns in soil microbial communities. *Ecol. Lett.* 2012, 15, 1230–1239. [CrossRef]
- 72. Liu, S.; Yu, H.; Yu, Y.; Huang, J.; Zhou, Z.; Zeng, J.; Chen, P.; Xiao, F.; He, Z.; Yan, Q. Ecological stability of microbial communities in Lake Donghu regulated by keystone taxa. *Ecol. Indic.* **2022**, *136*, 108695. [CrossRef]
- Banerjee, S.; Schlaeppi, K.; van der Heijden, M.G.A. Keystone taxa as drivers of microbiome structure and functioning. *Nat. Rev. Microbiol.* 2018, 16, 567–576. [CrossRef] [PubMed]
- 74. Pratt, R.G. Johnsongrass, yellow foxtail, and broadleaf signalgrass as new hosts for six species of *Bipolaris*, *Curvularia*, and *Exserohilum* pathogenic to bermudagrass. *Plant Dis.* **2006**, *90*, 528. [CrossRef] [PubMed]
- Kumar, J.; Schäfer, P.; Hückelhoven, R.; Langen, G.; Baltruschat, H.; Stein, E.; Nagarajan, S.; Kogel, K.-H. *Bipolaris sorokiniana*, a cereal pathogen of global concern: Cytological and molecular approaches towards better control. *Mol. Plant Pathol.* 2002, *3*, 185–195. [CrossRef] [PubMed]
- 76. Flory, S.L.; Kleczewski, N.; Clay, K. Ecological consequences of pathogen accumulation on an invasive grass. *Ecosphere* **2011**, 2, 120. [CrossRef]
- 77. Stricker, K.B.; Harmon, P.F.; Goss, E.M.; Clay, K.; Luke Flory, S. Emergence and accumulation of novel pathogens suppress an invasive species. *Ecol. Lett.* **2016**, *19*, 469–477. [CrossRef] [PubMed]
- Merzaeva, O.V.; Shirokikh, I.G. Colonization of plant rhizosphere by actinomycetes of different genera. *Microbiology* 2006, 75, 226–230. [CrossRef]
- 79. Kaushik, R.; Pandit, M.K.; Meyerson, L.A.; Chaudhari, D.S.; Sharma, M.; Dhotre, D.; Shouche, Y.S. Contrasting composition, diversity and predictive metabolic potential of the rhizobacterial microbiomes associated with native and invasive *Prosopis* congeners. *Curr. Microbiol.* **2021**, *78*, 2051–2060. [CrossRef]
- 80. Franco-Correa, M.; Quintana, A.; Duque, C.; Suarez, C.; Rodríguez, M.X.; Barea, J.-M. Evaluation of actinomycete strains for key traits related with plant growth promotion and mycorrhiza helping activities. *Appl. Soil Ecol.* **2010**, *45*, 209–217. [CrossRef]

- 81. Stefan, L.; Hartmann, M.; Engbersen, N.; Six, J.; Schöb, C. Positive effects of crop diversity on productivity driven by changes in soil microbial composition. *Front. Microbiol.* **2021**, *12*, 660749. [CrossRef] [PubMed]
- 82. Olanrewaju, O.S.; Babalola, O.O. Streptomyces: Implications and interactions in plant growth promotion. *Appl. Microbiol. Biotechnology* **2019**, *103*, 1179–1188. [CrossRef] [PubMed]
- 83. Li, Q.; Wan, F.; Zhao, M. Distinct soil microbial communities under Ageratina adenophora invasions. *Plant Biol.* **2022**, 24, 430–439. [CrossRef] [PubMed]