

Article

From Abundance to Extinction: Evolutionary History of European Aedemonini (Curculionidae) with a Description of the First Representative from Rovno Amber

Andrei A. Legalov ^{1,2,3,*} , Dmitry V. Vasilenko ^{4,5}  and Evgeny E. Perkovsky ^{6,7} 

¹ Institute of Systematics and Ecology of Animals, Siberian Branch, Russian Academy of Sciences, Novosibirsk 630091, Russia

² Department of Ecology, Biochemistry and Biotechnology, Altai State University, Barnaul 656049, Russia

³ Department of Forestry and Landscape Construction, Tomsk State University, Tomsk 634050, Russia

⁴ Laboratory of Arthropod, A. A. Borissiak Paleontological Institute of the Russian Academy of Sciences, Moscow 117647, Russia

⁵ Paleontological Laboratory, Cherepovets State University, Cherepovets 162600, Russia

⁶ Department of Entomology and Collection Management, I. I. Schmalhausen Institute of Zoology, National Academy of Sciences of Ukraine, 01030 Kiev, Ukraine

⁷ Natural History Museum of Denmark, 2100 Copenhagen, Denmark

* Correspondence: fossilweevils@gmail.com

Abstract: Fossil weevils of the subfamily Molytinae are currently represented by 103 species in 42 genera from 14 tribes. Fossil records of the tribe Aedemonini are known from the Eocene of Europe. This tribe makes up 22% of the London Clay weevil specimens identified to the genus level and 16% of the Curculionidae genera. At present, the distribution of this tribe is mainly paleotropical, but it was very prominent in the paratropical biota of the London Clay in the early Eocene, and it was still represented in the biota of the amber forests of Europe that retained some tropical elements in the Priabonian. A new species, *Electrorhinus vlaskini* n. sp., from the tribe Aedemonini of the subfamily Molytinae, is described from Rovno amber. It differs from *E. friedhelmi* in the smaller body size, finely faceted eyes, a coarser rugose pronotum, and sparser scales on the body. A list of the fossil Molytinae is compiled. This is the first record of a weevil of the subfamily Molytinae in Rovno amber, and the second finding of a representative of the genus *Electrorhinus* in the late Eocene and the third record of a species of Aedemonini preserved as a fossil. A key to European species of Molytinae with a rostral channel from Europe is given.

Keywords: Molytinae; Aedemonini; new species; faunistic relations; Eocene



Citation: Legalov, A.A.; Vasilenko, D.V.; Perkovsky, E.E. From Abundance to Extinction: Evolutionary History of European Aedemonini (Curculionidae) with a Description of the First Representative from Rovno Amber. *Diversity* **2023**, *15*, 376. <https://doi.org/10.3390/d15030376>

Academic Editors: Vyacheslav Martemyanov, Natalia Kirichenko and Luc Legal

Received: 14 January 2023

Revised: 28 February 2023

Accepted: 3 March 2023

Published: 5 March 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Some weevils of the subfamily Molytinae were previously considered the subfamily Cryptorhynchinae [1]. This subfamily included species from the tribe Cryptorhynchini along with representatives of the subtribes Cryptorhynchina, Mecistostylina, and Tylodina, tribes Aedemonini, Camptorhinini, Gasterocercini, Psepholacini, Sophrorhinini, and Torneumatini [1], or it was considered in an even wider sense with the inclusion of the tribes Cleogonini and Ithyporini [2]. Currently, a number of these tribes are included in the subfamily Molytinae [3–5].

The first finding of the tribe Aedemonini was in the London Clay (the early Eocene of England) [1,6], and the second from late Eocene Baltic amber [7].

Species of the tribe Cryptorhynchini are known from early Eocene Oise amber [8], the early Eocene Green River [9], the middle Eocene of Germany [10], late Eocene Baltic amber [11–13], the terminal Eocene of the USA [14], the latest Oligocene of Germany [15], and by many species from early Miocene Dominican and Mexican amber [16,17]. A single species of the extinct genus *Camptorrhinites* Britton, 1960, was described from London

Clay [6] as well. A species from the recent genus *Episcirrus* Kuschel, 1958, of the tribe Gasterocercini was discovered in Dominican amber [17]. Weevils from the tribes Psepholacini, Sophrorhinini, and Torneumatini are not known as fossils [11,18].

The family Curculionidae are represented in Rovno amber by 13 species from the subfamilies Eirrhiniinae, Dryophthorinae, Cossoninae, Conoderinae, Curculioninae, and Entiminae [19–28]. This is the first record of the subfamily Molytinae in Rovno amber, and the second finding of the late Eocene genus *Electrorhinus* and the fifth of the Aedemonini.

2. Materials and Methods

The studied fossil weevil is preserved in a piece of Rovno amber from the Pugach quarry (Klesov deposit), Sarny District, Rovno Oblast, Ukraine [29]. The type specimen is deposited in the Schmalhausen Institute of Zoology of the National Academy of Sciences of Ukraine, Kiev (SIZK). Images were taken using a Leica Z16 APO stereomicroscope equipped with a Leica DFC 450 camera and a Canon EOS 6D with an MP-E 65 mm macro lens and Cognisys Stackshot 3×. Photographs were stacked using the ZereneStacker 1.04 (Zerene Systems LLC, Richland, WA, USA). The morphological terminology used in this paper follows Lawrence et al. [30].

Fossil weevils of the subfamily Molytinae have been described from the following localities:

Cerro Guido—Chile: Magallanes, Última Esperanza Province, Cerro Guido, Dorotea Formation, Upper Cretaceous, Campanian;

Assiniboine—Canada: Manitoba, Assiniboine River; Upper Cretaceous, Maastrichtian;

Arkharo—Russia: Amurskaya Oblast, Arkharinskii District, quarry at Arkharo Railway Station lower Paleocene, Danian, Cagayan;

Meant—France: Puy-de-Dôme, middle-upper Paleocene, Selandian–Thanetian, 61.0–59.0 Ma;

Mors—Denmark: Fur Formation, lower Eocene, Ypresian, 54.0 Ma;

London Clay—UK: England, Sussex, Bognor Regis, lower Eocene, Ypresian, 54.0–50.0 Ma;

Roan Mountain—USA: Colorado, Garfield County, Green River Formation; Lower Eocene, Ypresian, 50.6–48 Ma;

Green River—USA: Colorado, Wyoming, Utah states, 3–4 km west of railway crossing of Green River, Green River Formation, lower Eocene, Ypresian, 50.6–48 Ma;

Oise amber—France: Paris basin, Creil, Oise; lowermost Eocene, Ypresian, 53.0 Ma;

Kutschlin—Czech Republic: northwest Bohemia, near Bilina, upper Eocene, Priabonian;

Baltic amber—Russia: Kaliningrad Oblast, Baltic Sea coast and Jantarnyi quarry near Kaliningrad, Poland: Gdańsk city area, at the Wisla River estuary, Prussian Formation, upper Eocene, Priabonian, 36.8–36.4 Ma;

Rovno amber—Ukraine, Rovno Oblast, Klesov, upper Eocene, Priabonian, 37.7–33.9 Ma;

Florissant—USA: Colorado, Rocky Mountains near Pike's Peak, Florissant Formation; uppermost Eocene, Priabonian, 34.07 ± 0.10 Ma;

White River Badlands—USA: South Dakota, lower Oligocene, Rupelian, 34.0–31.2 Ma;

Brunnstatt—France: Alsace, Haut-Rhine Dept., 5 km southwest of Mulhouse, terminal Eocene, Priabonian, 33.9 Ma;

Corent—France: South France, Puy-de-Dôme Department, Gergovia Plateau, south of Clermont-Ferrand, lower Oligocene, Rupelian;

Sieblös—Germany: Hessen, Rhön, lower Oligocene, Rupelian;

Lužice—Czech Republic: northwest Bohemia, upper Oligocene, Chattian;

Rott—Germany: North Rhine-Westphalia, latest Oligocene, Upper Siebengebirge, near Bonn, Rott Formation Chattian, 24.0–23.0 Ma;

Aix-en-Provence—France: Bouches-du-Rhône, Aix-en-Provence Formation uppermost Oligocene, upper Chattian, until 23 Ma;

Dominican amber—Dominican Republic: mines in the Cordillera Septentrional, lower-middle Miocene, 20–15 Ma;

Mexican amber—Mexico: Yucatan, Chiapas; mines near Simojovel, lower-middle Miocene, 20–15 Ma;

Shanwang—China: Shandong Province, Linqu County, lower Miocene, Burdigalian, 17–18 Ma;

Altamira shale—USA: California, Los Angeles Country, Woodcrest Drive, Sherman Oaks, upper Miocene;

Willershausen—Germany: Niedersachsen, Harz, upper Pliocene.

Nomenclatural acts introduced in the present work are registered in ZooBank (www.zoobank.org) under LSID urn:lsid:zoobank.org:pub:8931A219-B0E7-4F2A-85BD-0C29E9C48538.

3. Results

Systematic

Family **Curculionidae** Latreille, 1802

Subfamily **Molytinae** Schoenherr, 1823

Tribe **Aedemonini** Faust, 1898

Genus ***Electrorhinus*** Legalov, 2020

Type species: *Electrorhinus friedhelmi* Legalov, 2020, by original designation.

Electrorhinus vlaskini Legalov, Vasilenko and Perkovsky, **sp. n.** (Figures 1–4)




Figure 1. *Electrorhinus vlaskini* n. sp., holotype, habitus: (A) dorso-lateral view, left; (B) ventro-lateral view, right.




Figure 2. *Electrorhinus vlaskini* n. sp., holotype, habitus: (A) lateral view, left; (B) lateral view, right.



Figure 3. *Electrorhinus vlaskini* n. sp., holotype, habitus, frontal view.




Figure 4. *Electrorhinus vlaskini* n. sp., holotype: (A) dorso-lateral view; (B) carinae in postcoxal portions of the prosternum; arrow shows carinae.

LSIDurn:lsid:zoobank.org:act:76FEF9E3-38ED-47E8-9EFC-86BDBA382C57

Description. Size. Body length (without rostrum) 4.6 mm; rostrum length 0.8 mm. Body is black, with decumbent quite narrow scales. Rostrum moderately long and stout, slightly shorter than the pronotum, weakly curved, densely punctate dorsally; antennal scrobe lateral, directed under the rostrum; forehead quite narrow, slightly narrower than rostrum base, flattened dorsally, densely punctate; compound eyes large, spherical, weakly protruding from margin of head, quite finely faceted; temples short; vertex weakly convex, densely and finely punctate; antennae geniculate, inserted near middle of rostrum, geniculate; scape elongate, not reaching eye. Pronotum bell-shaped; disc weakly convex, coarsely rugose-punctate; sides weakly rounded, with large puncture; pronotal base biconcave. Elytra weakly convex, quite elongate, 3.1 times as long as pronotum; greatest width in middle; humeri weakly flattened; striae regular and distinct; elytral interstriae wide, punctate, 2.5–3.0 times as long as width of the striae. Prosternum densely punctate, with weak postocular lobes and a rostral channel bounded by carinae; procoxal cavities round, separated; postcoxal portion of prosternum bounded by carinae; metaventrite weakly convex, and coarsely punctate. Abdomen weakly convex. Legs elongate; femora weakly clavate, weakly sulcate beneath, with teeth beyond middle; metafemora reaching ventrite 3; tibiae weakly curved, flattened, with distinct uncus; tarsi elongate, with pulvilli on underside.

Material examined. Holotype: SIZK K-31315, Klesov, Rovno amber, late Eocene.

Derivation of name. The species is named in honor of paleoentomologist A.P. Vlsakin (Rovno).

Diagnosis. The new species differs from *Electrorhinus friedhelmi* in the smaller body sizes, finely faceted eyes, coarser rugose pronotum, and sparser scales on the body.

Remarks. The geniculate antennae and uncinata tibiae support the placement of this species in the family Curculionidae. Placement in the subfamily Molytinae is based on the quite large tibial uncus. The prosternum with rostral channel bounded by carinae in the precoxal and postcoxal portions of the prosternum (Figures 4 and 5) suggest placement in the tribe Aedemonini. The new species belongs to the genus *Electrorhinus* based on the femora weakly sulcate beneath and the metafemora reaching ventrite 3.




Figure 5. *Electrorhinus friedhelmi*, holotype, carinae in postcoxal portions of the prosternum; arrow shows carinae.

Key to European species of Molytinae with a rostral channel

1. Pronotal outgrowth extending beyond the head.2
- Pronotal outgrowth not extending beyond the head.....3
2. Apex of rostral channel bounded by carina. 4.9 mm. London Clay.....
-*Korystina gracilis* Britton, 1960.
- Apex of rostral channel not bounded by carina. 4.2 mm. Oise amber.....
-*Sciabregma squamosa* Legalov, Kirejtshuk, and Nel, 2019.
3. Rostral channel not reaching mesoventrite. ~5 mm. London Clay.....
-*Camptorrhinites orarius* Britton, 1960
- Rostral channel reaching mesoventrite.....4
4. Postcoxal portion of prosternum bounded by carinae.....5
- Postcoxal portion of prosternum not bounded by carinae.....8
5. Pronotum is densely punctate. Body is smaller (4.4 mm). London Clay.....
-*Taylorius litoralis* Britton, 1960.
- Pronotum rugose-punctate. Body is larger (4.6–6.2 mm).....6
7. Body is larger (5.7–6.2 mm). Eyes are quite coarsely faceted. Pronotum is weakly rugose. Body with denser scales. Baltic amber.....*Electrorhinus friedhelmi* Legalov, 2020.
- Body is smaller (4.6 mm). Eyes are finely faceted. Pronotum is coarser and rugose. Body with sparser scales. Rovno amber.....*Electrorhinus vlaskini* Legalov, Vasilenko, and Perkovsky, sp. n.
8. Metaventricle is longer than the third ventrite. Abdominal process is narrower than coxa. Elytra with rectangular humeri.....9

- Metaventrite is short, and shorter than the third ventrite. Abdominal process as broad as coxa. Elytra with reduced humeri.....10
- 9. Vertex is densely punctured and covered with wide scales. Elytra wider than base of pronotum. Odd-numbered elytral interstices are non-raised. Elytra is 1.8 times as long as the width at the middle. 4.5–4.7 mm. Baltic amber.....*S. silvestris* Legalov, in lit.
- Vertex glabrous, covered with narrow scales. Elytra subequal to base of pronotum. Odd-numbered elytral interstices are weakly raised. Elytra are 2.6 times as long as the width at the middle. 4.0 mm. Baltic amber.....*Succinacalles unicus* Zherikhin, 1971
- 10. Femora denticulate. 5.2 mm. Baltic amber.....*Baltacalles triumurbium* Bukejs, Alekseev et Legalov, 2020.
- Femora without teeth.....11
- 11. Postocular lobes are weak. Elytral interstices are rather narrow. Elytral striae are wide. 3.0 mm. Baltic amber.....*Parvacalles kotthoffi* Legalov, in lit.
- Postocular lobes are stark. Elytral interstices are wide. Elytral striae are narrow. 3.3 mm. Oise amber.....*Oisecalles latosquamosus* Legalov, Kirejtshuk et Nel, 2019.

List of the fossil Molytinae

- Subfamily Molytinae Schoenherr, 1823
- Tribe Molytini Schoenherr, 1823
- Genus *Archaralites* Legalov, 2010
- A. zherichini* Legalov, 2010—Arkhar [31]
- Genus *Archaeoheilus* Legalov, 2018
- A. gallicus* Legalov, Kirejtshuk and Nel, 2019—Menat [32]
- A. ovalis* Legalov, 2018—Green River [33]
- A. packardii* (Scudder, 1893)—Green River [14]
- A. proectus* (Scudder, 1876)—Green River [9]
- A. scudderi* Legalov, 2018—Green River [33]
- A. deleticius* (Scudder, 1893)—White River [14]
- A. lacoiei* (Scudder, 1893)—Florissant [14]
- Genus *Furhylobius* Legalov, 2015
- F. troesteri* Legalov, 2015—Mors [11]
- Genus *Hylobius* Germar, 1817
- "H." antiquus* Heyden and Heyden, 1866—Rott [15]
- "H." medianus* Zhang, 1989—Shanwang [34]
- "H." plenus* Zhang, 1989—Shanwang [34]
- "H." turfatus* Zhang, 1989—Shanwang [34]
- H. abietis* (Linnaeus, 1758)—Willershausen [35]
- Genus *Anisorhynchus* Schoenherr, 1842
- "A." deletus* Deichmueller, 1881—Kutschlin [36]
- "A." offusus* Oustalet, 1870—Corent [37]
- Genus *Sinocurculia* Hong, 1984
- S. granulata* Hong, 1984—Shanwang [38]
- S. shanwangensis* Hong, 1985—Shanwang [39]
- Tribe Sciabregmini Legalov, Kirejtshuk, and Nel, 2019
- Genus *Sciabregma* Scudder, 1893
- S. rugosa* Scudder, 1893—Roan Mountain [14]
- S. squamosa* Legalov, Kirejtshuk, and Nel, 2019—Oise amber [8]
- S. tenuicornis* Cockerell, 1921—Green River [40]
- Tribe Conotrachelini Jakel, 1865
- Genus *Conotrachelus* Schoenherr, 1845
- C. florissantensis* Wickham, 1912—Florissant [41]
- Tribe Acicnemidini Lacordaire, 1866
- Genus *Electrotribus* Hustache, 1942
- E. rarus* Legalov, 2020—Baltic amber [7]

E. theryi Hustache, 1942—Baltic amber [42,43]
 =*Pissodes henningseni* Voss, 1972
E. weigangae (Ulke, 1947)—Baltic amber [44]
 =*Anchorthorrhinus incertus* Voss, 1953 [45]
 =*Isalcidodes macellus* Voss, 1953
E. wolfschwenningerae (Rheinheimer, 2007)—Baltic amber [46]
 Tribe Plinthini Lacordaire, 1863
 Subtribe Leiosomatina Reitter, 1913
 Genus *Leiosoma* Stephens, 1829
 Subgenus *Palaeoleiosoma* Legalov, 2016
L. (L.) klebsi Legalov, 2016—Baltic amber [47]
L. (L.) michalski Legalov, in lit.—Baltic amber
L. (L.) shevnini Legalov, in lit.—Baltic amber
 Subtribe Typoderina Voss, 1965
 Genus *Baltanchonidium* Legalov, in lit.
B. eocenicum Legalov, in lit.—Baltic amber
 Tribe Magdalini Pascoe, 1870
 Genus *Magdalis* Germar, 1817
 “*M.*” *sedimentorum* Scudder, 1893—Florissant [14]
M. moesta Schlechtendal, 1894—Rott [48]
 “*M.*” *deucalionis* (Heyden and Heyden, 1866)—Rott [15]
 “*M.*” *protogenius* (Heyden and Heyden, 1866)—Rott [15]
 Tribe Cleogonini Gistel, 1856
 Genus *Rhysosternum* Scudder, 1893
Rh. aeternabile Scudder, 1893—Florissant [14]
Rh. longirostre Scudder, 1893—Florissant [14]
 Genus *Rhyssomatus* Schoenherr, 1837
Rh. miocenae Pierce, 1965—Altamira shale [49]
 Tribe Anchonini Imhoff, 1856
 Genus *Anchonus* Schoenherr, 1825
A. acrolepidotus Poinar and Legalov, 2019—Dominican amber [50]
A. bothynus Poinar and Legalov, 2019—Dominican amber [50]
 Genus *Velatis* Poinar and Voisin, 2003
V. dominicana Poinar and Voisin, 2003—Dominican amber [51]
 Tribe Pissodini Gistel, 1856
 Genus *Lithopissodes* Beier, 1952
L. lushtzensis Beier, 1952—Lužice [52]
 Tribe Lymantini Lacordaire, 1865
 Subtribe Lymantina Lacordaire, 1865
 Genus *Bronchotibia* Poinar and Legalov, 2022
B. adunatus Poinar and Legalov, 2022—Dominican amber [53]
 Tribe incertae sedis
 Genus *Hylobiites* Scudder, 1895
H. cretaceus Scudder, 1895—Assiniboine [54]
 Genus *Dorotheus* Kuschel, 1959
D. guidensis Kuschel, 1959—Cerro Guido [55]
 Genus *Lutago* Britton, 1960
L. fetusus Britton, 1960—London Clay [6]
L. nanus Britton, 1960—London Clay [6]
 Genus *Pissodites* Britton, 1960
P. argillosus Britton, 1960—London Clay [6]
 Genus *Erirhinites* Britton, 1960
E. bognoensis Britton, 1960—London Clay [6]
 Genus incertae sedis

"Chalcodermus" kirschi Deichmueller, 1881—Kutschlin [36]

"Acalles" exhumatus Wickham, 1913—Florissant [56]

"Hylobius" carbo Oustalet, 1874—Aix-en-Provence [37]

"Plinthus" heerii Oustalet, 1874—Aix-en-Provence [37]

"Molytes" hassencampi Heyden, 1858—Sieblos [57]

"Pissodes" effossus Heyden, 1858—Sieblos [57]

"Pissodes" planatus Foerster, 1891—Brunnstatt [58]

"Hylobius" deletus Oustalet, 1870—Corent [59]

"Plinthus" redivivus Oustalet, 1870—Corent [59]

"Hylobius" morosus Oustalet, 1874—Corent [37]

"Cryptorhynchinae Schoenherr, 1825"

Tribe Camptorhinini Lacordaire, 1866

Genus *Camptorrhinites* Britton, 1960

C. orarius Britton, 1960—London Clay [6]

Genus *Korystina* Britton, 1960

Remarks. The genus *Korystina* was described in the tribe Sophrorhinini [6]. Alonso-Zarazaga and Lyal [1] placed it in the tribe Cryptorhynchini. Legalov [11] transferred this genus in the tribe Camptorhinini. A re-examination of the description and illustrations show that *Korystina* has the rostral channel of the postcoxal portion of the prosternum bounded by carinae, which characterize the tribe Aedemonini [5], but it is not clear whether it is present on the prothorax or is also located on the mesoventrite. To finally establish the systematic position of *Korystina*, a re-examination of the type is required.

K. gracilis Britton, 1960—London Clay [6]

Tribe Aedemonini Faust, 1898

Genus *Taylorius* Britton, 1960

T. litoralis Britton, 1960—London Clay [6]

Genus *Electrorhinus* Legalov, 2020

E. friedhelmi Legalov, 2020—Baltic amber [7]

E. vlaskini Legalov, Vasilenko and Perkovsky, sp. n.—Rovno amber

Tribe Cryptorhynchini Schoenherr, 1826

Subtribe Cryptorhynchina Schoenherr, 1826

Genus *Succinacalles* Zherikhin, 1971

S. silvestris Legalov, in lit.—Baltic amber

S. unicus Zherikhin, 1971—Baltic amber [12]

Genus *Cryptorhynchus* Illiger, 1807

"C." annosus Scudder, 1876—Green River [9]

"C." kerri Scudder, 1893—Florissant [14]

"C." profusus Scudder, 1893—Florissant [14]

Genus *Semnorhynchus* Faust, 1896

S. brachyrhinus Poinar and Legalov, 2014—Dominican amber [17]

S. camptostegus Poinar and Legalov, 2014—Dominican amber [17]

S. contorhinus Poinar and Legalov, 2014—Dominican amber [17]

S. euryaspus Poinar and Legalov, 2014—Dominican amber [17]

S. eurystegus Poinar and Legalov, 2014—Dominican amber [17]

S. leptostegus Poinar and Legalov, 2014—Dominican amber [17]

S. megasomus Poinar and Legalov, 2014—Dominican amber [17]

S. stenostegus Poinar and Legalov, 2014—Dominican amber [17]

S. tanyrhinus Poinar and Legalov, 2014—Dominican amber [17]

Genus *Neoulosomus* O'Brien and Wibmer, 1982

Subgenus *Neoulosomus* s. str.

N. (N.) hurdi (Zimmermann, 1971)—Mexican amber [16]

N. (N.) leptosomus Poinar and Legalov, 2014—Dominican amber [17]

N. (N.) megaholcus Poinar and Legalov, 2014—Dominican amber [17]

N. (N.) megus Poinar and Legalov, 2014—Dominican amber [17]

N. (N.) microholcus Poinar and Legalov, 2014—Dominican amber [17]
N. (N.) pedinus Poinar and Legalov, 2014—Dominican amber [17]
N. (N.) pediosomus Poinar and Legalov, 2014—Dominican amber [17]
N. (N.) platystegus Poinar and Legalov, 2014—Dominican amber [17]
N. (N.) scamposomus Poinar and Legalov, 2014—Dominican amber [17]
N. (N.) scambus Poinar and Legalov, 2014—Dominican amber [17]
N. (N.) stenocalypus Poinar and Legalov, 2014—Dominican amber [17]
N. (N.) stylolepus Poinar and Legalov, 2014—Dominican amber [17]
 Subgenus *Stenosomus* Poinar and Legalov, 2014
N. (S.) contorhinus Poinar and Legalov, 2014—Dominican amber [17]
N. (S.) tanyrhinus Poinar and Legalov, 2014—Dominican amber [17]
 Genus *Lemmasomus* Poinar and Legalov, 2014
L. anodontotus Poinar and Legalov, 2014—Dominican amber [17]
 Genus *Apharosoma* Poinar and Legalov, 2014
A. euryrhina Poinar and Legalov, 2014—Dominican amber [17]
 Genus *Anlemmus* Poinar and Legalov, 2014
A. leptorhinus Poinar and Legalov, 2014—Dominican amber [17]
 Genus *Odontamera* Poinar and Legalov, 2014
O. dolichosoma Poinar and Legalov, 2014—Dominican amber [17]
 Subtribe Tylodina Lacordaire, 1866
 Genus *Oisecalles* Legalov, Kirejtshuk et Nel, 2019
O. latosquamosus Legalov, Kirejtshuk et Nel, 2019—Oise amber [8]
 Genus *Baltacalles* Bukejs, Alekseev et Legalov, 2020
B. triumurbium Bukejs, Alekseev et Legalov, 2020—Baltic amber [13]
 Genus *Parvacalles* Legalov, in lit.
P. kotthoffi Legalov, in lit.—Baltic amber
 Genus *Acalles* Schoenherr, 1826
 “*A. icarus* Heyden and Heyden, 1866—Rott [15]
 Genus *Pseudomoides* Poinar and Legalov, 2014
P. clisaulis Poinar and Legalov, 2014—Dominican amber [17]
 Genus *Paraulosomus* Hustache, 1930
P. adenolepus Poinar and Legalov, 2014—Dominican amber [17]
 Genus *Paracamptopsis* Poinar and Legalov, 2014
P. stenis Poinar and Legalov, 2014—Dominican amber [17]
 Tribe Gasterocercini Zherichin, 1991
 Genus *Episcirrus* Kuschel, 1958
E. isolepus Poinar and Legalov, 2014—Dominican amber [17]

4. Discussion

Fossil weevils of the subfamily Molytinae are currently represented by 103 species in 42 genera from 14 tribes. The oldest forms were described from the Maastrichtian of Canada and Chile [54,55]. The most diverse Molytinae fauna is known from Miocene Dominican amber [17,50,51,53].

The tribe Aedemonini includes 33–35 extant genera [1,3] distributed in sub-Saharan Africa (excluding the desert regions of South Africa), Madagascar, Oriental Region (north to southern Sakhalin, Japan, and South Korea), Australia (except for desert areas), Micronesia, Melanesia, and Polynesia (Figure 6). Representatives of this group are absent in the New World, Western and Central Palaearctic. Fossil findings are known from the Eocene of Europe [6,7,18]. It is noted that extant species of the tribe avoid dry and cold areas. Almost all species do not cross the line of +15 °C of the mean annual temperature. Several species have adapted to colder conditions in South Korea, Japan, and southern Sakhalin. Perhaps this is due to the monsoon climate. The fossil records are all from the Eocene when the climate of Europe was warm and humid. Aedemonini make up 22% of the London Clay weevil specimens identified to the genus level, and 16% (or even a third if *Korystina*

belongs to the tribe) of the Curculionidae genera. The representation of the tribe was not so significant in any other extant or extinct fauna. Even in the Afrotropical region, where at least 40% of the species and 56% of the genera of Aedemonini are distributed, they make up less than 7% of the species and less than 2.5% of the genera of Curculionidae [60]. It is highly meaningful that hyperdiverse Cryptorhynchini appear to be largely absent from the Afrotropics, where they seem to be replaced by Aedemonini [61], our data. It is interesting to combine the abundance of Aedemonini and Molytinae in general in London Clay with the extreme rarity of bark beetles (represented by a single specimen of ? *Blastophagus* Eichhoff, 1864 (= *Tomicus* Latreille, 1802)), while xylophagous insects dominate in this locality [6]. Their mineralization, at least in part, could occur in a specific microtaphotope inside pyritized wood [62].




Figure 6. Distribution of the species of the tribe Aedemonini: Recent members—green shaded area; red circle—fossil record from Rovno amber; blue circle—fossil record from Baltic amber; pink circle—fossil record from London Clay.

Moreover, *Taylorius* is the second most abundant genus of the family Curculionidae (together with *Lutago*, they account for 62.5% of all weevils determined to be in the genus), and the third most abundant genus in all biotas, and accounts for 16% of all beetles identified to the genus from London Clay. Of course, an extraordinary abundance of Aedemonini in the early Eocene of Europe was not ubiquitous. For example, the species of the tribe are unknown from Oise amber, in which numerous bark beetles were found, as well as a representative of Cryptorhynchini (*Oisecalles latosquamosus* Legalov, Kirejtshuk et Nel, 2019) [8].

The abundance of Aedemonini in the London Clay may have been due to the rich and diverse tropical vegetation [63,64]. The age of the London Clay Formation is early and middle Ypresian [65]. Bognor remains the principal international source of pyritized insects [66] (p. 93). Rundle and Cooper [67] asserted that insects from the London Clay drifted out to sea with wood from the more westerly, near-shore location of the Hampshire Basin [68]. This is further supported by the presence of numerous flightless larvae [66], together with beetles with their elytra in the resting position [68]. The London Clay Formation contains one of the most important fruit and seed assemblages from the Paleogene [69]. Given its unparalleled diversity [68], more than 350 described species, and superb preservation as well as uncompressed, three-dimensionally preserved morphology [63,70], the flora offers a unique insight into the vegetation present in Europe during the hyperthermals of the

early Eocene [71,72]. Numerous clades that are now largely confined to tropical latitudes, e.g., Annonaceae, Arecaceae, Burseraceae, Icacinaceae, and Menispermaceae, are well documented in the flora [63,70]. As was indicated by Collinson and Cleal [68] “in early Eocene times, a zone of lush vegetation had developed in Britain, strongly reminiscent of today’s tropical rain forests, dominated by evergreen trees of the sumac, custard apple, dillenia, dogbane, frankincense, flacourtia, icacina, laurel, palm, sabia, soap berry and tea families, and mastic trees of the dogwood family”, with abundant lianas of the icacina, grape vine, and moonseed families and numerous mangrove palms and rare, true mangroves in coastal areas [73–76]. The vegetation, with some reservations, has been compared with the modern paratropical rain forests of lowland Asia [68], and references therein.

Since the late Eocene, there are no fossil records of these Aedemonini beetles [18]. They may have disappeared from Europe during the Oligocene, when the European climate became colder, seasonal, and drier. The new record shows the connection of the European Priabonian fauna with the extant Paleotropical fauna. The weaver ant genus *Oecophylla* Smith, 1860, is characterized by a similar geological and geographical distribution [77,78], except for a single early Eocene finding in Okanagan [79].

The climate of the Rovno amber forest was warmer than the Baltic one [80,81], so the thermophilic beetles are much better represented there [27,82–89], resulting in only 15% of known Rovno beetles having anything in common with Baltic amber [80,87]. The genus *Electrorhinus* could be a new addition to the list of Rovno thermophilic taxa that are much rarer in Baltic amber.

Author Contributions: A.A.L. and E.E.P. designed the study, prepared new species descriptions and plates. A.A.L. prepared the systematic placement of the new species. A.A.L., D.V.V. and E.E.P. authors drafted the manuscript and contributed to the writing and discussion. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Data Availability Statement: The specimens are deposited in the I. I. Schmalhausen Institute of Zoology, National Academy of Sciences of Ukraine (Kiev).

Acknowledgments: The authors thank A.P. Vlaskin (SIZK) for finding, cutting, and polishing the sample and S.A. Simutnik (SIZK) and J.M.J. Shaw (Zoological Museum, Natural History Museum of Denmark, Copenhagen, Denmark) for help with the photos and editing, and anonymous reviewers for improving the overall quality of the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Alonso-Zarazaga, M.A.; Lyal, C.H.C. *A World Catalogue of Families and Genera Curculionoidea (Insecta: Coleoptera) (Excepting Scolytidae and Platypodidae)*; Entomopraxis: Barcelona, Spain, 1999; p. 315.
2. Morimoto, K. On the genera of Oriental Cryptorhynchinae (Coleoptera: Curculionidae). *Esakia* **1978**, *11*, 121–143. [[CrossRef](#)] [[PubMed](#)]
3. Lyal, C.H.C. 3.7.7. Molytinae Schoenherr, 1823. In *Arthropoda: Insecta. Handbook of Zoology. Tb. 40: Coleoptera (Beetles). Vol. 3: Morphology and Systematics (Phytophaga)*; Leschen, R.A.B., Beutel, R.G., Eds.; Walter de Gruyter: Berlin, Germany; New York, NY, USA, 2014; pp. 529–569.
4. Alonso-Zarazaga, M.A.; Barrios, H.; Borovec, R.; Bouchard, P.; Caldara, R.; Colonnelli, E.; Gültekin, L.; Hlavá, P.; Korotyaev, B.; Lyal, C.H.C.; et al. Cooperative catalogue of Palaearctic Coleoptera Curculionoidea. *Monogr. Electrónicas* **2017**, *8*, 1–729.
5. Legalov, A.A. Annotated key to weevils of the world. Part 2. Subfamily Molytinae (Coleoptera, Curculionidae). *Ukr. J. Ecol.* **2018**, *8*, 340–350.
6. Britton, E.B. Beetles from the London clay (Eocene) of Bognor Regis, Sussex. *Bull. Brit. Mus. Nat. Hist. Geol.* **1960**, *4*, 27–50, +pl. 2–7. [[CrossRef](#)]
7. Legalov, A.A. A review of the Curculionoidea (Coleoptera) from European Eocene ambers. *Geosciences* **2020**, *10*, 16. [[CrossRef](#)]
8. Legalov, A.A.; Kirejtshuk, A.G.; Nel, A. New weevils (Coleoptera, Curculionoidea) from the earliest Eocene Oise amber. *Paleontol. J.* **2019**, *53*, 729–751. [[CrossRef](#)]
9. Scudder, S.H. Fossil Coleoptera from the Rocky mountain territories. *Bull. US Geol. Geogr. Surv. Terr.* **1876**, *2*, 77–87.

10. Wappler, T. Die Insekten aus dem Mittel-Eozän des Eckfelder Maares. *Vulkaneifel Mainzer Naturwiss. Arch.* **2003**, *27*, 1–234.
11. Legalov, A.A. Fossil Mesozoic and Cenozoic weevils (Coleoptera, Obrienioidea, Curculionoidea). *Paleontol. J.* **2015**, *49*, 1442–1513. [\[CrossRef\]](#)
12. Zherikhin, V.V. On weevils (Insecta, Coleoptera) from the Baltic amber. *Tr. Paleontol. Inst. Akad. Nauk. SSSR* **1971**, *130*, 197–209.
13. Bukejs, A.; Alekseev, V.I.; Legalov, A.A. A new Eocene genus of the subtribe Tylodina (Coleoptera: Curculionidae) and notes concerning local differences of Baltic amber in the Kaliningrad Region. *Foss. Rec.* **2020**, *23*, 75–81. [\[CrossRef\]](#)
14. Scudder, S.H. Rhynchophorus Coleoptera of the United States. *Monogr. US Geol. Surv. Tert.* **1893**, *21*, 1–206.
15. Heyden, C.; Heyden, L. Käfer und Polypen aus der Braunkohle des Siebengebirges. *Palaeontographica* **1866**, *15*, 131–156, pl. XXII–XXIV.
16. Zimmermann, E.C. Mexican Miocene amber weevils (Insecta: Coleoptera: Curculionidae). *Univ. Calif. Publ. Entomol.* **1971**, *63*, 103–106.
17. Poinar, G., Jr.; Legalov, A.A. New Cryptorhynchinae (Coleoptera: Curculionidae) in Dominican amber. *Hist. Biol.* **2014**, *26*, 502–534. [\[CrossRef\]](#)
18. Legalov, A.A. Fossil history of Curculionoidea (Coleoptera) from the Paleogene. *Geosciences* **2020**, *10*, 358. [\[CrossRef\]](#)
19. Legalov, A.A.; Nazarenko, V.Y.; Perkovsky, E.E. A new species of the genus Glaesotropis Gratshev and Zherikhin, 1995 (Coleoptera, Anthribidae) from Rovno amber. *Fossil Record* **2021**, *24*, 1–7. [\[CrossRef\]](#)
20. Nazarenko, V.Y.; Perkovsky, E.E. A new genus and species of dryophthorid weevils (Coleoptera, Dryophthoridae: Stromboscerinae) from the Rovno amber. *Paleontol. J.* **2009**, *43*, 1097–1100. [\[CrossRef\]](#)
21. Nazarenko, V.Y.; Perkovsky, E.E. A new species of derelomine weevils (Coleoptera, Curculionidae, Curculioninae: Acalyptini) from the Rovno amber. *Paleontol. J.* **2016**, *50*, 991–996. [\[CrossRef\]](#)
22. Nazarenko, V.Y.; Legalov, A.A.; Perkovsky, E.E. A new species of the genus Caulophilus Woll. (Coleoptera: Curculionidae: Cossoninae) from the Rovno amber. *Paleontol. J.* **2011**, *45*, 287–290. [\[CrossRef\]](#)
23. Legalov, A.A.; Nazarenko, V.Y.; Perkovsky, E.E. New weevils (Coleoptera: Curculionidae) from Rovno amber. *Paleontol. J.* **2019**, *53*, 1045–1059. [\[CrossRef\]](#)
24. Legalov, A.A.; Nazarenko, V.Y.; Perkovsky, E.E. A new species of the genus Dorytomus Germar, 1817 (Coleoptera, Curculionidae) from Rovno amber. *Zootaxa* **2021**, *5006*, 95–100. [\[CrossRef\]](#)
25. Legalov, A.A.; Nazarenko, V.Y.; Vasilenko, D.V.; Perkovsky, E.E. Ceutorhynchus Germar (Coleoptera, Curculionidae) as proxy for Eocene Brassicaceae: First record of the genus from Rovno amber. *J. Paleontol.* **2022**, *96*, 379–386. [\[CrossRef\]](#)
26. Legalov, A.A.; Vasilenko, D.V.; Perkovsky, E.E. The American tribes Anypotactini and Eudiagogini (Coleoptera, Curculionidae) in Eocene of Europe as indicators of Eocene climate with description a new species. *Diversity* **2022**, *14*, 767. [\[CrossRef\]](#)
27. Legalov, A.A.; Vasilenko, D.V.; Perkovsky, E.E. New proxy for Moraceae in Priabonian of Europe: First record of the genus Demimaea Pascoe, 1870 (Coleoptera: Curculionidae) from Eocene Rovno amber. *Hist. Biol.* **2022**. [\[CrossRef\]](#)
28. Bukejs, A.; Legalov, A.A. First record of the tribe Naupactini (Coleoptera: Curculionidae) in Rovno amber. *Foss. Rec.* **2019**, *22*, 25–30. [\[CrossRef\]](#)
29. Makarkin, V.N.; Perkovsky, E.E.; Anisyutkin, L.N.; Dubovikoff, D.A. First larvae of Raphidioptera from Eocene Sakhalinian and Rovno ambers. *Zootaxa* **2022**, *5219*, 456–466. [\[CrossRef\]](#)
30. Lawrence, J.F.; Beutel, R.G.; Leschen, R.A.B.; Slipinsky, S.A. Chapter 2. Glossary of morphological terms. In *Handbook of Zoology. Arthropoda: Insecta. Coleoptera, Beetles. Volume 2: Morphology and Systematics (Elateroidea, Bostrichiformia, Cucujiformia partim)*; Kristensen, N.P., Beutel, R.G., Eds.; Walter de Gruyter: Berlin, Germany; New York, NY, USA, 2010; pp. 9–20.
31. Legalov, A.A. Checklist of Mesozoic Curculionoidea (Coleoptera) with description of new taxa. *Balt. J. Coleopterol.* **2010**, *10*, 71–101.
32. Legalov, A.A.; Kirejtshuk, A.G.; Nel, A. A new species of the genus Archaeoheilus Legalov, 2018 (Coleoptera: Curculionidae) from the Paleocene of Menat (France). *Comptes Rendus Palevol* **2022**, *21*, 245–251. [\[CrossRef\]](#)
33. Legalov, A.A. New weevils (Coleoptera, Curculionoidea) from the Eocene of the Green River, United States: Part 1. *Paleontol. J.* **2018**, *52*, 294–302. [\[CrossRef\]](#)
34. Zhang, J. *Fossil Insects from Shanwang, Shandong, China*; Shandong Science and Technology Publication: Jinan, China, 1989; p. 459.
35. Gersdorf, E. Weitere Käfer (Coleoptera) aus dem Jungtertiär Norddeutschlands. *Geol. Jahrb.* **1971**, *88*, 629–670.
36. Deichmueller, J.V. Fossile Insecten aus dem Diatomeenschiefer von Kutschlin bei Bilin, Böhmen. *Nova Acta Leopold. Verh. Kais. Leopold. Carol. Dtsch. Akad. Nat.* **1881**, *42*, 293–331.
37. Oustalet, M.E. Recherches sur les insectes fossiles des terrains tertiaires de la France. 2me partie. Insectes fossiles d’Aix en Provence. *Ann. Sci. Géol. Paris* **1874**, *5*, 1–347.
38. Hong, Y. Fossil insects in the diatoms of Shanwang. *Bull. Tianjin Inst. Geol. Mineral. Res.* **1984**, *8*, 1–11.
39. Hong, Y. *Fossil Insects, Scorpions and Aranaeids in the Diatoms of Shandong*; Geological Publishing House: Peking, China, 1985; p. 80.
40. Cockerell, T.D.A. Some Eocene insects from Colorado and Wyoming. *Proc. Unit. Stat. Nat. Mus.* **1921**, *59*, 29–39.
41. Wickham, H.F. On some fossil Rhynchophorus Coleoptera from Florissant Colorado. *Bull. Am. Mus. Nat. Hist.* **1912**, *31*, 41–55, I–IV.
42. Hustache, A. Un curculionide de l’ambre de la Baltique. *Bull. Mens. Soc. Lin. Lyon.* **1942**, *11*, 108–109. [\[CrossRef\]](#)
43. Voss, E. Einige Rüsselkäfer der Tertiärzeit aus baltischen Bernstein (Coleoptera, Curculionidae). *Steenstrupia* **1972**, *2*, 167–181.
44. Ulke, T. A new genus and species of Curculionidae (Coleoptera) in Baltic amber. *Not. Nat.* **1947**, *19*, 1–5.

45. Voss, E. Einige Rhynchophoren der Bernsteinfauna (Col.). *Mitt. Geol. Staatsinst. Hambg.* **1953**, *22*, 119–140.
46. Rheinheimer, J. Neue fossile Rüsselkäfer (Coleoptera: Curculionidae) aus dem Eozän des Baltischen Bernsteins und der Grube Messel bei Darmstadt. *Staatliches. Mus. Nat.* **2007**, *365*, 1–24.
47. Legalov, A.A. New weevils (Curculionidae) in Baltic amber. *Paleontol. J.* **2016**, *50*, 970–985. [[CrossRef](#)]
48. Schlechtendal, D.H.R. Beiträge zur Kenntnis Fossiler Insekten aus dem Braunkohlengebirge von Rott am Siebengebirge. *Abh. Nat. Gesellsch. Halle* **1894**, *20*, 197–228, XII–XIV.
49. Pierce, W.D. Fossil arthropods of California. 26. Three new fossil insects sites in California. *Bull. South. Calif. Acad. Sci.* **1965**, *64*, 151–152.
50. Poinar, G., Jr.; Legalov, A.A. Two new species of the genus *Anchonus* Schoenherr, 1825 (Coleoptera: Curculionidae: Molytinae) in Dominican amber. *Palaeontol. Electron.* **2019**, *22*, 1–10. [[CrossRef](#)]
51. Poinar, G.O., Jr.; Voisin, J.F. A Dominican amber weevil, *Velatis dominicana* gen. n., sp. n. and key to the genera of the Anchonini (Molytinae, Curculionidae). *Nouv. Rev. Entomol.* **2003**, *19*, 373–381.
52. Beier, M. Miozäne und oligozäne Insekten aus Österreich und den unmittelbar angrenzenden Gebieten. In *Sitzungsberichte Österreichische Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse, Abteilung 1: Biologie, Mineralogie, Erdkunde und Verwandte Wissenschaften*; Springer: Berlin/Heidelberg, Germany, 1952; Volume 161, pp. 129–134.
53. Poinar, G.; Legalov, A.A. First record of the tribe Lymantini (Coleoptera: Curculionidae) from Dominican amber. *Hist. Biol.* **2022**, *34*, 67–71. [[CrossRef](#)]
54. Scudder, S.H. Canadian fossil insects: 2. The Coleoptera hitherto found fossil in Canada. *Contrib. Canad. Palaeontol. Geol. Surv. Can.* **1895**, *2*, 1–57.
55. Kuschel, G. Un Curculiónido del Cretácico Superior, primer insecto fósil de Chile. *Invest. Zool. Chil.* **1959**, *5*, 49–54.
56. Wickham, H.F. Fossil Coleoptera from the Wilson Ranch near Florissant, Colorado. *Bull. Lab. Nat. Hist. State. Univ. Iowa* **1913**, *6*, 3–29, I–VII.
57. Heyden, C.H.G. Fossile Insekten aus der Braunkohle der Sieblos. *Palaeontology* **1858**, *5*, 115–120.
58. Foerster, B. Die Insekten des ‘Plattigen Steinmergels’ von Brunstatt. *Abh. Geol. Spec. Elsass Lothringen* **1891**, *3*, 333–594, XI–XVI.
59. Oustalet, F.E. Recherches sur les insectes fossiles des terrains tertiaires de la France. *Ann. Sci. Géol. Paris* **1870**, *2*, 1–178.
60. O’Brien, C.W.; Wibmer, G.J. Numbers of genera and species of Curculionidae (Coleoptera). *Entomol. News* **1978**, *89*, 89–92.
61. Letsch, H.; Balke, M.; Toussaint, E.F.A.; Riedel, A. Historical biogeography of the hyperdiverse hidden snout weevils (Coleoptera, Curculionidae, Cryptorhynchinae). *Syst. Entomol.* **2019**, *45*, 312–326. [[CrossRef](#)]
62. Zherikhin, V.V.; Ponomarenko, A.G.; Rasnitsyn, A.P. *Introduction into Palaeoentomology*; KMK Press: Moscow, Russia, 2008; p. 371.
63. Chandler, M.E.J. *Paleocene Flora: The Lower Tertiary Floras of Southern England. I. Palaeocene Floras, London Clay Flora (Supplement). Text and Atlas (2 Vol.)*; British-Museum (Natural History): London, UK, 1961; p. 354, xi +.
64. Collinson, M.E. *Fossil Plants of the London Clay. Field Guides to Fossils Series, 1*; The Palaeontological Association: London, UK, 1983; p. 121.
65. Aldiss, D.T. *The Stratigraphical Framework for the Palaeogene Successions of the London Basin, UK*; British Geological Survey: Nottingham, UK, 2014; p. 87.
66. Jarzembowski, E.A. Fossil insects from the London Clay (Early Eocene) of southern England. *Tert. Res.* **1992**, *13*, 87–94.
67. Rundle, A.J.; Cooper, J. Occurrence of a fossil insect larva from the London Clay of Herne Bay, Kent. *Proc. Geol. Ass.* **1971**, *82*, 293–295. [[CrossRef](#)]
68. Collinson, M.E.; Cleal, C.J. Early and early-middle Eocene (Ypresian-Lutetian) palaeobotany of Great Britain. Mesozoic and tertiary palaeobotany of Great Britain. Ed. C.J. Cleal, B.A. Thomas, D.J. Batten, M.E. Collinson. *Geol. Conserv. Rev. Ser. Peterb.* **2001**, *22*, 185–226.
69. Stull, G.V.; Adams, N.F.; Manchester, S.R.; Sykes, D.; Collinson, M.E. Revision of Icacinaceae from the Early Eocene London Clay flora based on X-ray micro-CT1. *Botany* **2016**, *94*, 713–745. [[CrossRef](#)]
70. Reid, E.M.; Chandler, M.E.J. *The London Clay Flora*; British Museum (Natural History): London, UK, 1933; p. 561.
71. Zachos, J.; Pagani, M.; Sloan, L.; Thomas, E.; Billups, K. Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science* **2001**, *292*, 686–693. [[CrossRef](#)]
72. Zachos, J.C.; Dickens, G.R.; Zeebe, R.E. An early Cenozoic perspective on greenhouse warming and carbon-cycle dynamics an early Cenozoic perspective on Greenhouse warming and carbon cycle dynamics. *Nature* **2008**, *451*, 17. [[CrossRef](#)] [[PubMed](#)]
73. Collinson, M.E. Taphonomy and fruiting biology of Recent and fossil *Nypa*. *Spec. Pap. Palaeontol.* **1993**, *49*, 165–180.
74. Collinson, M.E. Plant macrofossils from the Bracklesham Group (early and middle Eocene), Bracklesham Bay, West Sussex, England: Review and significance in the context of coeval British Tertiary floras. *Tert. Res.* **1996**, *16*, 175–202.
75. Collinson, M.E. Fruit and seed floras from the Palaeocene/Eocene transition and subsequent Eocene in southern England: Comparison and palaeoenvironmental implications. *Trans. Geol. Soc.* **2000**, *122*, 36–37. [[CrossRef](#)]
76. Collinson, M.E. Cenozoic evolution of modern plant communities and vegetation. In *Biotic Response to Global Change: The Last 145 Million Years*; Culver, S., Rawson, P., Eds.; Cambridge University Press: Cambridge, UK, 2000; pp. 223–243.
77. Dlussky, G.M.; Wappler, T.; Wedmann, S. New middle Eocene formicid species from Germany and the evolution of weaver ants. *Acta Palaeontol. Pol.* **2008**, *53*, 615–626. [[CrossRef](#)]
78. Radchenko, A.G.; Perkovsky, E.E. Wheeler’s dilemma revisited: First *Oecophylla*-*Lasius* syninclusion and other ant syninclusions in the Bitterfeld amber (late Eocene). *Invertebrate Zool.* **2021**, *18*, 47–65. [[CrossRef](#)]

79. Perfilieva, K.S. Distribution and differentiation of fossil Oecophylla (Hymenoptera: Formicidae) species by wings imprints. *Paleontol. J.* **2021**, *55*, 76–89. [\[CrossRef\]](#)
80. Mänd, K.; Muehlenbachs, K.; McKellar, R.C.; Wolfe, A.P.; Konhauser, K. Distinct origins for Rovno and Balticambers: Evidence from carbon and hydrogen stable isotopes. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **2018**, *505*, 265–273. [\[CrossRef\]](#)
81. Perkovsky, E.E.; Olmi, M.; Vasilenko, D.V.; Capradossi, L.; Guglielmino, A. First Bocchus Ashmead (Hymenoptera: Dryinidae) from Upper Eocene Rovno amber: *B. schmalhauseni* sp. nov. *Zootaxa* **2020**, *4819*, 544–556. [\[CrossRef\]](#)
82. Perkovsky, E.E.; Nel, A. A new Rovno amber termite genus (Isoptera, Rhinotermitidae) from Styr River basin. *Palaeont. Electron.* **2021**, *24*, a05. [\[CrossRef\]](#)
83. Anisyutkin, L.N.; Perkovsky, E.E. Two new cockroaches (Dictyoptera: Ectobiidae) from Rovno amber. *Biologia* **2023**. in print.
84. Legalov, A.A.; Nazarenko, V.Y.; Perkovsky, E.E. A new genus of fungus weevils (Coleoptera: Anthribidae) in Rovno amber. *Foss. Rec.* **2018**, *21*, 207–212. [\[CrossRef\]](#)
85. Matalin, A.V.; Perkovsky, E.E.; Vasilenko, D.V. First record of tiger beetles (Coleoptera, Cicindelidae) from Rovno amber with the description of a new genus and species. *Zootaxa* **2021**, *5016*, 243–256. [\[CrossRef\]](#) [\[PubMed\]](#)
86. Telnov, D.; Perkovsky, E.E.; Vasilenko, D.V.; Yamamoto, S. The first fossil Coleoptera record from the Volyn Region, Ukraine, with description of a new Glesoconomorpha (Coleoptera, Mycteridae) in syninclusion with Winterschmidtidae (Acari) and a key to species. *ZooKeys* **2021**, *1068*, 189–201. [\[CrossRef\]](#)
87. Telnov, D.; Perkovsky, E.E.; Kundrata, R.; Kairišs, K.; Vasilenko, D.V.; Bukejs, A. Revealing Palaeogene distribution of the Ptilodactylidae (Insecta: Coleoptera): The first Ptilodactyla Illiger, 1807 records from Rovno amber of Ukraine. *Hist. Biol.* **2022**, *1*–10. [\[CrossRef\]](#)
88. Yamamoto, S.; Nazarenko, V.Y.; Vasilenko, D.V.; Perkovsky, E.E. First fossil species of ship-timber beetles (Coleoptera: Lymexyidae) from Eocene Rovno amber (Ukraine). *Foss. Rec.* **2022**, *25*, 65–74. [\[CrossRef\]](#)
89. Nabozhenko, M.V.; Perkovsky, E.E. *Oracula campbelli* sp. n.—The first fossil darkling beetle (Coleoptera: Tenebrionidae: Alleculinae) from the late Eocene Rovno amber (Ukraine). *Zootaxa* **2023**, *5230*, 245–250. [\[CrossRef\]](#)

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.