

The Terebrantia (Insecta: Thysanoptera) of the Maltese Islands

Godwin Degabriele ^{1,*} , Adriano Cavalleri ², Arturo Goldarazena ³  and David Mifsud ¹

¹ Institute of Earth Systems, Rural Sciences and Food Systems, University of Malta, MSD 2080 Msida, Malta

² Biological Sciences Institute, Carreiros Campus, Federal University of Rio Grande, Rio Grande 96203-900, Brazil

³ National Museum of Natural Sciences-CSIC, Department of Biodiversity and Evolutionary Biology National Reference Laboratory for Arthropods, C/Serrano Duplicado, CP 28006 Madrid, Spain

* Correspondence: godwin.degabriele@um.edu.mt; Tel.: +356-2590-7411

Abstract: Thirty-nine species of terebrantian Thysanoptera belonging to four families and 22 genera are here recorded from the Maltese Islands. Of these, 33 represent new records to this archipelago. Thrips were collected from 65 different locations over a seven-year period, covering the main habitat types found across the Maltese Islands, namely steppe, garigue, maquis and woodland, but also sand dunes and saltmarshes as well as roadsides, private and public gardens, greenhouses and cultivated fields. An illustrated dichotomous key to identify the Terebrantia of the Maltese Islands is presented. Chorological data for the species researched in the current study shows that the majority of these insects are of a European Mediterranean origin, though the geographical distribution of some of them extends to Africa and the Middle East. Seven species associated with agricultural commodities were found to be of alien origin; however they were locally found in small numbers and do not pose a threat to horticulture.

Keywords: Malta; thrips; Mediterranean; identification keys



Citation: Degabriele, G.; Cavalleri, A.; Goldarazena, A.; Mifsud, D. The Terebrantia (Insecta: Thysanoptera) of the Maltese Islands. *Diversity* **2023**, *15*, 514. <https://doi.org/10.3390/d15040514>

Academic Editors: Dimitrios Kontogiannatos, Anna Kourti and Piero G. Giulianini

Received: 29 January 2023

Revised: 8 March 2023

Accepted: 14 March 2023

Published: 3 April 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Thrips belong to the insect order Thysanoptera which comprises more than 6400 described species worldwide. These insects are found on plants, leaf litter or dead twigs where they feed on pollen, plant sap and fungi. The order is divided into two sub-orders: Terebrantia and Tubulifera, which are physiologically and behaviorally different. Around 100 thrips species are reported to impact crop productivity around the world, almost all belonging to Terebrantia [1].

Generally, thrips are associated with plants, but these plants are only considered as host-plants if the insect completes its life cycle on the mentioned plant [2]. Thus, the establishment of whether a plant is a true host of a particular thrips relies on the presence of immature stages. Although a small number of thrips species do not stray from their host-plant, the majority, as adults, tend to use other plants as a source of food or shelter, but these should never be regarded as host plants.

The Maltese archipelago consists of the habitable islands of Malta and Gozo and a number of much smaller islands including Comino, Selmunett and Filfla. The archipelago is situated in the center of the Mediterranean basin and covers an area of 316 km² [3]. The climate of the archipelago alternates between a mild rainy cold season and a dry hot summer [4] lasting from early June to the end of September, with temperatures averaging about 30 °C, though sometimes peaking at 40 °C. Water availability is rather scarce, with typical average rainfall for the period 1900–2000 being 550 mm [5]. Abstraction of groundwater, necessary to sustain the relatively limited perennial surface water ecosystems, exceeds aquifer recharge rate [5].

The Maltese Islands lack many habitats such as mountains, rivers, large forests, alpine environments and others and thus insects associated with these habitat types are lacking.

Nonetheless a range of natural terrestrial habitats still occur, which can be categorized into four main types: the steppe, the garigue, the maquis and the woodland [6]. A number of valleys with temporary fresh water courses as well as a few degraded sand dunes and saltmarshes can also be found. All of these habitats can potentially host thrips on the vegetation therein. Despite the limited variety of habitats, the biodiversity of the flora of the Maltese Islands is rather rich, with some 1100 indigenous plants [7] and a relatively large number of cultivated plants, some of which have become established escapees.

The literature on the Thysanoptera of the Maltese Islands is very fragmentary. Seven species of Terebrantia have been recorded. These include *Thrips tabaci* a widely known agricultural pest that affects onions, garlic and cauliflowers [8–10], *Tenothrips discolor* [11], *Frankliniella occidentalis*, *Heliethrips haemorrhoidalis* [10,12], *Aeolothrips tenuicornis*, *Melanthrips fuscus* [13] and *Melanthrips lybicus* [14,15].

The Maltese Islands are becoming urbanized at a rapid rate due to the high population density, scoring at 1649 per km² [16]. This problem is compounded by the fact that the economy of the Maltese Islands heavily depends on tourism, with c. 1.5 million tourists visiting the islands every year. These factors leave a considerable negative impact on the Maltese natural environment. Extensive use of plant protection products is also causing the decline of a number of indigenous insect species, many of which serve as ecosystem providers. A knowledge of insect biodiversity is a prerequisite to enable proper conservation programs, thus avoiding loss of species diversity due to climate change, among other issues.

This present study was thus undertaken to document the biodiversity of the terebrantians that inhabit the Maltese Islands. It also provides an illustrated dichotomous key.

2. Materials and Methods

During the current study, thrips specimens were collected from 65 different locations from Malta and Gozo (as shown in Figure 1) between Spring 2015 and Spring 2022. The habitats and flora therein are described in Table 1. Thrips were collected from the main habitats found in the Maltese Islands as mentioned earlier, but also from private and public gardens, town and country roadsides, public green areas, private farms and cultivated open fields.




Figure 1. Map of the Maltese Islands (Gozo and Malta) showing the locations where data were collected between 2015–2022 (with names, habitat description and predominant vegetation of locations indicated in Table 1).

Table 1. Locations where data were collected for this study: habitat type and brief description.

Site No.	Location	Habitat Type and Description	
1	Ta' Sabbara, Żabbar	Steppe, with typical vegetation including <i>Foeniculum vulgare</i> , <i>Vicia sativa</i> and, especially when in a degraded state, <i>Asphodelus ramosus</i> , <i>Galactites tomentosa</i> and <i>Stipa capensis</i> , and. Most plants in this habitat dry up during the dry summer period [6].	
2	Wied Anġlu, Birguma		
3	Dingli Cliffs, Dingli	Garigue, characterised by low lying, often aromatic shrubs adapted to resist drought and strong winds, e.g., <i>Thymbra capitata</i> . Commonly found species include <i>Drimia</i> (= <i>Urginea</i>) <i>maritima</i> , <i>Erica multiflora</i> , and <i>Euphorbia melitensis</i> [6].	
4	Country road, l/o Rabat		
5	Manikata		
6	Ċirkewwa		
7	Lapsi, Siġġiewi		
8	Xrobb l-Għajin		
9	Għaxqet l-Għajn, Għargħur		
10	Mġiebaħ, Xemxija		
11	Lippija Tower, Mġarr		
12	Pembroke		
13	l/o Majjistral Park, Għajn Tuffieħa		
14	Xlendi Tower, Xlendi, Gozo		
15	Ta Ċenċ, Gozo		
16	Il-Maqluba, Qrendi		Maquis, with plants consisting largely of evergreen small trees such as <i>Ceratonia siliqua</i> and <i>Olea europaea</i> , large shrubs such as <i>Pistacea lentiscus</i> and climbers such as <i>Hedera helix</i> and <i>Smilax aspera</i> , as well as herbaceous shade loving species such as <i>Acanthus mollis</i> [6].
17	Il-Fawwara, Siġġiewi		
18	Mġarr ix-Xini, Gozo		
19	Has-Saptan, Gudja	Woodland, comprising of Mediterranean sclerophyll forest with trees including <i>Quercus ilex</i> and <i>Pinus halepensis</i> and an undergrowth of smaller shrubs. <i>Tamarix</i> spp. prevails on coastal wood remnants, while <i>Populus alba</i> occurs in riparian woodland remnants [6].	
20	Buskett, Dingli		
21	Wied Ghollieqa, San Ġwann	Valleys, with slopes having plants typically found in maquis environments and the valley itself. Plants occurring in these habitats include <i>Hyparrhenia hirta</i> , <i>Rubus ulmifolius</i> and <i>Potentilla reptans</i> and, where water courses are present also some hydrophytes such as <i>Ranunculus saniculifolius</i> .	
22	Wied Hesri, Siġġiewi		
23	Wied Musa, Siġġiewi		
24	Wied Xkora, Siġġiewi		
25	Wied Qirda, Żebbuġ		
26	Wied il-Baqquieħa, Żebbuġ		
27	Wied Speranza, Mosta		
28	Wied Qlejgħa, Rabat		
29	Wied tal-Fiddien, Rabat		
30	Wied Babu, Żurrieq		
31	Il-Qattara, Dwejra, Gozo		
32	Ramla l-Ħamra, Gozo.	Sand dune, with vegetation that needs to be suited to harsh conditions including high temperatures, dryness, occasional inundation by seawater and accumulation of sand. These plants include <i>Cakile maritima</i> , <i>Elytrigia juncea</i> , <i>Eryngium maritimum</i> , <i>Lotus cytisoides</i> and <i>Pancratium maritimum</i> .	
33	Il-Ballut, Marsaxlokk	Saltmarshes, with plants that tolerate high salinity such as <i>Athrocneum macrostachyum</i> , <i>Salsola soda</i> and <i>Salicornia ramosissima</i> .	
34	Siġġiewi (roads)	Town roads. Plants growing along these roads include a variety of both indigenous species (e.g., <i>Diplotaxis tenuifolia</i>) as well as non-indigenous species. These sites yield plants which are cultivated, non-indigenous species.	
35	Naxxar (road)		
36	Rabat (roads)		
37	Żabbar (road)		
38	Attard (road)		
39	Victoria, Gozo (roads)		

Table 1. Cont.

Site No.	Location	Habitat Type and Description
40	Popeye Village, Mellieħa road	Country roads. These host a number of indigenous plants including <i>Glebionis coronaria</i> and <i>Antirrhinum siculum</i> .
41	Ġhajn Tuffieħa road	
42	Ġebel San Pietru, Madliena	
43	Il-Kuncizzjoni, l/o Rabat	
44	Żebbuġ (roads)	
45	Luqa road	
46	Mosta road	
47	Mellieħa by-pass	By-passes. A number of central strips and roundabouts on the proximity of these roads include cultivated species such as <i>Solandra maxima</i> , <i>Albizia julibrissin</i> and <i>Nerium oleander</i> .
49	Zabbar (private gardens)	Private gardens. These include a variety of cultivated herbaceous plants and fruit trees such as <i>Citrus</i> .
50	Mosta garden	
51	Attard garden	
52	Junior College Ringroad, Msida	
53	University of Malta Grounds, Msida	
54	Siggiewi plant nursery	Public gardens. These feature a combination of cultivated plants, e.g., <i>Yucca gloriosa</i> and <i>Salvia coccinea</i> and indigenous species including <i>Tamarix africana</i> and <i>Atriplex halimus</i> .
55	St. Thomas Bay, Marsaskala garden	
56	Sta. Lucia garden	
57	Golden Bay garden	
58	Qbajjar, Gozo garden	
59	Floriana Car Park	Public cultivated green areas with plants as described in the “Town roads” section.
60	Qormi (roundabout)	
61	San Ġwann farm	Private farms, which grow a variety of crop plants.
62	Qormi farm	
63	Siggiewi farm	
64	Ta’ Qali l/o Attard field	Private open cultivated fields with a variety of plants as described in private farms.
65	Tal-Ferħa, Ġhargħur field	

Thrips were collected from 398 different species of indigenous and cultivated plants. The literature on some 700 thrips species recorded mostly from the Palaearctic region was consulted, e.g., zur Strassen (2003) [14] in order to find out which plant species these thrips were collected from. If these plant species were present in the Maltese Islands, they were chosen for sampling purposes. Moreover, other plants which prevailed in the different habitats of the Maltese archipelago were also sampled. Plants were beaten using a plastic rod onto a white plastic tray [17]. All specimens collected in the tray were transferred into collecting Eppendorf tubes using a fine brush (e.g., size 00) dipped in AGA fluid mixture consisting of 10 parts of 60% ethyl alcohol, one part of glycerin and one part of glacial acetic acid. Additional specimens of thrips were collected from Malaise traps present in two locations in Malta (Fawwara and Buskett).

Four hundred and sixty of the collected specimens were mounted on microscope slides for examination under compound microscopy. Prior to mounting, thrips specimens were macerated by immersing in 5% sodium hydroxide. Dark specimens were immersed in this agent for about 12 h while lighter specimens were left for less time depending on the colour intensity of the specimen. The specimens were subsequently dehydrated by placing the specimens in increasing concentrations of ethanol (75, 80, 95, and 100%) [18]. The specimens were finally cleared by immersing in clove oil prior to mounting in Canada balsam [18], after which they were mounted by placing them on the cover slip and then placing the glass slide on top [18].

Mounted specimens were viewed using a Leica DVM6 microscope and identified with the use of a number of printed [14,17,19–24] and on-line sources [25–28] as well as published works having identification keys and species descriptions. Different morphological features observed in the species identified were used to construct the dichotomous key.

The majority of the specimens examined for the current study form part of the private collection of Godwin Degabriele (GD), though some 15 specimens come from the private collection of David Mifsud (DM), which were collected during the late 1990s to the mid 2000s. A number of species from the collection of GD were donated to Arturo Goldarazena (AG) who will deposit them in the collection of the Museo Nacional de Ciencias Naturales, Madrid.

Abbreviations used in the following sections include the following: Godwin Degabriele (GD); Charles Farrugia (CF); Sylvan Farrugia (SF); David Mifsud (DM); E.R. Speyer (ERS); I.A. Speyer (IAS); Gillian Watson (GW); J.W. Ismay (JWI); Niki Young (NY); slide mounted specimens (sm); specimens conserved in AGA (aga); specimens housed in the British Museum of Natural History (BMNH); thrips which represent new record for the Maltese Islands (+). Representative species from the private collection of GD will eventually be donated to the BMNH.

3. Results

Thirty-nine species of Terebrantia belonging to four families and 22 genera were identified, with 33 of these being new records to the Maltese Islands. These species are discriminated by the illustrated dichotomous key provided below.

3.1. Key to the Terebrantia of the Maltese Islands

- 1 Abdominal segment X conical in females (Figure 2a) and rounded posteriorly in males; fore wings with one or two longitudinal veins (Figure 2c); females with external ovipositor (Figure 3e,f) **Terebrantia** .. 2
- Abdominal segment X tubular in both sexes (Figure 2b); fore wings with no veins (Figure 2d); females lacking external ovipositor (Figure 2b) **Tubulifera**




Figure 2. Abdominal segment X (a,b): (a) conical; (b) tubular; fore wing (c,d): (c) with longitudinal veins; (d) with no veins.

- 2 Antennal segments III and IV with sensoria consisting of a ridge-like structure set in parallel (Figure 3a) or perpendicular (or oblique) to each segment (Figure 3b), never produced as trichomes; fore wing broad, being five to ten times as long as broad (Figure 3c); females with ovipositor curving upwards towards abdominal segments (Figure 3e) 3
- 3 Antennal segments III and IV with emergent sensoria, consisting of an acorn-shaped structure (Figure 3g), or a simple (Figure 3h) or forked trichome (Figure 3i); fore wing narrow, being 11–18 times as long as broad and often tapering apically (Figure 3d); females with ovipositor curving downwards (Figure 3f) 15




Figure 3. Sensoria on antennal segments III and IV (a,b): (a) parallel to length of segment; (b) perpendicular to length of segment; fore wing tip (c,d): (c) broad; (d) narrows apically; ovipositor (e,f): (e) curving upwards towards body; (f) curving downwards; antennal sensoria on segments III and IV (g,i): (g) cone-shaped; (h) simple or hair-like; (i) forked.

- 3 Antennal segments III and IV with sensoria set in parallel to length of segment (Figure 3a); antennal segments VI–IX broadly joined basally (Figure 4a) **Aeolothripidae** .. 4
 Antennal segments III and IV with sensoria set perpendicular or oblique to length of segment (Figure 3b); antennal segments VI–IX constricted basally and clearly distinct from each other (Figure 4b) **Melanthripidae** .. 12




Figure 4. Antennal segments VI–IX: (a) broadly joined basally; (b) distinct from each other.

- 4 Antennal segment III about 15 times as long as wide (Figure 5a); antennal segments III and IV with sensoria consisting of a multitude of dot-like structures lined up parallel to length of segment (Figure 5b); fore wings with dark band sub-apically (Figure 5c); abdominal segments I–III significantly narrower than other segments, giving the appearance of a wasp-like waist (Figure 5e) **Franklinothrips megalops** Trybom
 Antennal segments III 3.3–5 times as long as wide; antennal segments III and IV with sensoria consisting of one or a small number of continuous ridge shapes lined up parallel to length of segment (Figure 5b); fore wings with light coloured sub-apical region (Figure 5d); abdominal segments I–III only slightly narrower than other segments (Figure 5f) 5




Figure 5. (a) antennal segments III–IX; (b) sensorium on antennal segment III; fore wing tip (c,d): (c) with dark band sub-apically; (d) with light-coloured sub-apical region; upper abdominal segments (e,f): (e) being considerably narrower than other segments; (f) only slightly narrower than other segments.

- 5 Antennal segments III and IV with sensoria consisting of one continuous ridge-like structure lined up parallel to length of segment (Figure 6a); fore wings broad, being less than six times as long as broad and with two dark bands (Figure 6c); pronotum postero-marginal setae as long as discal setae *Aeolothrips* Haliday .. 6
- 6 Antennal segments III and IV with sensoria consisting of lens-shaped structures lined up parallel to length of segment (Figure 6b); fore wings narrow, being more than six times as long as broad and of a uniformly pale brown hue (Figure 6d); pronotum bearing a pair of postero-marginal setae that are longer than discal setae..... *Rhipidothrips* Uzel .. 9




Figure 6. Antennal sensoria on segment III (a,b): (a) continuous and ridge-shaped; (b) lens-shaped; fore wing (c,d): (c) broad and banded; (d) narrow and monochrome.

- 6 Fore wing margin at tip as light as wing membrane (Figure 7a)..... 7
- Fore wing margin at tip darker than wing membrane (Figure 7b) 8




Figure 7. Fore wing: (a) pale coloured margin at tip; (b) dark coloured margin at tip.

- 7 Body bicoloured, with head and thorax yellow with brown areas medially (Figure 8a); antennal segment III bicoloured, with the basal half of segment pale yellow, while the apical half dark brown (Figure 8c) *Aeolothrips gloriosus* Bagnall
 Body uniformly dark brown, though upper abdominal segments can sometimes be lighter coloured (Figure 8b); antennal segment III yellow or yellowish brown with the distal fifth brown, paler than segment II (Figure 8d)
 *Aeolothrips intermedius* Bagnall




Figure 8. Body (a,b): (a) bicoloured; (b) uniformly brown; antennal segment III (c,d): (c) distal half brown; (d) only distal fifth brown.

- 8 Antennal segment III with sensorium being one third the length of the segment (Figure 9a); antennal segment IV with sensorium two thirds the length of the segment (Figure 9a); pronotum pale, often yellow (Figure 9c) *Aeolothrips melisi* Priesner
 Antennal segment III and IV with sensoria about half the length of the segment (Figure 9b); pronotum dark, usually brown (Figure 9d)
 *Aeolothrips tenuicornis* Bagnall




Figure 9. Antennal segments III and IV (a,b): (a) sensoria about one third the length of segment in III, two thirds the length of segment in IV; (b) sensoria about half the length of segment in both III and IV; pronotum (c,d): (c) pale; (d) dark.

- 9 Antennal segment II yellow (Figure 10c); pronotum yellow to light brown and with three to four long pairs of postero-marginal setae (Figure 10a); females always macropterous (Figure 10f); abdominal tergites with little or no sculpture or microtrichia (Figure 10g) *Rhipidothrips gratiosus* Uzel.
 Antennal segment II brown (Figure 10d); pronotum brown and with two to three long pairs of postero-marginal setae (Figure 10b); females usually micropterous (Figure 10e); abdominal tergites sculptured, sometimes lined with microtrichia (Figure 10h) 10




Figure 10. Antennal segment II (a,b): (a) yellow; (b) brown; pronotum (c,d): (c) with four pairs of long postero-marginal setae; (d) with three pairs of postero-marginal setae; fore wings (e,f): (e) micropterous; (f) macropterous; abdominal tergites (g,h): (g) with no sculpture; (h) with sculpture.

- 10 Internal body colour orange brown (Figure 11a); females macropterous. On *Stipa capensis* *Rhipidothrips unicolor* zur Strassen.
 Internal body colour often carmine red (Figure 11b); females can be macropterous or micropterous (Figure 10g). On different species of Poaceae 11




Figure 11. Abdominal segments II and III showing internal body color: (a) brown; (b) red.

- 11 Abdominal tergites with microtrichia on sculpture lines (Figure 12a); females usually macropterous..... *Rhipidothrips niveipennis* O.M. Reuter
 Abdominal tergites with no microtrichia on sculpture (Figure 12b); females usually micropterous *Rhipidothrips brunneus* Williams




Figure 12. Section of abdominal tergites: (a) with microtrichia on sculpture; (b) with no microtrichia on sculpture.

- 12 Fore wing banded (Figure 13a) *Melanthrrips ficalbii* Buffa
 Fore wing evenly brown, not banded (Figure 13b) and with a small pale sub-basal area 13



Figure 13. Fore wing: (a) banded; (b) not banded.

- 13 Anterior margin of fore wing between transverse veins with one row of setae (Figure 14a) *Melanthrips knechteli* Priesner
- Anterior margin of fore wing between transverse veins with two rows of setae (Figure 14b) 14




Figure 14. Fore wing: (a) with anterior margin between transverse veins with one row of setae; (b) with anterior margin between transverse veins with two rows of setae.

- 14 Hind tibia with one long seta (Figure 15a) *Melanthrips fuscus* (Sulzer)
- Hind tibia with two long setae (Figure 15b) *Melanthrips lybicus* Priesner




Figure 15. Hind tibia: (a) with one long seta; (b) with two long setae.

- 15 Antennae nine-segmented (Figure 16a); antennal segments III and IV with cone-shaped sensoria (Figure 3g) **Stenurothripidae**
- *Holarthrothrips tenuicornis* Priesner
- Antennae six-, seven-, or eight-segmented (Figure 16b,d); antennal segments III and IV with sensoria consisting of a single (simple or hair-like) (Figure 3h) or forked filamentous structure (Figure 31) **Thripidae** .. 16



Figure 16. Antenna (a–d): (a) nine-segmented; (b) eight-segmented; (c) seven-segmented; (d) six-segmented.

- 16 Metathoracic endofurca lyre-shaped, with arms extending well beyond length of segment (Figure 17a); median tergal setae on abdomen arising close to each other (Figure 17c) **Dendrothripinae**
 *Dendrothrips saltator* Uzel
- Metathoracic endofurca not lyre-shaped (Figure 17b), and usually not extending beyond length of segment; median tergal setae set far apart (Figure 17d) 17




Figure 17. Pterothorax (a,b): (a) with lyre-shaped metathoracic endofurca; (b) with metathoracic endofurca not lyre-shaped; tergites (c,d): (c) median setae close to each other; (d) median setae far away from each other.

- 17 Fore wing with one row of postero-marginal cilia (Figure 18a); antennal segments VI–VIII long and slender when compared with basal segments. (Figure 18c).....
 **Panchaetothripinae** .. 18
- Fore wing with two rows of postero-marginal cilia (Figure 18b); antennal segments VI–VIII not sharply elongated (Figure 18d) **Thripinae** .. 19




Figure 18. Fore wing (a,b): (a) one row of postero-marginal cilia; (b) two rows of postero-marginal cilia; terminal antennal segments (c,d): (c) ending in a sharp point; (d) not ending in a sharp point.

- 18 Fore tarsi one-segmented (Figure 19a); fore wings uniformly light brown; postero-marginal cilia straight (Figure 19c); fore wing apex rounded (Figure 19c) *Heliothrips haemorrhoidalis* (Bouchè)
 Fore tarsi two-segmented (Figure 19b); fore wings brown with white stripes; postero-marginal cilia wavy (Figure 19d); fore wing apex pointed (Figure 19d) *Hercinothrips femoralis* Reuter




Figure 19. Fore tarsus (a,b): (a) one-segmented; (b) two-segmented; fore wing (c,d): (c) postero-marginal cilia straight and apex rounded; (d) postero-marginal cilia wavy and apex pointed.

- 19 Pronotum considerably narrower at anterior margin than at posterior margin (Figure 20a)..... *Chirothrips* Haliday .. 20
 Pronotum having the same width at both margins (Figure 20b) :..... 22




Figure 20. Pronotum: (a) considerably narrower at anterior margin than at posterior margin; (b) being almost the same width at both margins.

- 20 Antennal segment II symmetrical *Chirothrips hamatus* Trybom
 Antennal segment II asymmetrical, projecting laterally (Figure 21a,b) 21




Figure 21. Antennal segment II showing outer edge (a,b): (a) nearly straight; (b) at an obtuse angle from the base of the segment; antennal segment IV (c,d): (c) with a forked sense cone; (d) with a simple sense cone.

- 21 Antennal segment II with the outer edge nearly straight (Figure 21a); antennal segment IV with forked sense cone (Figure 21c); antennal segment VII longer than VIII; males usually macropterous *Chirothrips meridionalis* Bagnall
- Antennal segment II with the outer edge at an obtuse angle from the base of segment, creating a tip with bulge (Figure 21b); antennal segment IV with simple sense cone (Figure 21d); antennal segment VII shorter than VIII; males micropterous *Chirothrips manicatus* Haliday
- 22 Abdominal segment X bearing a pair of thorn-shaped setae (Figure 22a) *Limothrips* Haliday .. 23
- Abdominal segment X without thorn-shaped setae (Figure 22b) 24




Figure 22. Abdominal segment X: (a) with thorn-shaped setae; (b) lacking thorn-shaped setae.

- 23 Antennal segment II asymmetrical, with elongated extremity at outer side of segment (Figure 23a); antennal segments III and IV with forked sense cones (Figure 23c) *Limothrips angulicornis* Jablonowski
- Antennal segment II rounded and symmetrical (Figure 23b); segments III and IV with simple sense cones (Figure 23d) *Limothrips cerealium* Haliday




Figure 23. Antennal segment II (a,b): (a) asymmetrical; (b) symmetrical; antennal segments III and IV (c,d): (c) with forked sense cones; (d) with simple sense cones.

- 24 Antennal segments III and IV with simple sense cones (Figure 3h) 25
- Antennal segments III and IV with forked sense cones (Figure 3i) 28
- 25 Antennae six-segmented (Figure 16d); head and body having a distinct bright yellow colour (Figure 24a) *Aptinothrips rufus* Haliday
- Antennae eight-segmented (Figure 16b); head and body pale yellow, brown, or bi-coloured (Figure 24b) 26




Figure 24. Male habitus: (a) yellow body colour; (b) body bicoloured.

- 26 Fore wing with setae on vein always long, dark and capitate (blunt tipped) (Figure 25b); second vein with no setae (Figure 25a); antennal segment VIII ending in a sharp point (Figure 18c); males macropterous *Echinothrips americanus* Morgan
- Fore wing setae on vein not always long and dark and tapering to a point; second vein with setae that have pointed tips (Figure 25c); antennal segment VIII rounded at tip (Figure 18d); males micropterous 27




Figure 25. (a) Fore wing with second vein with no setae; (b) setae on vein of fore wing with capitate tips; (c) fore wing with second vein with setae.

- 27 Females micropterous; head with reticulate (network-like) sculpture; ridge present between antennal sockets (Figure 26a) *Prosopothrips nigriceps* Bagnall
- Females macropterous; head with no distinct sculpture; ridge lacking between antennal sockets (Figure 26b) *Bregmatothrips dimorphus* (Priesner)



Figure 26. Head: (a) with reticulate sculpture and with a ridge between antennal sockets; (b) with no distinct sculpture and with no ridge between antennal sockets.

- 28 Pronotum with no long postero-angular setae (Figure 27a); fore wing with dark band at proximal region (Figure 27c) *Anaphothrips sudanensis* Trybom
- Pronotum with one to two pairs of long postero-angular setae (Figure 27b); fore wing with no dark band at proximal region (Figure 27d) 29




Figure 27. Pronotum (a,b): (a) with no long postero-angular setae; (b) with long postero-angular setae present; fore wing (c,d): (c) with a dark band at proximal region; (d) with no dark band at proximal region.

- 29 Both sexes micropterous; antennal segment V yellow with very little brown on distal margin (Figure 28a) *Asphodelothrips croceicollis* (Karny)
- Both sexes macropterous; antennal segment V either evenly brown (Figure 28b) or distinctly bicoloured (Figure 28c) 30




Figure 28. Antennal segment V: (a) pale yellow; (b) evenly brown; (c) bicoloured.

- 30 Pronotum with five (rarely 4) pairs of prominent postero-marginal setae, with submedian pair more than twice as long as the discal setae (Figure 29a); male antennal segment VI as long as segments IV and V together (Figure 29c); male with a row of around 12 pore plates on each of abdominal sternites II–VII (Figure 29e) *Pezothrips kellyanus* (Bagnall)
 Pronotum with variable number of prominent postero-marginal setae, usually fewer than five and submedian pair never twice as long as discal setae (Figure 29b); male antennal segment VI shorter than segments IV and V together (Figure 29d); male with only one or no pore plate on each of abdominal sternites II–VII (Figure 29f) 31




Figure 29. Pronotum (a,b): (a) with five pairs of prominent postero-marginal setae; (b) with four pairs of prominent postero-marginal setae; antennal segments V–VIII (c,d): (c) antennal segment VI as long as IV + V; (d) antennal segment VI shorter than IV + V; male abdominal sternite (e,f): (e) with around 12 small pore plates per sternite; (f) with one large pore plate per sternite.

- 31 Antennal segment VI with a sensorium of which the length of the base (insertion) of the sense cone is longer than the width of segment VII (Figure 30a).....
 *Odontothrips meliloti* Priesner
 Antennal segment VI with sensorium of which base is shorter than the width of segment VII (Figure 30b) 32




Figure 30. Sensorium on antennal segment VI: (a) with base of sensorium longer than width of segment VII; (b) with base of sensorium shorter than width of segment VII.

- 32 Abdominal tergite VIII without paired lateral ctenidia (Figure 31a) often replaced by rows of microthrichia 33
- Abdominal tergite VIII with paired lateral ctenidia (Figure 31b) sometimes with only few and sparse (not arranged in a single row) 35




Figure 31. Left side of abdominal segment VII: (a) lacking ctenidia; (b) with ctenidia.

- 33 Fore wing clavus with five marginal setae (Figure 32a); second vein of fore wing with 8–10 setae *Tenothrips discolor* (Karny)
 Fore wing clavus with six marginal setae (Figure 32b); second vein of fore wing with 11 or more setae 34




Figure 32. Clavus (a,b): (a) with six marginal setae; (b) with five marginal setae.

- 34 Abdominal sternites bearing discal setae; ocellar setae iii arising just antero-lateral to ocellar triangle (Figure 33a); pronotum with one pair of postero-angular setae longer than discal setae (Figure 33c)..... *Oxythrips ajugae* Uzel
 Abdominal sternites with no discal setae; ocellar setae iii either arising outside ocellar triangle (Figure 33b) or between hind ocelli; pronotum with two pairs of postero-angular setae longer than discal setae (Figure 33d) *Ceratothrips ericae* (Haliday)




Figure 33. Head (a,b): (a) showing ocellar setae iii just anterolateral to ocellar triangle; (b) showing ocellar setae iii arising outside ocellar triangle; pronotum (c,d): (c) with one pair of postero-angular setae longer than discal setae; (d) with two pairs of postero-angular setae longer than discal setae.

- 35 Ctenidia on abdominal segment VIII situated antero-laterally to spiracle (Figure 34a); anterior margin of pronotum bearing at least one pair of long setae *Frankliniella* Karny .. 36
 Ctenidia on abdominal segment VIII situated postero-mesad to spiracles (Figure 34b). anterior margin of pronotum with no long setae *Thrips* Linnaeus 37




Figure 34. Position of ctenidia on abdominal segment VIII: (a) antero-laterally to spiracle; (b) postero-mesad to spiracles.

- 36 Ocellar setae iii arising on anterior margins of ocellar triangle (Figure 35a); meta-thoracic campaniform sensilla present (Figure 35e); microtrichial comb at the posterior margin of tergite VIII fully developed (Figure 35d) *Frankliniella occidentalis* Pergande
 Ocellar setae iii arising close together between hind ocelli (Figure 35b); meta-thoracic campaniform sensilla absent (Figure 35d); microtrichial comb at the posterior margin of tergite VIII not developed or with few short teeth on lateral margins (Figure 35f) *Frankliniella schultzei* Trybom




Figure 35. Head (a,b): (a) with ocellar setae iii arising on anterior margins of ocellar triangle; (b) with ocellar setae iii arising close together between hind ocelli; metanotum (c,d): (c) with campaniform sensilla present; (d) with no campaniform sensilla; abdominal tergite VIII (e,f): (e) with complete microtrichial comb; (f) with no microtrichial comb.

- 37 Antennae eight-segmented (Figure 16b) *Thrips simplex* Morison
 Antennae seven-segmented (Figure 16c) 38
- 38 Abdominal tergite II with four lateral setae (Figure 36a); fore wing clavus with six
 marginal setae (Figure 32a) *Thrips australis* (Bagnall)
 Abdominal tergite II with three lateral setae (Figure 36b); fore wing clavus with five
 marginal setae (Figure 32b) 39




Figure 36. Abdominal tergite II (a,b): (a) with four lateral setae; (b) with three lateral setae.

- 39 Abdominal tergite VIII with microtrichial comb medially incomplete (Figure 37a).....
 *Thrips major* Uzel
 Abdominal tergite VIII with microtrichial comb complete (Figure 37b)
 *Thrips tabaci* Lindeman




Figure 37. Abdominal tergite VIII: (a) medially incomplete; (b) complete.

3.2. Species Catalogue

FAMILY AEOLOTHIRIPIDAE

Aeolothrips gloriosus Bagnall, 1914 †

Material examined: MALTA: Buskett, 19.iv.2016, 2 ♀♀(sm) on *Laurus nobilis*, GD; Buskett, 21.ii.2018, 1 ♀(sm) on *Rhamnus alaternus*, GD; Xemxija, 3.iii.2018, 2 ♀♀(sm) on *Pistacea lentiscus*, GD

Body length: ♀: 1760–1940 µm; ♂: no records.

Wing type: Both sexes are macropterous.

Aeolothrips gloriosus occurs on woody, often deciduous plants such as *Tilia* (Malvaceae), *Fraxinus* (Oleaceae), *Sambucus* (Adoxaceae), *Prunus* (Rosaceae), but also *Citrus* (Rutaceae) and *Olea* (Oleaceae) [14,20,26,27]. In the Maltese Islands, *A. gloriosus* was exclusively found on trees and shrubs including *Laurus nobilis* (Lauraceae), *Pistacea lentiscus* (Anacardiaceae)

and *Rhamnus alaternus* (Rhamnaceae) in woodland and maquis habitats. This species was recorded as a facultative predator [29]. This Aegean species has a distribution which spans from Anatolia to Morocco and the Azores [14]. It has also been recorded from Iran [30] and the Near East [13].

Aeolothrips intermedius Bagnall, 1934 †

Material examined: MALTA: 03.iv.1959, 1 ♀ and instar larva (sm) on *Gladiolus* sp., ERS (BMNH); Wied Qirda, 06.iv.2016, 1 ♂ (sm) on *Glebionis coronaria*, GD; Wied Hesri, 22.iv.2016, 2 ♀♀ (sm) on *Avena* sp., GD (BMNH); Wied Qirda, 17.iii.2017, 1 ♀ (sm) on *Asphodelus ramosus*, GD; Manikata, 29.iv.2017, 1 ♀ (sm) on *Pallenis spinosa*, GD; Fawwara, 22.v.2017, 1 ♀ (sm) from Malaise trap, DM; Lapsi, 09.x.2018, 1 ♀ (sm) on *Hyparrhenia hirta*, GD.

Body length: ♀: 1700–1840 µm; ♂: 1300 µm.

Wing type: Both sexes are macropterous.

A. intermedius has been recorded on a wide variety of unrelated plant species, often on yellow flowered plants such as those of Asteraceae, Cruciferae, Leguminosae, and Poaceae [15,21,24,26,31–34]. In the Maltese Islands, a larval instar was found on *Gladiolus* (Iridaceae) implying that this plant is used as a host by this species. Adults were found on annual herbaceous plants namely Asparagaceae, Asteraceae and Poaceae. This species was recorded as a facultative predator feeding on different insects including other thrips [32]. This species is widespread in Europe and Palearctic Asia and the Middle East [13,14]. It has also been intercepted in the US from plant material originating from Europe [35].

Aeolothrips melisi Priesner, 1936 †

Material examined: MALTA: Wied Hesri, 22.iv.2016, 1 ♀ (sm) on *Capparis orientalis*, GD; Popeye Village, 29.iv.2017, 1 ♀ (sm) on *Tamarix africana*, GD. GOZO: Ramla Bay, 18.iv.2017, 2 ♀♀ and 1 ♂ (sm) on *Cakile maritima* and 2 ♀♀ (sm) on *Medicago marina*, 18.iv.2017, GD.

Body length: ♀: 1960–2550 µm; ♂: 1780 µm

Wing type: Both sexes are macropterous.

Aeolothrips melisi was recorded on flowers of unrelated plants [14] including Brassicaceae [22,23] and Fabaceae [30] and this was also the case with the material collected from the Maltese Islands. This species was recorded as a facultative predator [29]. *A. melisi* is widespread throughout the Mediterranean basin [13,19].

Aeolothrips tenuicornis Bagnall, 1926

Material examined: MALTA: Għammieri, 19.xii.1996, on *Galium* sp., 2 ♀♀ (sm), DM; M'Scala, 02.ii.1997, on *Hedysarum coronarium*, 2 ♀♀ (sm), DM; Siġġiewi (private garden), 04.xi.2015, 2 ♀♀ (sm) on *Rosa* sp., GD; Siġġiewi (private garden), 28.x.2015, 1 ♀ (sm) on *Rosa* sp., GD; Wied Hesri, 15.i.2016, 1 ♂ (sm) on *Silene colorata*, GD; Siġġiewi (private garden), 29.iii.2016, 4 ♀♀ (sm) and 2 ♂♂ (sm) on *Ranunculus asiaticus*, GD; Siġġiewi (road), 04.iv.2016, 1 ♀ and 1 ♂ (sm) copula pair on *Glebionis coronaria*, GD; Wied Hesri, 15.iv.2016, 1 ♀ (sm) on *Convolvulus arvensis*, GD; Wied Qirda, 06.iv.2016, 2 ♀♀ (sm) on *Glebionis coronaria*, GD; Wied Hesri, 22.iv.2016, 1 ♀ (sm) on *Capparis orientalis*, GD; Kuncizzjoni, 22.iv.2016, 1 ♂ (sm) on *Reichardia picroides*, GD; Siġġiewi (road), 23.v.2016, 1 ♀ (sm) on *Glebionis coronaria*, GD; Siġġiewi (road), 21.vi.2016, 3 ♀♀ (sm, aga) on *Glebionis coronaria*, GD; Msida, Junior College grounds, 08.x.2016, 1 ♀ (sm) on *Cynodon dactylon*, GD; Lapsi 09.i.2017, 1 ♀ (sm) on *Periploca angustifolia*, GD; Siġġiewi (road), 21.iv.2017, 1 ♀ (sm) on *Argyranthemum frutescens*, GD; Mellieħa road l/o Popeye Village, 29.iv.2017, 1 ♂ (sm) on *Convolvulus althoides*, GD; Wied Hesri, 30.iv.2017, 1 ♀ (sm) on *Gladiolus communis*, GD; Pembroke, 05.xi.2018, 1 ♀ (sm) on *Reichardia picroides*, GD. GOZO: Ramla Bay, 18.iv.2017, 1 ♀ (sm) on *Malva arborea*, GD.

Body length: ♀: 1900–2400 µm; ♂: 1200–1680 µm.

Wing type: Both sexes are macropterous.

Adult thrips of this species are found on flowers of unrelated plant species including Asteraceae, Brassicaceae, Fabaceae, [15,23,24,33,36], *Citrus* (Rutaceae) [24] and *Cratageus* (Rosaceae) [28,29], *Triticum aestivum* (Poaceae) and also on *Verbascum* (Scrophulariaceae) [24],

Vitis (Vitaceae). In the Maltese Islands, this species was also found on flowers of unrelated indigenous plants including Asteraceae, Convolvulaceae, Fabaceae and Ranunculaceae. *Aeolothrips tenuicornis* is recorded as a facultative predator [32]. It occurs throughout most of Europe, the Middle East, the Mediterranean Region and the Atlantic Islands (Azores, Madeira, Canaries) [13,14,20,22,28,31].

***Franklinothrips megalops* (Trybom, 1912) †**

Material examined: MALTA: Msida, University of Malta grounds, 04.xi.2016, 1 ♀(sm) on *Rosmarinus officinalis*, DM; Fawwara, 22.v.2017, 1♀(sm) from Malaise trap, DM.

Body length: ♀: 2380–2480 µm, ♂: no records.

Wing type: ♀: macropterous; ♂: no records.

Franklinothrips megalops is a predatory species [37], and has been observed to feed on other small arthropods including juvenile and adult stages of other Thysanoptera such as *Heliothrips hemorrhoidalis* and *Thrips* spp [14,23]. This species was described originally from Kenya but is in fact widespread in Africa [36], Spain and Israel, Yemen and southern India [14,23,36].

***Rhipidothrips brunneus* Williams, 1913 †**

Material examined: MALTA: Wied Ħesri, 24.ii.2017, 2 ♀♀(micropterous) (sm) on *Bromus diandrus*, GD.

Body length: ♀: 1880–2060 µm; ♂: no records.

Wing type: ♀: micropterous; ♂: no records.

This phytophagous species feeds on flowers of Poaceae [14,23,24,27,28,34] but was also recorded as having predatory tendencies [27]. *R. brunneus* is distributed throughout the western Palaearctic and the Near East, and has been accidentally introduced to North America and Australia [13,14,23] where it is an established species [28].

***Rhipidothrips gratiosus* Uzel, 1895 †**

Material examined: MALTA: Wied Ħesri, 03.iv.2016, 1 ♀(sm) on *Avena sterilis*, GD; Wied Ħesri, 22.iv.2016, 2 ♀♀(sm) on *Avena sterilis*, GD; Fiddien, 14.iv.2016, 1 ♀(sm) on *Avena* sp., GD; Wied Ghollieqa 02.xii.2016, 1 ♂(micropterous–sm) on *Cynodon dactylon*, GD; Wied Ħesri, 03.iv.2017, 4 ♀♀(sm, aga) on *Gladiolus communis*, GD; Fiddien, 24.iv.2017, 1 ♀(sm) on *Avena* sp., 1 ♀(sm) on *Medicago* sp., GD; Wied Qirda, 17.iii.2017, 1 ♀(sm) on *Medicago* sp., GD.

Body size: ♀: 1900–2120 µm; ♂: 1240 µm.

Wing type: ♀: macropterous; ♂: micropterous.

Males of this species have been described as being macropterous just like females [14,26], however, the male specimen examined from the Maltese Islands was found to be micropterous and marginally smaller than other males described from continental Europe [14]. *Rhipidothrips gratiosus* is phytophagous, typically found on the flowers of Poaceae [26,27,34,38,39] but also on other plants such as *Erysimum cheiranthoides* (Brassicaceae) [39] and *Vicia faba* (Fabaceae) [40]. In the Maltese Islands, this species has been largely collected from Poaceae. *R. gratiosus* was also recorded as having predatory habits [26]. The distribution of this species extends across the Western Palaearctic, the Near East and North Africa and has been introduced [13] and has become established in North America [14,24].

***Rhipidothrips niveipennis* Reuter, 1899 †**

Material examined: MALTA: Wied Ħesri, 04.iv.2016, 1 ♀(sm) on *Avena sterilis*, GD; Wied Ħesri, 22.iv.2016, 1 ♀(sm) on *Avena sterilis*, GD; Gudja, 1/o Malta International Airport, 26.ix.2016, 1 ♀(sm) on *Ficus microcarpa*, GD.

Body size: ♀: 2000–2014 µm; ♂: no records

Wing type: ♀: macropterous; ♂: no records

Although material from the Maltese Islands consisted of only macropterous specimens, micropterous individuals have been described from continental Europe [14]. This species is phytophagous on different Poaceae [14] as was also largely the case for the specimens collected from the Maltese Islands. *R. niveipennis* is recorded from Finland, Sweden, the French Alps, [14] and Norway [41].

Rhipidothrips unicolor zur Strassen, 1965 †

Material examined: MALTA: Ghajn Tuffieha, 29.iv.2017, 7 ♀♀(sm, aga) and instar larva (sm) on *Stipa capensis*, GD.

Body size: ♀: 1760–2250 µm; ♂: n/a.

Wing type: Both sexes are macropterous.

Rhipidothrips unicolor has been recorded, often in large numbers, on *Stipa capensis* (Poaceae) which has also been described as the host plant for this species [14,42]. In the Maltese Islands, both adults and larvae were also collected from the mentioned host plant. The geographical distribution for *R. unicolor* includes Greece, Italy, Spain, Canary Islands [14,42,43], Morocco [13,14] and Iran [31].

FAMILY MELANTHRIPIDAE

Melanthrips ficalbii Buffa, 1907 †

Material examined: GOZO: Victoria, (road), 20.iv.2019, 2 ♀♀(sm) on *Galium aparine*, GD.

Body length: ♀: 1840–1960 µm; ♂: n/a.

Wing type: Both sexes are macropterous.

Melanthrips ficalbii is a phytophagous species with larval stages being found on Rubiaceae [33] indicating that these plants are used as hosts by this species. Adults have also been recorded on *Galium* (Rubiaceae), *Euphorbia cyparissias* (Euphorbiaceae), *Reseda* (Resedaceae) as well as on different Poaceae [14]. In the current study, *M. ficalbii* has been recorded from *Galium aparine* (Rubiaceae). *M. ficalbii* is widespread in western Europe and the western Mediterranean Region [13,14,25,35].

Melanthrips fuscus Sulzer, 1776

Material examined: MALTA: Siggiewi (private garden), 04.xi.2015, 3 ♂♂(sm) on *Rosa* sp., GD; Siggiewi (private garden), 12.i.2016, 3 ♀♀ and 3 ♂♂(sm) on *Rosa* sp., GD; Siggiewi (private garden), 15.i.2016, 2 ♀♀(sm) and 2 ♂♂(sm) on *Mercurialis annua*, GD; Wied Hēsri, 15.i.2016, 1 ♀(sm) on *Silene colorata*, GD; Wied Qirda, 29.i.2016, 2 instar larvae (aga) and 1 ♂(sm) on *Brassica rapa*, GD; Siggiewi (private garden), 29.iii.2016, 1 ♀(sm) and 1 ♂(sm) on *Ranunculus asiaticus*, GD; Siggiewi (private garden), 16.x.2016, 1 ♀(sm) on *Lobularia maritima*, GD; Wied Qirda, 17.iii.2017, 1 ♂(sm) on *Asphodelus ramosus*, GD; Wied Hēsri, 03.iv.2017, 2 ♀♀(sm) and 1 ♂(sm) on *Acacia saligna*, GD; Dingli Cliffs, 27.iv.2017, 1 ♂(sm) on *Brassica rapa*, GD; Lapsi, 09.x.2017, 1 ♂(sm) on *Potentilla reptans*, GD; Wied Baqqiegha, 22.i.2018, 1 ♀(sm) on *Cerintho major*, GD.

Body length: ♀: 1540–2140 µm; ♂: 1360–1560 µm.

Wing type: Both sexes are macropterous.

This species is phytophagous with overwhelming records from plants in the Brassicaceae [14,23,24,26,32,34,36]. It has also been recorded from annual plants [24,32,34,36,44], grasses and trees [45]. In The Maltese Islands, larvae were found on *Brassica rapa* (Brassicaceae). *Melanthrips fuscus* adults were collected together with *M. lybicus* on flowers of many unrelated indigenous and cultivated plants including Brassicaceae, Rosaceae, Caryophyllaceae, Xanthorrhoeaceae and Fabaceae. *M. fuscus* is found across Europe, the East Palaearctic, the Nearctic and North Africa [22,23]. It has also been intercepted in US from plant material imported from Europe, the Mediterranean and Africa [35].

Melanthrips knechteli Priesner, 1936 †

Material examined: MALTA: Siggiewi (private garden), 15.i.2016, 1 ♀(sm) on *Mercurialis annua*, GD; Siggiewi (private garden), 29.iii.2016, 1 ♂(sm) on *Ranunculus asiaticus*, GD; Wied Xkora, 30.x.2017, 1 ♂(sm) on *Mercurialis annua*, GD; Wied Baqqiegħa, 22.i.2018, 1 ♀(sm) on *Cerinth major*, GD.

Body length: ♀: 1580 µm; ♂: 1160–1240 µm.

Wing type: Both sexes are macropterous.

Melanthrips knechteli has been recorded on *Cerinth major* (Boraginaceae) and *Thymus* (Lamiaceae) [14,23]. In the Maltese Islands, this species was found on *Cerinth major* but also on *Ranunculus asiaticus* (Ranunculaceae) and *Mercurialis annua* (Euphorbiaceae). The geographical distribution for this species includes Turkey, Bulgaria, Romania, Albania, Czech Republic (Moravia), Spain and Iran [14,26,46,47].

Melanthrips libycus Priesner, 1936

Material examined: MALTA: Siggiewi (private garden), 12.i.2016, 1 ♀(sm) and 1 ♂(sm) on *Rosa* sp., GD; Wied Ħesri, 15.i.2016, 1 ♀(sm) and 2 ♂♂(sm) on *Silene colorata*, GD; Wied Qirda, 29.i.2016, 1 ♀(sm) on *Trifolium nigrescens*, 1 ♀(sm) on *Brassica* sp., GD; Qormi (private farm), 07.ii.2016, 1 ♀(sm) on *Brassica oleracea* var. *botrytis*, GD; Wied Qirda, 24.ii.2016, 1 ♀(sm) on *Bromus diandrus*, GD; Ta' Qali, 27.ii.2016, 5 instar larvae (aga) and 2 ♀♀(sm) on *Diplotaxis tenuifolia*, SF; Siggiewi (private garden), 29.ii.2016, 1 ♀(sm) on *Ranunculus asiaticus*, GD; Wied Qirda, 17.iii.2017, 1 ♀(sm) on *Medicago* sp., GD; Wied Ħesri, 03.iv.2017, 1 ♀(sm) on *Acacia saligna*, GD; Dingli Cliffs, 29.x.2017, 1 ♀(sm) on *Brassica rapa*, GD. GOZO: Ramla Bay, 18.iv.2017, 1 ♂(sm) on *Cakile maritima*, GD.

Body length: ♀: 1600–2300 µm; ♂: 1400–1725 µm.

Wing type: Both sexes are macropterous.

The long setae present on the head of this species were described to be as long as three pairs of post-ocular setae [14,48] but in specimens collected from the Maltese Islands these setae were found to be shorter. *Melanthrips libycus* has been recorded on Poaceae [24], Brassicaceae [15,23,44,48] and also from *Citrus x paradisi* (Rutaceae) [15], though the literature does not specify which species are used as host plants. In the current study, larvae of this species were found on *Diplotaxis tenuifolia* (Brassicaceae). *M. libycus* adults together with *M. fuscus* has been recorded on a variety of unrelated plants such as Brassicaceae, Fabaceae and Poaceae. *M. libycus* is widely distributed across southern Europe and the Mediterranean basin [14,15,23,49].

FAMILY STENUROTHRIPIDAE

Holarthrothrips tenuicornis Bagnall, 1927 †

Material examined: MALTA: Qormi (roundabout), 14.ix.2017, 13 ♀♀(sm, aga) on male *Phoenix dactylifera*, GD; Qormi (roundabout), 03.x.2017, 13 ♀♀(sm, aga) on male flowers of *Phoenix dactylifera*, GD; Msida, Junior College grounds, 10.x.2017, 1 ♀ on male flowers of *Phoenix dactylifera*, GD; Siggiewi (private garden), 30.vii.2018, 2 ♀♀(sm), 10 ♀♀(aga) and 1 ♂(sm), 1 ♂(aga) on male flowers of *Phoenix dactylifera*, GD.

Body length: ♀: 1550–1820 µm; ♂: 1260 µm.

Wing type: Both sexes are macropterous.

Thrips of the genus *Holarthrothrips* are mostly associated with male flowers of *Phoenix* trees (Arecaceae) [15,50] where both larvae and adults occur. Adults of this species have also been recorded on other plants such as *Vitis* (Vitaceae), *Citrus* (Rutaceae), and *Rubia* (Rubiaceae) [14,24,29] which, however, should not be considered as host-plants. In the Maltese Islands larvae and adults were always found on male flowers of *Phoenix dactylifera*. *H. tenuicornis* has been recorded from across the Mediterranean basin [13–15,23,29,43].

FAMILY THRIPIDAE

Subfamily Dendrothripinae

Dendrothrips saltator Uzel, 1895 †

Material examined: MALTA: Wied Babu, 15.xii.1996, 7 ♀♀(sm) on *Ferula melitensis*, DM; Wied Qirda, 06.iv.2016, 13 instar larvae (aga) on *Foeniculum vulgare*, GD; Maqluba, 1/o Qrendi, 08.iv.2016, 9 ♀♀(sm, aga) on *Foeniculum vulgare*, GD; Qormi (road), 19.iv.2016, 1 instar larva (aga) on *Ferula melitensis*, GD; 09.vii.2018, 6 ♀♀(sm, aga) and 1 ♂(sm) on *Foeniculum vulgare*, GD.

Body length: ♀: 1260–1440 µm; ♂: 860 µm.

Wing type: Both sexes are macropterous.

Dendrothrips saltator has been reported from leaves of various deciduous trees and a variety of unrelated plants [14,26], particularly on Apiaceae, Asteraceae [14,23,26], Fabaceae, as well as plants such as *Triticum aestivum* (Poaceae) and *Olea europaea* (Oleaceae) [34]. In the Maltese Islands, both adults and larvae of *D. saltator* were found on *Foeniculum vulgare* and *Ferula melitensis* (both in Apiaceae). Thus, most likely, the host plants for *Dendrothrips saltator* are only plants within the family Apiaceae. In fact, in the UK, this species is only recorded on *Paucedanum officinale* (Apiaceae) [27]. *D. saltator* is widespread in Europe, the Oriental Region and the Near East [13,14]. It was also intercepted in the US on plant material originating from Europe [35,51].

Subfamily Panchaetothripinae

Heliothrips haemorrhoidalis (Bouché, 1883)

Material examined: MALTA: Żabbar, 06.v.1996, 1 ♀(sm) on *Viburnum* sp., CF; Msida, University of Malta grounds, 28.iv.2017, 1 ♀(sm) on *Apium graveolens*, GD.

Body length: ♀: 1660–1900 µm; ♂: no records.

Wing type: ♀: macropterous; ♂: no records.

Heliothrips haemorrhoidalis is a polyphagous species recorded on different unrelated host-plants [23,24,26,32]. It is an agricultural pest and can potentially inflict damage on a number of crop plants in both greenhouses [23] and open fields [26]. In the Maltese Islands *H. haemorrhoidalis* was collected on *Viburnum* (Adoxaceae), *Apium graveolens* (Apiaceae) as well as from plants with low levels of nitrogen [12]. This species is preyed upon by *Franklinothrips megalops* [26]. *H. haemorrhoidalis* is believed to have originated in South America, probably Peru [51], but has become widespread around the world in tropical and subtropical areas [24], being introduced with exotic ornamental plants. In northern Europe, *H. haemorrhoidalis* thrives in greenhouses and indoor places [14].

Hercinothrips femoralis (Reuter, 1881) †

Material examined: MALTA: Msida, University of Malta grounds, 04.v.2016, 1 ♀(sm) on *Origanum majorana*, GD; Siġġiewi (private garden), 05.xi.2016, 14 ♀♀(sm, aga) and 1 instar larva (aga) on *Hippeastrum* sp., GD; Birguma, 20.viii.2020, 13 ♀♀(sm, aga) on *Ocimum basilicum*, NY; Siġġiewi private garden, 04.xi.2020, 8 ♀♀(sm, aga) on *Calendula officinalis*, GD.

Body length: ♀: 1320–1580 µm; ♂: no records.

Wing type: ♀: macropterous; ♂: no records.

Hercinothrips femoralis is a polyphagous species occurring on herbs and cultivated plants, mostly in greenhouses [23,24] and often on Amaryllidaceae [14] but also on other crops [26]. In the Maltese Islands, *H. femoralis* was found on different unrelated plants but only the *Hippeastrum* sp. (Amaryllidaceae) can be regarded as a host-plant for the mentioned thrips. Although this species was recorded in Europe from greenhouses, it was found in plants grown outside in the Maltese Islands, probably due to the warmer climate of the archipelago. *H. femoralis* has been recorded as a pest, inflicting damage in greenhouse crops [27]. *H. femoralis* is native to Central Africa, but is now subcosmopolitan in distribution with many records in tropical and subtropical countries [14,27,35].

Subfamily Thripinae

Anaphothrips sudanensis Trybom, 1911 †

Material examined: MALTA: Wied Hesri, 04.xi.2016, 1 ♀(sm) on *Cynodon dactylon*, GD.

Body length: ♀: 1340 µm; ♂: no records.

Wing type: the locally collected female specimen is macropterous.

Anaphothrips sudanensis is a phytophagous species associated with Poaceae [23]. It is commonly found in the leaf axils and tend to develop streaks on leaves as these expand and mature [28]. This species has been recorded on different Poaceae, particularly on *Panicum maximum*, *Pennisetum*, *Sorghum halepense*, *Zea mays* and *Saccharum officinarum* [14,26,28], as well as on *Allium cepa* (Amaryllidaceae) [24] and *Tagetes minuta* (Asteraceae) [26]. In the Maltese Islands, the single female specimen of this species was collected on Poaceae (*Cynodon dactylon*). *A. sudanensis* is recorded from the East Palaearctic, Afrotropical, Australian, Neotropical and Oriental Regions, the Near East and North Africa [13].

Aptinothrips rufus (Haliday, 1836) †

Material examined. MALTA: Wied Hesri, 04.iv.2016, 1 ♀(sm) and 1 instar larva (aga) on *Hyparrhenia hirta* and 4 ♀♀(sm) and 1 instar larva (aga) on *Avena sterilis*, GD; Wied Qirda, 06.iv.2016, 1 ♀(sm) on *Plantago major*, GD; Maqluba l/o Qrendi, 15.iv.2016, 8 ♀♀(sm) on *Triticum aestivum*, GD; Wied Hesri, 15.iv.2016, 1 ♂(sm) on *Convolvulus arvensis*, GD; Msida, Junior College grounds, 27.iv.2016, 1 ♀(sm) and 2 instar larvae (aga) on *Hordeum leporinum*, GD; Kuncizzjoni, 22.i.2017, 1 ♀(sm) on *Hordeum leporinum*, GD; Msida, Junior College grounds, 03.iv.2017, 2 ♀♀(sm) on *Hordeum leporinum*, GD; Ghajn Tuffieħa, 29.iv.2017, 1 ♀(sm) on *Stipa capensis*, GD. GOZO: Ramla Bay, 18.iv.2017, 1 ♀(sm) on *Medicago marina*, GD.

Body length: ♀: 1460–1700 µm; ♂: 900 µm.

Wing type: Both sexes are apterous.

Aptinothrips rufus is associated with Poaceae [15,22,26,44,52] but adults were recorded on unrelated plants belonging to the following families: Fabaceae, Lamiaceae, Plumbaginaceae [44], Polygonaceae [34], and Asteraceae [26]. In the Maltese Islands, both adults and larvae of *A. rufus* were only found on Poaceae (*Avena* sp., *Hordeum leporinum* and *Hyparrhenia hirta*), though adults were also found on other plants as well as grasses. *A. rufus* is a species that originated from Europe, but has become widespread in temperate regions around the world [14,32,52]. In fact, it is recorded from the East Palaearctic, Afrotropical, Australian, Nearctic, Neotropical and Oriental Regions, the Near East and North Africa [13]. It has been intercepted in the US from plants originating from Europe, the Mediterranean and Africa [34,53].

Asphodelothrips croceicollis (Karny, 1914) †

Material examined: MALTA: Dingli Cliffs, 27.i.2016, 4 ♂♂(sm) on *Asphodelus ramosus*, GD; Buskett, 19.ii.2016, 2 instar larvae (aga) on *Asphodelus ramosus*, GD; Dingli Cliffs, 24.i.2017, 5 ♀♀(sm) on *Asphodelus ramosus*, GD; Kuncizzjoni, 28.i.2017, 2 ♀♀(sm) and 1 ♂(sm) on *Asphodelus ramosus* (ag) and 1 ♀(sm) on *Erica multiflora*, GD; Wied Hesri, 24.ii.2017, 1 ♀(sm) and 1 ♂(sm) and on *Asphodelus ramosus*, GD; Buskett, 23.ii.2018, 2 ♀♀(sm) on *Asphodelus ramosus*, GD; Kuncizzjoni, 29.i.2018, 1 ♀(sm) on *Glebionis coronaria*, GD.

Body length: ♀: 1760–2160 µm; ♂: 1000–1580 µm.

Wing type: Both sexes of locally recorded specimens are micropterous.

Both micropterous as well as macropterous forms for this species exist, with the macropterous females being dark brown in colour and the micropterous forms being light brown [14]. All specimens collected from Malta were found to be micropterous, with females being of a dark brown colour. *Asphodelothrips croceicollis* is associated with *Asphodelus* spp. (Asphodelaceae) [14,29] and in the Maltese Islands it was almost exclusively collected on flowers of *Asphodelus ramosus*. *A. croceicollis* is found throughout the Ponto-Mediterranean area and the Canary Islands, occurring up to 1500 m altitude [14], as well in the Near East [13].

Bregmatothrips dimorphus (Priesner, 1919) †

Material examined: MALTA: Wied Hesri, 15.i.2016, 5 ♀♀(aga) and 4 ♂♂(sm, aga) on *Pipatherum miliaceum*, GD; Buskett, 03.ii.2016, 1 ♀(sm) on *Cynodon dactylon*, 3 ♀♀(sm, aga) and 2 ♂♂(sm, aga) on *Pipatherum miliaceum*, GD.

Body length: ♀: 1440–1460 µm; ♂: 1140–1240 µm.

Wing type: ♀: macropterous; ♂: micropterous.

The female specimens collected in the Maltese Islands were all macropterous, but micropterous forms are also described from the European continent [32]. *Bregmatothrips dimorphus* is associated with Poaceae [34] and this was also the case for the material collected from the Maltese Islands which was found on the grass species *Cynodon dactylon* and *Pipatherum miliaceum*. This species is widespread in Europe but is also recorded from Yemen, Sudan, southern Africa [14,23,26], the Near East and North Africa [13].

Ceratothrips ericae (Haliday, 1836) †

Material examined: MALTA: Lapsi, 9.x.2017, 1 ♀(sm) on *Limbarda crithmoides*, GD. GOZO: Qbajjar, 31.iii.2018, 1 ♀(sm) on *Helychrysum melitense*, GD.

Body length: ♀: 1200–1460 µm; ♂: no records.

Wing type: ♀: macropterous; ♂: no records.

Different authors have recorded this species from unrelated plants [14,27,29,35,54,55]. In the Maltese Islands, *Ceratothrips ericae* was found on *Avena sterilis* (Poaceae) and *Helychrysum melitense* (Asteraceae). *C. ericae* is an important pollinator, especially in areas where larger pollinators do not occur [56]. This species is of Euro-Siberian origin, but has been accidentally introduced to other parts of the world [14], including the Near East and the Australian Region [13]. It has also been intercepted in the US from plants originating from Europe [35].

Chirothrips hamatus Trybom, 1895 †

Material examined: GOZO: Xlendi, 21.vii.1956, 1 ♀(sm) on *Lithospermum arvense* (JIS & ERS), (BMNH).

Body length: ♀: 1453 µm; ♂: no records.

Wing type: ♀: macropterous; ♂: no records.

Males of this species are described as being micropterous [14,24,28]. *Chirothrips hamatus* is mostly associated with Poaceae, particularly *Alopecurus pratensis* [14,24,27] and thus, *Lithospermum arvense* (Boraginaceae), on which the single female specimen was collected during the present study, cannot be considered as a host plant. *C. hamatus* is recorded from western Siberia, the Near East and the Nearctic Region; it is also widespread in Europe being generally less common in southern areas [13,14,27].

Chirothrips manicatus (Haliday, 1836) †

Material examined: MALTA: Siggiewi (private farm), 03.v.2016, 1 ♀ and 1 ♂(sm) on *Koeleria cristata*, GD; Wied Hesri, 02.xi.2016, 1 ♂(sm) on *Phragmites australis*, GD; Wied Hesri, 04.xi.2016, 4 ♀♀(sm, aga) and 5 ♂♂(sm, aga) on *Arundo donax*, GD; Naxxar (road), 29.iv.2017 1 ♂(sm) on *Triticum aestivum*, GD; Wied Qirda, 31.v.2018, 1 ♂(sm) on *Hyparrhenia hirta*, GD; Fiddien 16.vi.2021, 1 ♀(sm) on *Avena sterilis*, GD.

Body length: ♀: 1400–1700 µm; ♂: 960–1440 µm.

Wing type: ♀: macropterous; ♂: micropterous.

Chirothrips manicatus is a species which is mostly associated with Poaceae and sometimes Cyperaceae [23]. It has been also recorded on other unrelated plants such as *Malva alcea* (Malvaceae) and *Malus* sp. (Rosaceae) [54,57–59]. In the Maltese Islands, this species was only collected from Poaceae. *C. manicatus* may transmit phytopathogenic bacteria and fungi [25]. *Chirothrips manicatus* probably originated in the Palaearctic but has a Holarctic distribution and is nowadays considered as subcosmopolitan in distribution, with numerous records in temperate parts of the world [14].

Chirothrips meridionalis Bagnall, 1927 †

Material examined: MALTA: Wied Ħesri, 04.iv.2016, 2 ♂♂(sm) on *Hyparrhenia hirta*, GD. Wied Qirda, 04.xi.2016, 2 ♀♀(sm) on *Arundo donax*, GD; Wied Qirda, 31.v.2018, 3 ♀♀(sm) on *Hyparrhenia hirta*, GD (1 ♀ag).

Body length: ♀: 1600–1680 µm; ♂: 1240 µm.

Wing type: Both sexes are macropterous.

Similar to other species of the genus *Chirothrips*, *C. meridionalis* is also associated with Poaceae [14,22,23,26]. In the Maltese Islands this thrips was also collected on plants belonging to the mentioned plant family. *C. meridionalis* has been recorded from South Europe [22] and is widespread across the Mediterranean basin, Madeira, Canary Islands, India, Pakistan, Iran, Yemen, Nigeria and South Africa [14]. It has also been intercepted in the US from plants originating from Europe, the Mediterranean and Africa [35].

Echinothrips americanus Morgan, 1913 †

Material examined: MALTA: Siġġiewi (private farmhouse), 21.v.2017, 1 ♀(sm) on *Salvia officinalis*, GD; Siġġiewi (private garden), 14.vii.2017, 13 ♀♀(sm, aga) on *Azalea indica*, Siġġiewi (private garden), 16.vii.2017, 8 ♀♀, 3 instar larvae (aga) on *Azalea indica*, GD.

Body length: ♀: 1100–1340 µm; ♂: no records.

Wing type: ♀: macropterous; ♂: no records.

This phytophagous species has been recorded from unrelated plants including many vegetables [50], *Impatiens* (Balsaminaceae), *Dendranthema* (Asteraceae) and *Euphorbia* (Euphorbiaceae) [32]. It is believed to have been introduced to many countries around the world with trade of ornamental plants. This species has been recorded as a pest on cultivated plants such as *Philodendron* (Araceae) and *Syngonium* (Araceae), *Impatiens*, and *Euphorbia* [32,48]. *Echinothrips americanus* is native to eastern North America and has been imported with ornamental plants to Europe [14,32], Neotropical and Oriental Regions [13].

Frankliniella occidentalis (Pergande, 1895)

Material examined: MALTA: Żabbar, 11.iii.1994, 2 ♀♀(sm) (BMNH) on *Gerbera* sp., DM; St. Paul's Bay, 14.iii.1994, 2 ♀♀(sm) (BMNH) on *Solanum melanogena*, GW; St. Paul's Bay, 14.iv.1994, 1 ♀(sm) (BMNH) on *Dianthus caryophyllus*, JWI; St. Paul's Bay, 14.iv.1994, 2 ♀♀(sm) (BMNH) on *Chrysanthemum* sp., JWI; St. Paul's Bay (glasshouse), 14.iii.1994, 2 ♀♀(sm) (BMNH) on *Fragraria x ananassa*, GW; Żabbar, 13.i.1997, 1 ♀(sm) (dark morph) on *Dianthus caryophyllus*, DM; Siġġiewi (road), 14.x.2015, 1 ♀(sm) (yellow morph) on *Diplotaxis tenuifolia*, GD; Siġġiewi (private garden), 29.i.2016, 1 ♀(sm) (yellow morph) on *Narcissus tazetta*, GD; Wied Qirda, 06.iv.2016, 1 ♂(sm) on *Matricaria chamomilla*, GD; Wied Ħesri, 04.iv.2016, 1 ♂(sm) on *Avena sterilis*, GD; Siġġiewi (private garden), 13.iv.2016, 1 ♀(sm) (yellow morph) on *Foeniculum vulgare*, GD; Manikata, 21.iv.2017, 1 ♂(sm) on *Pallenis spinosa*, GD; Mellieħa (road l/o Popeye Village), 21.iv.2017, 1 ♀(sm) (yellow morph), 1 ♀(sm) (dark form) on *Tamarix africana*, GD; Il-Ballut, l/o M'Xlokk, 05.v.2017, 1 ♀(sm) (pale morph), 1 ♀(sm) (dark morph) on *Malva arborea*, GD; road, l/o Wied Ħesri, 07.viii. 2017, 1 ♀(sm) (pale morph), 3 ♀♀(aga) on *Ipomoea carnososa*, GD; Qormi (road), 18.viii.2017, 1 ♀(sm) (pale morph), 3 ♀♀(aga) (pale morph) on *Yucca gloriosa*, GD; San Ġwann (private farm), 29.x.2018, 2 ♀♀(sm) (yellow morph), 3 ♀♀(aga) (pale form) and 1 ♂ on *Lactuca sativa*, GD; Wied Ħesri, 12.iii.2018, 2 ♀♀(sm) (dark morph) on *Trifolium nigrescens*, GD; Fiddien, 16.v.2021, 1 ♀(sm) on *Avena sterilis*, GD.

Body length: ♀: 1440–1800 µm; ♂: 1080–1300 µm.

Wing type: Both sexes are macropterous.

The extent of the brown color pattern on antennae and overall color intensity of this species is likely dependent on temperature, with darker specimens developing in cooler temperatures [27]. Both dark and yellow morphs were collected from the Maltese Islands. *Frankliniella occidentalis* is polyphagous, with adults and larvae commonly found on both flowers and leaves of unrelated plants [29] including numerous crops [23] and indigenous

plants [27,35,40,54,55,59]. In the Maltese Islands this species was recorded on seed crops, cut flowers such as *Dahlia* (Asteraceae), *Dianthus* (Caryophyllaceae) and *Gerbera* (Asteraceae), nursery stock, peaches, plums, nectarines, strawberries, sweet peppers, grapes, cotton and other crops including tomatoes [12]. This species is particularly harmful to crops, fruits and cultivated flowers and is a vector of a number of orthotospoviruses such as Tomato Spotted Wilt Virus (TSWV). Nonetheless, *F. occidentalis* has also been described to feed on the eggs and adults of mites species which can be harmful to agricultural crops [60]. *F. occidentalis* is native to Pacific North America, but has been imported with cultivated plants to virtually all countries in the world during the early 1980s, thriving at first in greenhouses but subsequently spreading out in the open especially in areas with warmer climates such as the Mediterranean basin [14,32].

Frankliniella schultzei (Trybom, 1910) †

Material examined: MALTA: Siggiewi (private garden), 04.xi.2015, 1 ♀(sm) (yellow morph form) on *Rosa* sp.

Body length: ♀: 1400 µm; ♂: no records.

Wing type: ♀: mactopterous; ♂: no records.

The specimen collected from the Maltese Islands was yellow but dark morphs of this species also exist as in the case of *Frankliniella occidentalis*. *F. schultzei* was recorded on flowers and leaves of a wide variety of different unrelated plants [14,25,26,30,33,61], crops [25,48] and stored bulbs [28]. It is known to cause damage to different crops and is an orthotospovirus vector [14,25]. *F. schultzei* is generally considered as native to South America, but it could also be of Africotropical origin [26]. It is widespread in circum-tropical and subtropical regions, where it is recorded as a common pest [14].

Limothrips angulicornis Jablonowski, 1894 †

Material examined: MALTA: Fiddien, 24.iv.2017, 1 ♀(sm) on *Avena* sp., GD.

Body length: ♀: 2080 µm; ♂: no records.

Wing type: ♀: macropterous; ♂: no records.

Limothrips angulicornis is phytophagous and associated with leaves and leaf axils of grasses [14,44]. Larvae have been found on Poaceae [62] but adults were also recorded from *Populus* sp. (Salicaceae) and *Pinus* sp. (Pinaceae) [34]. In the Maltese Islands, this thrips was collected from *Avena* (Poaceae). *L. angulicornis* originates from the West Palaearctic, but has been imported to other parts of the world such as Western Asia [22], California, Chile and southern Australia where it has become established [14,26,27].

Limothrips cerealium (Haliday, 1836) †

Material examined: MALTA: Wied Qirda, 10.iv.2016, 2 ♀♀(sm) on *Bromus diandrus*, GD; Wied Hēsri, 22.iv.2016, 1 ♀(sm) on *Hyparrhenia hirta*, GD; Wied Speranza, 26.iv.2016, 1 ♀(sm) on *Mentha pulegium*, GD; Junior College grounds, 27.iv.2016, 1 ♀(sm) on *Hordeum leporinum*, GD; Dingli Cliffs, 24.v.2016, 1 ♀(sm) on *Brassica rapa*, GD; Dingli Cliffs, 27.v.2016, 1 ♀(sm) on *Glebionis coronaria*, GD; Fiddien, 24.iv.2017, 2 ♀♀(sm) on *Avena* sp., GD; Naxxar, road, 29.iv.2017, 1 ♀(sm) on *Avena sterilis* and 1 ♂(sm) on *Triticum aestivum*, GD; Wied Hēsri, 12.iii.2018, 1 ♀(sm) on *Trifolium nigrescens*, GD.

Body length: ♀: 1520–2200 µm; ♂: 1260 µm.

Wing type: ♀: macropterous; ♂: micropterous.

Limothrips cerealium is phytophagous and generally found on Poaceae, namely cereals [25] and grasses [22,34,35,55,63] where larvae were also recorded [27]. In the Maltese Islands, this species was mainly found on Poaceae, but also on unrelated herbaceous plants such as *Brassica rapa* (Brassicaceae) and *Glebionis coronaria* (Asteraceae). This species can cause damage to grains, causing “silvering” on leaves [23,26,48] and is possibly a mechanical vector of phytopathogenic bacteria and fungi [23]. *L. cerealium* probably originated in Atlantic western Europe [14], but has been accidentally introduced on cereals and grains in many parts of the world and is now considered subcosmopolitan in distribution [14].

Odontothrips meliloti Priesner, 1951 †

Material examined: MALTA: Wied Qirda, 17.iii.2017, 1 ♀(sm) on *Hedysarum coronarium*, GD; Fiddien, 24.iv.2017, 1 ♀(sm) on *Medicago* sp., GD; Il-Ballut l/o M'Xlokk, 05.v.2017, 1 ♀(sm) on *Medicago arborea*, GD; Siggiewi (private residence), 18.iii.2018, 1 ♀ on glass window pane, GD. GOZO: Xlendi, 31.iii.2016, 2 ♀♀(sm) on *Lotus ornithopodioides*, GD; Ramla Bay, 17.iii.2017, 1 ♂(sm) on *Medicago marina*, GD.

Body length: ♀: 1900–2460 µm; ♂: 1500 µm.

Wing type: Both sexes are macropterous.

Odontothrips meliloti is recorded from flowers of Fabaceae, specifically on those of the genus *Melilotus* (*M. albus* and *M. officinalis*) [23,33]. In the Maltese Islands, it was also collected from flowers of Fabaceae (*Hedysarum coronarium*, *Lotus ornithopodioides*, *Medicago* sp.). This species has a Euro-Siberian distribution and is widespread in central and southern Europe [14].

Oxythrips ajugae Uzel, 1895 †

Material examined: MALTA: Wied Ghollieqa, 10.iii.2017, 10 ♀♀(aga, sm) on *Pinus halepensis*, GD; Wied Qirda, 17.iii.2017, 6 ♀♀(aga, sm) on *Pinus halepensis*, GD; Kuncizzjoni, 29.i.2018, 5 ♀♀(sm) and 3 ♂♂(sm) on *Pinus halepensis*, GD; Xemxija, 03.iii.2018, 5 ♀♀(sm) on *Pinus halepensis*, GD.

Body length: ♀: 1500–1700 µm; ♂: 920 µm.

Wing type: Both sexes are macropterous.

Oxythrips ajugae is a spring occurring [14] dendrophilous foliivore [64]. It has been recorded on Pinaceae such as *Pinus* and *Cupressus* [14,29], but also on a number of hardwoods and herbaceous plants [34]. In the Maltese Islands, adults and larvae of this species were found in spring on *Pinus halepensis* (Pinaceae), indicating that this plant is used as a host by the mentioned thrips. *O. ajugae* is widespread in Europe [14] and Asia [32], the Nearctic Region, the Near East and North Africa [13].

Pezothrips kellyanus (Bagnall, 1916) †

Material examined: MALTA: Siggiewi (private garden), 04.xi.2015, 3 ♂♂(sm) on *Rosa* sp.

Body length: ♀: no records; ♂: 1460–1660 µm.

Wing type: ♀: no records; ♂: macropterous.

Pezothrips kellyanus is found on highly scented white flowers [28,65] especially those of *Citrus* (Rutaceae) [48,65,66], but also on *Gardenia* (Rubiaceae) [65], *Jasminum* (Oleaceae) [32,65–67], *Lonicera japonica* (Caprifoliaceae) [67], *Olea europea* (Oleaceae) [34] and *Pittosporum tobira* (Pittosporaceae) [27,67]. Immature stages of this species were only found on lemon, grapefruit, *Jasminum* and *Gardenia* flowers [65], however, the flowers of *Camellia* (Theaceae), *Chrysanthemum* (Asteraceae), *Rosa* (Rosaceae), *Prunus* (Rosaceae), *Jasminum* (Oleaceae), and *Passiflora* (Passifloraceae) can sustain the breeding of this species in times when citrus plants are not in bloom and at times when the climate becomes unfavourable, to then reinfest citrus trees when the climate temperature becomes warmer and the citrus trees bloom again [68]. In the Maltese Islands this species was collected from flowers of *Rosa* sp. (Rosaceae) found in close proximity to *Citrus* trees (Rutaceae). *P. kellyanus* is often recorded as an agricultural pest, causing scarring on fruits, particularly those of citrus trees [48,66]. This species may have originated in Australia [27] but was accidentally introduced in many territories, particularly in Europe [27].

Prosopothrips nigriceps Bagnall, 1927 †

Material examined: MALTA: Pembroke, 05.xi.2018, 1 ♀(sm) on *Reichardia picroides*, GD; Xrobb l-Għagin, 14.iii.2018, 2 ♀♀(aga) on *Hedysarum glomeratum*, GD. GOZO: Ramla Bay, 18.iv.2017, 6 ♀♀(aga, sm) on *Silene colorata*, GD.

Body length: ♀: 1340–1500 µm; ♂: no records.

Wing type: ♀: apterous; ♂: no records.

Prosopothrips nigriceps is mainly recorded from coastal regions on dune grasses such as *Lygeum spartum* and *Poa annua*, as well as on low-growing vegetation [14]. In the Maltese Islands, this species was mostly collected from coastal garigue areas on different unrelated plants such as *Reichardia picroides* (Asteraceae), *Hedysarum glomeratum* (Fabaceae) and *Silene colorata* (Caryophyllaceae). *P. nigriceps* is distributed throughout the Mediterranean basin [14]. It has also been intercepted in the US from plants originating from southern Europe and the Mediterranean [35].

***Tenothrips discolor* (Karny, 1907)**

Material examined: MALTA: St. Paul's Bay, vii.1956, 1 ♀(sm) (BMNH) on *Limbarda crithmoides*, ERS.

Body length: ♀: 1200 µm; ♂: no records.

Wing type: ♀: macropterous; ♂: no records.

Although mostly associated with Asteraceae, *Tenothrips discolor* is regarded as polyphagous [14] and recorded on several unrelated plants [29,34,44,55,57]. In the Maltese Islands, this species is recorded from a specimen in the collection of the BMNH [11] collected on *Limbarda crithmoides* (Asteraceae). This species is known from the Ponto-Mediterranean area extending its distribution range to Kyrgyzstan and the Canary Islands [14].

***Thrips australis* (Bagnall, 1915) †**

Material examined: MALTA: Wied Hesri, 10.x.2015, 6 ♀♀(sm) and 2 ♂♂(sm) on *Sambucus nigra*, GD; Wied Hesri, 16.iv.2016, 1 ♀(sm), 5 ♀♀(aga) and 1 ♂(sm) on *Schinus terebinthifolius*, GD; Dingli Cliffs, 27.i.2017, 1 ♂(sm) on *Asphodelus ramosus*, GD; Wied Musa l/o Siggiewi, 30.x.2017, 1 ♀(sm) on *Smilax aspera*, GD; Rabat road 2, 23.xi.2017, 2 ♀♀(sm) on *Ceratonia siliqua*, GD; Luqa (by-pass), 14.i.2018, 1 ♀(sm) on *Solandra maxima*, GD; Siggiewi (private farm), 29.i.2018, 3 ♀♀(sm) and 5 ♀♀(aga) on *Eriobotrya japonica*, GD.

Body length: ♀: 1330–1540 µm; ♂: 800–1260 µm.

Wing type: Both sexes are macropterous.

Thrips australis is mostly, though not exclusively, found on *Eucalyptus*, and other Myrtaceae, but may disperse to other white flowering plants (e.g., *Melaleuca* sp.) once *Eucalyptus* flowers are no longer available [52]. This species has been also recorded on a number of agricultural crops [26], *Juncus* (Poaceae) [57], *Lantana* (Verbenaceae), *Pentanisia ouranogyne* (Rubiaceae) [26] and *Acacia* (Fabaceae) [26,50], but their presence on these species is probably accidental. In the Maltese Islands, this species was never collected on *Eucalyptus* flowers, but was always found on white flowers of unrelated trees and shrubs. An Australia species in origin, this species is now present wherever *Eucalyptus* plantations were introduced [53].

***Thrips major* Uzel, 1895 †**

Material examined: MALTA: Wied Babu, 15.xii.1996, 2 ♀♀(sm), on *Erica multiflora*, DM; Siggiewi (private garden), 04.xi.2015, 4 ♀♀(sm) on *Rosa* sp., GD; Siggiewi (private garden), 12.i.2016, 1 ♀(sm) on *Rosa* sp., GD; Siggiewi (private garden), 01, 29.i.2016, 4 ♀♀(sm) on *Narcissus tazetta*, GD; Siggiewi (private garden), 21.iii.2016, 2 ♀♀(sm) on *Pittosporum tobira*, GD; Buskett, 19.iv.2016, 1 ♀(sm) on *Laurus nobilis*, GD; Żabbar (private garden), 05.iv.2016, 1 ♀(sm) on *Stephanotis floribunda*, GD; Wied Qirda, 15.iv.2016, 1 ♀(sm) on *Bromus madritensis*, GD; Siggiewi (road), 20.iv.2016, 3 ♀♀(sm) on *Olea europaea*, GD; Wied Speranza, 26.iv.2016, 1 ♀(sm) on *Tropaeolum majus*, GD; Siggiewi (private garden), 10.v.2016, 2 ♀♀(sm) on *Lonicera caprifolium*, GD; Siggiewi, 21.viii.2016, 3 ♀♀(sm) on *Glebionis coronaria*, GD; Msida, University of Malta grounds, 11.x.2016, 3 ♀♀(sm) on *Quercus robur*, GD; Siggiewi (private garden), 21.x.2016, 4 ♀♀(sm) on *Rosa* sp., GD; Kuncizzjoni, 19.i.2017, 1 ♀(sm) on *Erica multiflora*, GD; Dingli Cliffs, 27.i.2017, 2 ♀♀(sm) on *Asphodelus ramosus*, GD; Msida, University of Malta grounds, 28.iv.2017, 2 ♀♀(sm) on *Olea europaea*, GD; Luqa (by-pass), 14.i.2018, 1 ♀(sm) on *Solandra maxima*, GD; Siggiewi (private garden), 14.vii.2017, 1 ♀(sm) on *Azalea indica*, GD; Siggiewi (private farm), 29.i.2018, 4 ♀♀(sm) on *Eriobotrya japonica*, GD; Buskett, 20.ii.2018, 1 ♀(sm) on *Rhamnus alaternus*, GD; Wied Ghollieqa, 11.iv.2018, 2 ♀♀(sm) on *Cercis siliquastrum*,

GD; Wied Għollieqa, 14.iv.2018, 1 ♀(sm) on *Cercis siliquastrum*, GD; Għaxqet l-Għajn, l/o Naxxar, 28.iv.18, 2 ♀(sm) on *Pistacea lentiscus*, GD; San Ġwann (private farm), 05.vii. 2019, 2 ♀♀(sm) on *Cucurbita* sp., GD; Siġġiewi, (private garden), 22.iv.2020, 1 ♀(sm) on *Citrus limon*, GD; Mġiebaħ l/o Xemxija, 2.iii.2018, 2 ♀♀(sm) on *Tamarix africana*, GD; Siġġiewi(private garden), 22.iv.2020, 4 ♀♀(sm) on *Prunus persica*, GD.

Body length: ♀: 1300–1580 µm; ♂: no records.

Wing type: ♀: macropterous; ♂: no records.

Thrips major is a polyphagous species, found on flowers and occasionally on leaf buds of unrelated plants [14,24,26,29,34,35,44,54,55,63]. In the Maltese Islands it was also collected on flowers and leaves of different crops, indigenous shrubs and trees belonging to the families Rosaceae, Amaryllidaceae, Pittosporaceae, Fabaceae, Tamaricaceae, Rhamnaceae, Poaceae and Lauraceae amongst others. *T. major* is distributed across the Palaearctic [14] and has been intercepted in the US from Europe, the Mediterranean basin and Africa [69].

Thrips simplex (Morison, 1930) †

Material examined: MALTA: 3.vi.1959, 1♀(sm—BMNH) on *Gladiolus* sp., ERS; St. Paul's Bay, 14.iv.1994, 1♀, *Gladiolus* sp., JWI;

Body length: ♀: 1697 µm; ♂: no records.

Wing type: ♀: macropterous; ♂: no records.

This species is associated with *Gladiolus* spp. (Iridaceae) [14,27,55,58], though it has also been recorded on *Calla* (Araceae), *Pancreatium* (Amaryllidaceae), *Protea* (Proteaceae), *Vitis* (Vitaceae) [14] and *Dianthus* (Caryophyllaceae) [58]. In the Maltese Islands, *Thrips simplex* was only found on *Gladiolus* (Iridaceae). This species originated from southern Africa [27], but has become cosmopolitan in distribution and occurs wherever *Gladiolus* is cultivated [14].

Thrips tabaci Lindeman, 1889

Material examined: MALTA: St. Paul's Bay, vii.1956, 1 ♀(sm—BMNH) on *Limbarda crithmoides* (JIS & ERS); Żabbar, 11.iii.1994, 2 ♀♀(sm—BMNH) on *Matthiola bicornis*, GW; Armier, 14.vi.1994, 1 ♀(sm—NMNH) on *Fragraria x ananassa*, GW; Lapsi, 26.i.1997, 1 ♀(dark morph) on *Asphodelus ramosus*, DM; Siġġiewi (private garden), 21.iii.2016, 1 ♀(dark morph) on *Pittosporum tobira*, GD; Żabbar (private garden), 05.iv.2016, 1 ♀(dark morph) on *Stephanotis floribunda*, GD; Siġġiewi (private garden), 13.iv.2016, 2 ♀♀(dark morph—sm) on *Foeniculum vulgare*, GD; Wied Ħesri, 16.iv.2016, 1 ♀(dark morph) on *Schinus terebinthifolius*, GD; Dingli, Buskett, 19.iv.2016, 1 ♀(pale form—sm) on *Laurus nobilis*, GD; Wied Speranza, 26.iv.2016, 2 ♀♀(dark morph—sm) on *Tropaeolum majus*, and 3 ♀♀(dark morph—sm) on *Parietaria judaica*, GD; Siġġiewi (private garden), 10.v.2016, 1 ♀(dark morph—sm) on *Prunus persica*, GD; Wied Ħesri, 15.v.2016, 2 ♀♀(pale morph—sm) on *Hypericum sinuatum*, GD; Wied Qirda, 30.v.2016, 1 ♀ on *Hyparrhenia hirta*, GD; Msida, University of Malta grounds, 11.x.2016, 1 ♀(dark morph—sm) on *Quercus robur*, GD; Dingli Cliffs, 27.i.2017, 1 ♀(dark morph—sm) on *Asphodelus ramosus*, GD; Fiddien, 24.iv.2017, 2 ♀♀(dark morph—sm) on *Medicago* sp., GD; Popeye Village, 29.vii.2017, 1 ♀(pale morph—sm) on *Tamarix africana*, GD; Siġġiewi, (road l/o Wied Ħesri), 07.viii.2017, 3 ♀♀(pale morph—sm) on *Ipomoea carnososa*, GD; Siġġiewi (private farm), 13.xi.2017, 1 ♀(pale form—sm) on *Eriobotrya japonica*, GD; Xemxija, 03.iii.2018, 7 ♀♀(dark morph—sm) on *Anacamptis urvilleana*, GD; Qormi (private farm), 7.vii.2018, 1 ♀(pale morph—sm), 3 ♀♀(dark morph—sm) on *Foeniculum vulgare*, GD; Pembroke, 05.xi.2018, 1 ♀(dark morph—sm) on *Reichardia picroides*, GD. GOZO: Ramla Bay, 02.vii.2018, 2 ♀♀(pale form—sm) on *Pancreatium maritimum*, GD.

Body length: ♀: 1060–1360 µm; ♂: no records.

Wing type: ♀: macropterous; ♂: no records.

Both the pale and the dark morphs of this species were recorded in the Maltese Islands, with the pale morphs being more common in the warmer months. *Thrips tabaci* is a polyphagous species, being recorded on numerous unrelated plants, particularly Asteraceae [24], but also on a wide variety of crops, cultivars and indigenous

plants [27,34–36,54,55,57,58,63,70]. In the Maltese Islands, this species has been originally reported on *Allium* (Amaryllidaceae) [7], though in the current study, it has been found on a number of indigenous and cultivated herbaceous annuals and trees including species of Brassicaceae, Rosaceae, Tropaeolaceae, Poaceae, Asteraceae, Apiaceae, Orchidaceae, Amaryllidaceae, Convolvulaceae and Fabaceae amongst others. *T. tabaci* tends to perform thelytokous reproduction, therefore most individuals are female [71] while males are always very rare. This accounts for the fact that in the current study, no male specimens were found despite the large number of specimens collected. *T. tabaci* is a widely distributed pest, causing both direct damage to agricultural crops as well as acting as a vector of orthospoviruses [14,26] but it is also recorded as an important pollinator [56]. *T. tabaci* is probably of Eastern Mediterranean origin [14] but is now cosmopolitan [14,27] and present wherever onion and garlic are grown [27].

4. Discussion

The insect fauna of the Maltese Islands has been well studied, with works covering practically all orders [72–74]. The order Thysanoptera is however an exception, owing to the fact that insects belonging to this order are small, difficult to identify and few people are interested to study them. As stated in the results section, the current study has recorded 39 species of Terebrantia from the Maltese Islands. This species richness compares well to other Mediterranean islands such as Sicily with 58 [13], Sardinia with 29 [13] and the Canary Islands in the Atlantic ocean with 75 species [75], despite the fact that all of these islands are considerably larger than the Maltese Islands and feature a larger variety of habitats.

Thirty eight of the identified species are phytophagous, with the four species of the genus *Aeolothrips*, as well as *Frankliniella occidentalis* and *Thrips tabaci* being described as facultative predators. The only terebrantian recorded in the Maltese Islands which is an obligate predator is *Franklinothrips megalops*.

An analysis of the zoogeographical composition of the thrips fauna based on distributional types (chorotypes) as defined by Vigna Taglianti et al. [76] (Table 2, Table A1 in Appendix A) reveals that the majority of the species (26 species–66.7%) come from Europe and the Mediterranean Region, with 12 (30.7%) of these species extending their geographical distribution towards the Middle East and the Asian continent and four species (10.2%) reaching the African continent.

Table 2. Proportion of chorotypes for the Terebrantia of the Maltese Islands.

Chorotype	Number of Species	%
Subcosmopolitan	7	17.9
Cosmopolitan	5	12.8
Palearctic	5	12.8
Turano Mediterranean	4	10.3
Asiatic European	3	7.7
Afrotropico Mediterranean	3	7.7
Mediterranean	3	7.7
Turano European	2	5.1
Afrotropico-Indo-Mediterranean	1	2.6
European	1	2.6
Europeo Mediterranean	1	2.6
Holarctic	1	2.6
Sibero European	1	2.6
Turano-Europeo-Mediterranean	1	2.6
West Palearctic	1	2.6
Total:	39	100

These findings reflect the geographical locations of the Maltese Islands, which are located in the center of the Mediterranean Sea, around 250 km north of the Libyan coast and 90 km south of Sicily. Previous local studies on insects with the approximate size and mobility of thrips [77] indicate that most of the local fauna is of European origin whereas African representation is very limited. This also seems to be the case for the terebrantian thrips. The limited amount of habitats found on the Maltese Islands may also have contributed to the fact that no endemic species of Terebrantia were recorded in this study.

Twelve (30.7%) of the species occurring in the Maltese Islands have a cosmopolitan or subcosmopolitan distribution. All of these species, with the exception of *Limothrips angulicornis*, are indicated as agriculturally important [26,32] and are capable of potentially damaging crops and cultivated plants by either feeding directly on plant tissues or by transmitting disease agents, including orthospoviruses. It is highly likely that there is a link between the chorotypes of these species and the fact that these species are widely indicated as polyphagous (since larval forms were recorded on a large variety of unrelated plants). Being able to use a wider range of plant species as hosts increases the chances of settling into new environments.

Seven (17.9%) of the species with a cosmopolitan or subcosmopolitan distribution were found to be of alien origin. These species include *Heliothrips haemorrhoidalis*, *Hercinothrips femoralis*, *Echinothrips americanus*, *Frankliniella occidentalis*, *F. schultzei*, *Thrips australis* and *T. simplex*. The geographical distribution of these seven species either extends from distant world regions into the Mediterranean, or else is found in different unconnected world regions since they were imported from their countries of origin to other areas together with crops or cultivated plants. During the current study, the above-listed species, with the exception of *Thrips australis*, were collected from cultivated fields, greenhouses or gardens, on crops or cultivated plants and were absent in indigenous habitats situated close to the locations where these species were collected from. It is interesting to note that these species were generally not collected in numbers large enough to suggest that these can act as pests in the Maltese Islands, though some of the species such as *Echinothrips americanus* were observed to have the potential of quickly establishing sizeable populations that can quickly decimate cultivated plants given the right environmental conditions. Species such as *Thrips australis* may have occurred in the Maltese Islands for quite some time. *T. australis* is closely related to *Eucalyptus* trees and may therefore have been present in the Maltese Islands as far back as the 1950s, when fast growing trees, including *Eucalyptus gomphocephala*, were imported from different parts of the world in an attempt to increase the local tree populations in urban environments.

In the present study, a number of specimens were also collected which could only be identified to genus level. These specimens belonged to the genera *Chirothrips* (1 species), *Odontothrips* (2 species), *Tenothrips* (1 species) and *Thrips* (2 species) and additional material will be required to properly identify them in future.

In due course, the thrips fauna of the Maltese Islands may change due to a number of factors, namely building over natural habitats due to urbanization requirements and the extensive use of plant protection products in agricultural fields. Another important contributing factor is climate change, with the weather of the Maltese Islands becoming hotter and drier. This can affect the distribution of certain indigenous plants that serve as hosts or even as food or shelter plants and consequently impact the thrips diversity. Importation of cultivated plants will continue to introduce additional alien species to the Maltese Islands. Species such as *Scirtothrips aurantii*, *S. citri* and *S. dorsalis* are recorded and already established in Sicily and are considered as pests on citrus trees [48]. Without proper quarantine monitoring of imported plant material and the devising of eradication policies, these species can definitely become established in the Maltese Islands in the very near future.

Author Contributions: Conceptualization, D.M. and G.D.; methodology, D.M. and G.D.; validation, A.C., A.G. and D.M.; formal analysis, G.D.; investigation, G.D.; resources, G.D.; data curation, G.D.; writing—original draft preparation, G.D.; writing—review and editing, D.M., A.C. and A.G.; visualization, D.M.; supervision, D.M., A.C. and A.G.; project administration, D.M. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: The authors would like to acknowledge Laurence A. Mound from CSIRO for the identification of *A. gloriosus* and *R. niveipennis* and G. (Bert) Vierbergen from the Netherlands Institute for Vectors, Invasive plants and Plant health, for providing information about *Tenothrips discolor*. Thanks also go to Paul Brown from the BMNH who kindly lent us specimens of *Chirothrips hamatus* and *Thrips simplex*, as well as to Edwin Lanfranco for the help in identifying some of the plants on which the thrips specimens were collected.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A

Table A1. Chorotypes for the Thysanoptera of the Maltese Islands (based on data from previously published material).

Species	Chorotype	Notes
<i>Aeolothrips gloriosus</i>	West Palaearctic	
<i>Aeolothrips intermedius</i>	Asiatic European	
<i>Aeolothrips melisi</i>	Mediterranean	
<i>Aeolothrips tenuicornis</i>	Turano Mediterranean	
<i>Franklinothrips megalops</i>	Afrotropico Mediterranean	
<i>Rhipidothrips brunneus</i>	Palaeartic	also Australian region
<i>Rhipidothrips gratiosus</i>	Palaeartic	
<i>Rhipidothrips niveipennis</i>	European	
<i>Rhipidothrips unicolor</i>	Turano Mediterranean	
<i>Melanthrips ficalbii</i>	Palaeartic	
<i>Melanthrips fuscus</i>	Palaeartic	
<i>Melanthrips knechteli</i>	Turano European	
<i>Melanthrips lybicus</i>	Mediterranean	
<i>Holarthrothrips tenuicornis</i>	Mediterranean	
<i>Dendrothrips saltator</i>	Asiatic European	also Oriental region
<i>Heliothrips haemorrhoidalis</i>	Cosmopolitan	
<i>Hercinothrips femoralis</i>	Cosmopolitan	
<i>Anaphothrips sudanensis</i>	Subcosmopolitan	
<i>Aptinothrips rufus</i>	Cosmopolitan	
<i>Asphodelothrips croceicollis</i>	Turano Europeo Mediterranean	
<i>Bregmatothrips dimorphus</i>	Afrotropico Mediterranean	also East Palaearctic
<i>Ceratothrips ericae</i>	Asiatic European	also Australian region
<i>Chirothrips hamatus</i>	Sibero European	Also Great Britain. Does not extend beyond Turkey
<i>Chirothrips manicatus</i>	Subcosmopolitan	
<i>Chirothrips meridionalis</i>	Afrotropico Indo Mediterranean	
<i>Echinothrips americanus</i>	Subcosmopolitan	
<i>Franklinella occidentalis</i>	Cosmopolitan	
<i>Frankliniella schultzei</i>	Subcosmopolitan	
<i>Limothrips angulicornis</i>	Subcosmopolitan	
<i>Limothrips cerealium</i>	Subcosmopolitan	
<i>Odontothrips meliloti</i>	Turano European	
<i>Oxythrips ajugae</i>	Holarctic	
<i>Pezothrips kellyanus</i>	Turano Mediterranean	also the Netherlands and the Australian region
<i>Prosopothrips nigriceps</i>	Turano Mediterranean	
<i>Tenothrips discolor</i>	Turano Europeo Mediterranean	
<i>Thrips australis</i>	Afrotropico Mediterranean	also Australian region
<i>Thrips major</i>	Palaeartic	
<i>Thrips simplex</i>	Subcosmopolitan	
<i>Thrips tabaci</i>	Cosmopolitan	

References

- Mound, L.A.; Wang, Z.; Lima, É.F.B.; Marullo, R. Problems with the Concept of “Pest” among the Diversity of Pestiferous Thrips. *Insects* **2022**, *13*, 61. [[CrossRef](#)] [[PubMed](#)]
- Burckhardt, D.; Ouvrard, D.; Queiroz, D.; Percy, D. Psyllid Host-Plants (Hemiptera: Psylloidea): Resolving a Semantic Problem. *Fla. Entomol.* **2014**, *97*, 242–246. [[CrossRef](#)]
- Pedley, M.; Hughes Clarke, M.; Galea, P. *Limestone Isles in a Crystal Sea*; PEG: San Gwann, Malta, 2002; p. 109.
- Chetcuti, D.; Buhagiar, A.; Schembri, P.J.; Ventura, F. *The Climate of the Maltese Islands*; Malta University Press: Msida, Malta, 1992; p. 108.
- FAO. *Malta Water Resources Review*; FAO: Rome, Italy, 2006.
- MEPA Terrestrial Habitats. Available online: <http://www.mepa.org.mt/biodiversity-habitats-terrestrial#8> (accessed on 25 February 2015).
- Mifsud, S. Malta Wild Plants. Available online: www.maltawildplants.com (accessed on 14 July 2018).
- Saliba, L.J. *Insect Pests of Crop Plants in the Maltese Islands*; Department of Information: Msida, Malta, 1963.
- Farrugia, C. *Insect Pests on Cauliflower (Brassica oleracea var. botrytis) in Gozo (Maltese Islands, Central Mediterranean)*; University of Malta: Msida, Malta, 1997.
- Mifsud, D. Biological control in the Maltese Islands—Past initiatives and future programmes. *Bull. OEPP* **1997**, *27*, 77–84. [[CrossRef](#)]
- Mound, L.A.; Palmer, J.M. Notes on Thysanoptera from Israel. *Entomol. Mon. Mag.* **1973**, *109*, 102–106.
- Mifsud, D.; Watson, G.W. *Introduced Sap-Feeding Insect Pests of Crop Plants in the Maltese Islands*; University of Malta: Msida, Malta, 1999.
- Vierbergen, G. Fauna Europaea—All European animal species on the web: Thysanoptera Section. *Biodivers. Data J.* **2014**, *2*, e4034.
- Zur Strassen, R. *Die Terebrantien Thysanopteren Europas; Die Tierwelt Deutschlands*; Goecke & Evers: Kelttern, Germany, 2003; p. 277.
- Zur Strassen, R.; Kuslitzky, W. An annotated checklist of the thrips of Israel (Thysanoptera). *Isr. J. Entomol.* **2012**, *41*, 53–66.
- National Statistics Office. *Census of Population and Housing 2021; Preliminary Report*; National Statistics Office: Valletta, Malta, 2021; p. 100.
- Mound, L.A.; Kibby, G. *Thysanoptera*; CAB International: Wallingford, UK, 1998; p. 70.
- Palmer, J.M.; Du Heaume, G.J.; Betts, C. 1959 CIE Guides to Insects of Importance to Man. 2, Thysanoptera. Betts, C.R., Ed.; CAB International: Wallingford, UK, 1989; p. 73.
- Marullo, R. I Tisanotteri dell’ Italia meridionale. II Contributo. Le specie italiane del genere *Aeolothrips* Haliday. *Boll. Lab. Entomol. Agrar. Filippo Silvestri* **1993**, *50*, 121–140.
- Marullo, R. I Tisanotteri dell’ Italia meridionale. III Contributo. La Collezione del Museo Civico di Storia Naturale “G. Doria” di Genova. *Boll. Lab. Entomol. Agrar. Filippo Silvestri* **1996**, *51*, 57–65.
- Marullo, R. Thysanoptera of southern Italy. V Contribution. Morphological remarks and biological notes on some southern Mediterranean species. *Boll. Lab. Entomol. Agrar. Filippo Silvestri Portici* **2004**, *59*, 49–57.
- Priesner, H. *A Monograph of the Thysanoptera of the Egyptian Deserts*; Desert Institute: El Mataria, Egypt, 1960; p. 582.
- Priesner, H. *Ordnung Thysanoptera (Fransenfluger Thripse)*; Akademie-Verlag: Berlin, Germany, 1964; p. 242.
- Mound, L.A.; Morrison, G.D.; Pitkin, B.R.; Palmer, J.M. *Thysanoptera. Handbooks for the Identification of British Insects*; Royal Entomological Society of London: London, UK, 1976; Volume 1, p. 79.
- Zhang, S.M.; Mound, L.A.; Hastings, A. Thysanoptera Chinensis—Thripidae Genera from China. Available online: https://keys.lucidcentral.org/keys/v3/thysanoptera_chinensis/ (accessed on 9 May 2018).
- Moritz, G.; Brandt, S.; Sseruwagi, P.; Waiganjo, M.; Subramanian, S. Pest thrips of Eastern Africa—Identification and Information based on LucID 3.5. *Mitt. Dtsch. Ges. Allg. Angew. Entomol.* **2012**, *18*, 533–539.
- Mound, L.A.; Collins, D.W.; Hastings, A. Thysanoptera Britannica et Hibernica—Thrips of the British Isles. Available online: https://keys.lucidcentral.org/keys/v3/british_thrips/.html (accessed on 9 May 2018).
- Mound, L.A.; Hoddle, M.S.; Hastings, A. Thysanoptera Californica—Thrips of California. Available online: https://keys.lucidcentral.org/keys/v3/thrips_of_california_2019/index.html (accessed on 26 April 2019).
- Marullo, R.; De Grazia, A. Territorial distribution, classification and relationships amongst Italian Thysanoptera. *Bull. Insectology* **2013**, *66*, 127–134.
- Bhatti, J.S.; Alavi, J.; zur Strassen, R. Thysanoptera in Iran 1938–2007: An Overview. *Thrips* **2009**, *7–8*, 373.
- Canale, A.; Conti, B.; Petacchi, R.; Rizzi, I. Thysanoptera collected in an olive growing area of the northern Tuscany (Italy). *Entomol. Probl.* **2003**, *33*, 105–110.
- Moritz, G. *Thripse*; Meiling Druck: Haldensleben, Germany, 2006; p. 384.
- Kirk, W.D.J. *Thrips. Naturalist’s Handbook No 25*; Richmond Publishing Co., Ltd.: Richmond, UK, 1996; p. 70.
- Fallahzedah, M.; Elaheh, A.; Nazila, S.; Hassan, A.; Jalil, A. Faunistic survey of Thysanoptera in Fars province, Iran. *Munis Entomol. Zool.* **2011**, *6*, 251–261.
- Nickle, D. A Checklist of Commonly Intercepted Thrips (Thysanoptera) from Europe, the Mediterranean, and Africa at U.S. Ports-of Entry (1983–1999). Part 1. Key to Genera. In Proceedings of the Entomological Society of Washington, Washington, DC, USA, 2 May 2003; Volume 105, pp. 80–99.
- Tunç, İ.; Bahşi, Ş.Ü.; Göçmen, H. Thysanoptera fauna of the Aegean region, Turkey, in the spring. *Turk. J. Zool.* **2012**, *36*, 592–606. [[CrossRef](#)]

37. Mound, L.A.; Reynaud, P. *Franklinothrips*; a pantropical Thysanoptera genus of ant-mimicking obligate predators (Aeolothripidae). *Zootaxa* **2005**, *864*, 1–16. [[CrossRef](#)]
38. Alavi, J.; zur Strassen, R.; Bagherani, N. Thrips (Thysanoptera) species associated with wheat and barley in Golestan province, Iran. *J. Entomol. Soc. Iran* **2007**, *27*, 1–28.
39. Mirab-balou, M.; Chen, X.; Tong, X. *Tenothrips bhatti*, a newly recorded genus of Thripinae (Thysanoptera: Thripidae) from China. *Entomotaxonomia* **2012**, *34*, 162–166.
40. Razi, S.; Laamari, M. Thysanoptera survey on *Vicia faba* (broad bean) in the arid Biskra region of Algeria. *Agric. Biol. J. North Am.* **2013**, *4*, 268–274. [[CrossRef](#)]
41. Gertsson, C. An annotated checklist of Thysanoptera (thrips) from the Nordic countries. [Provinsförteckning över Nordens tripsar.]. *Entomol. Tidskr.* **2015**, *134*, 185–198.
42. Marullo, R. *Un Profilo della Tisanotterofauna italiana: Le Interazioni tra Specie e Piante Ospiti e gli Effetti delle Coltive sulla Diversità delle Popolazioni Naturali*; Congresso Nazionale Italiano di Entomologia: Rome, Italy, 2002; pp. 179–184.
43. Stoch, F. Fauna Italica. Available online: <http://www.faunaitalia.it/> (accessed on 20 February 2015).
44. Zur Strassen, R. Thysanoptera on islands of the northern Sporades in the Aegean (Greece) (Insecta: Thysanoptera). *Senckenberg. Biol.* **1986**, *67*, 85–129.
45. Tunc, I. Studies on the Thysanoptera of Antalya I. Aeolothripidae Uzel. *Türk. Entomoloji Derg.* **1991**, *15*, 129–141.
46. Sierka, W.; Fedor, P.; Vasiliu-Oromulu, L.; Jenser, G.; Bărbuceanu, D. The state of knowledge of thrips (Insecta: Thysanoptera) of the Carpathian Mountains. *Acta Phytopathol. Entomol. Hung.* **2008**, *43*, 355–366. [[CrossRef](#)]
47. Karadjova, O.; Krumov, V. Thysanoptera of Bulgaria. *ZooKeys* **2015**, *504*, 93–131.
48. Marullo, R. *Conoscere i Tisanotteri*; Edizioni Agricole de Il Sole 24 ore Edagricole S.r.l. via Goito 13—40126; Università degli Studi Mediterranea di Reggio Calabria: Reggio Calabria, Italy, 2003; p. 75.
49. ThripsWiki Contributors zur Strassen Distribution Lists. Available online: http://thrips.info/w/index.php?title=Zur_Strassen_distribution_lists&oldid=41628 (accessed on 6 December 2016).
50. ThripsWiki. *ThripsWiki—Providing Information on the World's Thrips in the Catalogue of Life*; ThripsWiki: Leiden, The Netherlands, 2018.
51. Nakahara, S.; O'donnell, C.A.; Mound, L.A. *Heliothrips haemorrhoidalis* and its relatives, with one new species and one new genus (Thysanoptera: Thripidae). *Zootaxa* **2015**, *4021*, 578–584. [[CrossRef](#)]
52. Mound, L.A.; Nielsen, M.; Hastings, A. Thysanoptera Aaotearoa—Thrips of New Zealand. Available online: https://keys.lucidcentral.org/keys/v3/nz_thrips/ (accessed on 3 April 2019).
53. Nickle, D.A. Commonly Intercepted Thrips at U.S. Ports-of-Entry from Africa, Europe, and the Mediterranean. IV. Miscellaneous Thripine Genera Excluding Frankliniella, Iridothrips, and Thrips (Thysanoptera: Thripidae). In Proceedings of the Entomological Society of Washington, Washington, DC, USA, 1 January 2009; Volume 111, pp. 215–238.
54. Raspudić, E.; Ivezić, M.; Brmež, M.; Trdan, S. Distribution of Thysanoptera species and their host plants in Croatia. *Acta Agric. Slov.* **2009**, *93*, 275–283. [[CrossRef](#)]
55. Trdan, S. Thrips in Slovenia: Thrips and tospoviruses. In Proceedings of the 7th International Symposium on Thysanoptera, Reggio Calabria, Italy, 2–7 July 2001; pp. 351–356.
56. Garcia-Fayos, P.; Goldarazena, A. Role of Thrips in Pollination of *Arctostaphylos uva-ursi*. *Int. J. Plant Sci.* **2008**, *169*, 776–781. [[CrossRef](#)]
57. Jenser, G. Data to the Thysanoptera fauna of Tunisia. *Folia Entomol. Hung.* **1982**, *43*, 55–57.
58. Trdan, S.; Andjus, L.; Raspudić, E.; Kač, M. Distribution of *Aeolothrips intermedius* Bagnall (Thysanoptera: Aeolothripidae) and its potential prey Thysanoptera species on different cultivated host plants. *J. Pest Sci.* **2005**, *78*, 217–226. [[CrossRef](#)]
59. Badieritakis, E.G.; Thanopoulos, R.C.; Fantinou, A.A.; Emmanouel, N.G. Emmanouel qualitative and quantitative study of thrips (Thysanoptera) on alfalfa and records of thrips species on cultivated and wild Medicago species of Greece. *Biologia* **2015**, *70*, 504–515. [[CrossRef](#)]
60. Mound, L.A.; Teulon, D.A.J. *Thysanoptera as Phytophagous Opportunists*; Thrips Biology and Management; Springer: Boston, MA, USA, 1995; pp. 3–19.
61. Tyagi, K.; Kumar, V. Thrips of Economic importance in India—An Identification guide. *Zool. Surv. India* **2020**, *2020*, 96.
62. Goldarazena, A. Orden Thysanoptera. *Rev. SEA* **2015**, *52*, 1–20.
63. Jenser, G.; Tzanakakis, M.E. Records of Thysanoptera from Northern Greece. *Entomol. Hell.* **2017**, *3*, 59. [[CrossRef](#)]
64. Kucharczyk, H.; Zawirska, I. The occurrence of Thysanoptera in Poland; Thrips and Tospoviruses. In Proceedings of the 7th International Symposium on Thysanoptera, Reggio Calabria, Italy, 2–7 July 2001; pp. 341–344.
65. Vassiliou, V.A. Ecology and Behavior of *Pezothrips kellyanus* (Thysanoptera: Thripidae) on Citrus. *J. Econ. Entomol.* **2010**, *103*, 47–53. [[CrossRef](#)]
66. Conti, F.; Tumminelli, R.; Amico, C.; Fiscaro, R.; Frittitta, C.; Perrotta, G.; Marullo, R. Monitoring *Pezothrips kellyanus* on citrus in eastern Sicily. Thrips and Tospoviruses. In Proceedings of the 7th International Symposium on Thysanoptera, Reggio Calabria, Italy, 2–7 July 2001; pp. 207–210.
67. Navarro Campos, C. *Pezothrips Kellyanus* (Thysanoptera: Thripidae), Nueva Plaga en Cítricos; Comportamiento de sus Poblaciones, Muestreo y Enemigos Naturales. Unpublished. Ph.D. Thesis, Universitat Politècnica de València, Valencia, Spain, 2013; p. 166.

68. Varikou, K.; Tsitsipis, I.; Alexandrakis, V.; Hoddle, M. Effect of Temperature on the Development and Longevity of *Pezothrips kellyanus* (Thysanoptera: Thripidae). *Ann. Entomol. Soc. Am.* **2009**, *102*, 835–841. [[CrossRef](#)]
69. Nickle, D.A. Commonly Intercepted Thrips at U.S. Ports-of-Entry from Africa, Europe, and the Mediterranean. III. The Genus Thrips Linnaeus, 1758 (Thysanoptera: Thripidae). *Proc. Entomol. Soc. Wash.* **2008**, *110*, 165–185. [[CrossRef](#)]
70. Kucharczyk, H.; Kucharczyk, M. Characteristic and diagnostic features of the most frequently occurring species of the Thripidae family (Insecta, Thysanoptera) in crown canopies of Central European forests. *Leśne Pr. Badaw.* **2013**, *74*, 5–11.
71. Khan, F.; Roy, M.C.; Kim, Y. Thelytokous Reproduction of Onion Thrips, *Thrips tabaci* Lindeman 1889, Infesting Welsh Onion and Genetic Variation among Their Subpopulations. *Insects* **2022**, *13*, 78. [[CrossRef](#)]
72. Mifsud, D. Present knowledge of the Entomofauna of the Maltese Islands. *Entomol. Basiliensia* **2000**, *22*, 75–86.
73. Schembri, P.J. Current state of knowledge of the Maltese non-marine fauna. In Malta Environment and Planning Authority Annual report and accounts. *Malta Environ. Plan. Auth.* **2003**, *2003*, 33–65.
74. Cassar, T.; Mifsud, D. Insects of the Maltese Islands. *Entomol. Soc. Malta* **2023**, *12*, 711.
75. Berzosa, J. Thysanoptera checklist (Insecta, Thysanoptera) of the Canary Islands. Geographical distribution, host-plants and bibliography. *Boll. Real Soc. Esp. Hist. Nat. Sec. Biol.* **2000**, *96*, 93–112.
76. Vigna Taglianti, A.; Audisio, P.A.; Biondi, M.; Bologna, M.A.; Carpaneto, G.M.; De Biase, A.; Fattorini, S.; Piattella, E.; Sindaco, R.; Venchi, A.; et al. A proposal for a chorotype classification of the Near East fauna, in the framework of the Western Palearctic region. *Biogeogr. J. Integr. Biogeogr.* **1999**, *20*, 1. [[CrossRef](#)]
77. Carapezza, A.; Mifsud, D. New records of true bugs (Hemiptera, Heteroptera) from the Maltese Islands 2015. *Bull. Entomol. Soc. Malta* **2015**, *7*, 27–50.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.