diversity

MDPI

Article

A New Species of *Enicospilus* Stephens, 1835 (Ichneumonidae, Ophioninae), from Southern Mexico, Parasitic on *Zanola verago* Cramer, 1777 (Lepidoptera, Apatelodidae), Feeding on *Piper neesianum* C. DC. (Piperaceae)

Diego Fernando Campos-Moreno [1,*], Edgard Palacio [2], Luis Alberto Lara-Pérez [3], James B. Whitfield [4], Carmen Pozo [5] and Lee A. Dyer [6,*]

[1] Programa de Maestría en Uso, Manejo y Conservación de Bosques, Facultad de Medio Ambiente y Recursos Naturales, Universidad Francisco José de Caldas, Bogotá 110321, Colombia
[2] Instituto Colombiano Agropecuario (ICA), Barranquilla 110221, Colombia
[3] Tecnológico Nacional de México, Instituto Tecnológico de la Zona Maya, Chetumal 77965, Mexico
[4] Department of Entomology, University of Illinois, Urbana-Champaign, Urbana, IL 61801, USA
[5] El Colegio de la Frontera Sur. Unidad Chetumal. Av. Centenario, km 5.5 s/n. Col. Pacto Obrero. Chetumal, Quintana Roo. CP 77014, Chetumal 77014, Mexico
[6] Biology Department, University of Nevada-Reno, Reno, NV 89557, USA
* Correspondence: dfcamposm@unal.edu.co (D.F.C.-M.); ldyer@unr.edu (L.A.D.)

Abstract

Plant–herbivore–parasitoid systems are poorly studied in the tropics. *Enicospilus carmenae* Campos and Palacio sp. nov. are described, originating from southern Mexico in the Yucatan Peninsula and establishing a new tri-trophic interaction. This species is a koinobiont larval endoparasitoid of the American silkworm moth caterpillar *Zanola verago* (Cramer) (Lepidoptera: Apatelodidae) feeding on the shrub *Piper neesianum* C.DC. (Piperaceae) in a semi-evergreen forest. The host plant *P. neesianum* had no herbivore records to date, and a single collection event yielded the rearing of a new species of *Enicospilus* (Ichneumonidae, Ophioninae). Morphological, molecular (COI), biological, ecological, and geographical data are integrated to delineate the new species.

Keywords: species description; natural history; tri-trophic interaction; Yucatán Peninsula

check for updates

Academic Editors: Luc Legal and Michael Wink

Received: 14 July 2024
Revised: 4 March 2025
Accepted: 21 March 2025
Published: 3 July 2025

Citation: Campos-Moreno, D.F.; Palacio, E.; Lara-Pérez, L.A.; Whitfield, J.B.; Pozo, C.; Dyer, L.A. A New Species of *Enicospilus* Stephens, 1835 (Ichneumonidae, Ophioninae), from Southern Mexico, Parasitic on *Zanola verago* Cramer, 1777 (Lepidoptera, Apatelodidae), Feeding on *Piper neesianum* C. DC. (Piperaceae). *Diversity* **2025**, *17*, 466. https://doi.org/10.3390/d17070466

Copyright: © 2025 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).

1. Introduction

The discovery of new species and their interactions with other organisms is critical to developing a deeper understanding of biodiversity [1,2]. Interactions between plants, herbivores, and natural enemies involve over half of all described species [3], and documenting the natural history of these interactions contributes to understanding evolutionary and ecological outcomes, such as host specificity [4,5]. Studies of tri-trophic interactions between plants, herbivorous insects, and their natural enemies have contributed substantially to basic research in ecology [6–8], as well as applied sciences, including biological control [9] and conservation biology [10,11]. These tri-trophic studies are important for revealing ecological and evolutionary patterns that cannot be understood by separately studying each trophic level [12].

The genus *Piper* (Piperaceae), with a pantropical distribution, is one of the most diverse taxa within the angiosperms, with more than 1300 species in the Neotropics [13]. *Piper neesianum* C. DC. is native to the "ramonales" forests of the Yucatán Peninsula, which

are semi-evergreen forests on calcareous soils dominated by *Brosimum alicastrum* Sw. [14]. Currently, the known distribution includes southern Mexico, Guatemala, and Belize [15]. There are no published herbivore associations for this species [16,17].

The American silkworm moth genus *Zanola* Cramer, 1777 (Bombycoidea, Apatelodidae), includes seven valid species [18]. *Zanola verago* is distributed throughout the Neotropical Region in relatively well-preserved forests [19–21]. The larva is grey-black and orange, with numerous secondary setae and distinctive subdorsal red spots. The head and tail have tufts of long black hairs with the tips widened into flat triangular scales [20]. *Zanola verago* is a polyphagous species that feeds on at least 12 plant species within 11 families across its distribution [17,20–22].

Ophioninae is an ichneumonid subfamily comprising slender wasps with orange to brown coloration. All are koinobiont solitary endoparasitoids of crepuscular and nocturnal Lepidoptera [23]. *Enicospilus* is the most species-rich genus within the subfamily, with more than 150 described species in the New World [24–28] and 61 species in Mexico [29]. The Mesoamerican species of *Enicospilus* are parasitoids of large moth caterpillars including Saturnidae, Lasiocampidae, and Erebidae, and, as far as is known, there is only one record of *Enicospilus* developing on Apatelodidae [17].

As a result of intensive surveys in semi-evergreen forests of the Yucatán Peninsula [30], a new tri-trophic interaction involving *Piper neesianum* (Piperaceae), *Zanola verago* (Lepidoptera, Apatelodidae), and a larval koinobiont endoparasitoid belonging to a new species of *Enicospilus* (Ichneumonidae, Ophioninae) is reported.

2. Materials and Methods

The parasitoid was reared from a manually collected caterpillar feeding on *P. neesianum* (Figure 1A) in a semi-evergreen forest in the southeast of Yucatán Peninsula (19.28° N–88.62° W). This locality is part of INIFAP, a forest reserve established in 1971. The caterpillar, identified as *Zanola verago* (Apatelodidae) (Figure 1B), was reared under ambient conditions (i.e., an outdoor facility, as is common for large rearing projects [4,17]) at the ECOSUR laboratory in a plastic cup covered with mesh and fed with leaves of *P. neesianum* until adult parasitoids emerged. The host remains were preserved in ethanol. A Nikon D850 digital camera attached to a Leica DMLB stereomicroscope was used to take photographs of the caterpillar.



Figure 1. Habit of the host plant and host larva. (**A**) *Piper neesianum* (photo by E. Tepe); (**B**) *Zanola verago* (photo by D. Campos).

Morphological terminology, measurements, and indices for the species description follow [23,24]. Layer photos of the new species were taken at the Instituto Colombiano Agropecuario (ICA) using a LEICA MC170 HD camera attached to a LEICA S8APO stereomicroscope. The female holotype was compared with layer photos of *E. gamezi* taken at NHMUK using a Canon SLR EOS 5DSR with a 65 mm macro lens mounted on a copy

stand with an automated Z-stepper; images were aligned using Helicon Focus software, version 6.6.1. The holotype is preserved at the Arthropod Collection of ECOSUR, Chetumal (ECO-CH-AR).

DNA Protocols

DNA extraction and PCR were carried out at El Colegio de la Frontera Sur (ECOSUR) Chetumal, using standardized protocols. DNA was extracted (from a single EtOH-preserved leg) according to the protocol in [31]. PCR amplification of the COI gene used primers LCO1490: 5′-GGTCAACAAATCATAAAGATATTGG-3′ and HCO2198: 5′-TAAACTTCAGGGTGACCAAAAAATCA-3′ [32]. The faster rate of evolution of the mitochondrial genes is sometimes helpful for separating closely related species, and they are frequently used for the species delimitation of parasitoids [33,34]. We sent PCR products to the High Throughput Genomics Center in Seattle, USA, for sequencing. Newly generated sequences were deposited in GenBank database under accession number MZ960307. Additionally, sequences of species closely related (BLAST best hit) to new *Enicospilus* species were obtained from the Bold system [35,36]. Sequences were aligned in MEGA 10 [37,38]. Interspecific genetic distances were calculated using the Kimura-2-parameter model (K2P) [37] to produce the neighbour-joining (NJ) tree to display divergences [39], using *Thyreodon maculipennis* and *Cryptophion inaequalis* as outgroups.

3. Results

Enicospilus carmenae Campos & Palacio sp. nov.

ZooBank 965D83BA-07AF-4E22-AB69-157EE838792E (http://zoobank.org/965D83BA-07AF-4E22-AB69-157EE838792E, accessed on 21 April 2025).

3.1. Description

Holotype:

Zoobank: 3C55513F-C906-416F-927F-52CE8C57BA51;

Locality: 19.28° N; 88.62° W semi-evergreen forest in the southeast of Yucatán Peninsula; Mexico; female; adult; Catalog Code: AR730

Female: Length of body 21.0 mm. Fore wing length 16 mm. AI = 1.285; CI = 0.421; ICI = 0.633; SDI = 1.254.

Head: Clypeus in profile weakly convex, in frontal view 1.6× as broad as long, margin subtruncate (Figure 2A). Lower face as broad as long, polished, punctate. Head in dorsal view with genae rounded behind eyes; posterior ocellus very close to eye; FI = 0.651 (65%); occipital carina mediodorsally distinct, not joining hypostomal carina. Antenna with 61 flagellomeres, central flagellomeres at least 2.0× as long as broad.

Mesosoma: Mesoscutum polished, with fine shallow punctures, in profile evenly rounded, with anterior margin slightly turned forwards, notauli vestigial. Mesopleuron polished, the upper and lower parts punctostriate; epicnemial carina abruptly curved towards anterior margin of pleuron. Scutellum, in profile weakly convex, laterally carinate entire length; in dorsal view 1.11× as long as anteriorly wide, virtually smooth. Metapleuron strongly convex, with fine diagonal striation; submetapleural carina broadened towards base; posterior transverse carina of mesosternum complete. Propodeum in profile declivous; anterior transverse carina complete, strong (Figure 3A); posterior transverse carina absent; anterior area coriaceous to striate; spiracular area short, slightly rugulose; posterior area strongly rugose; metapleural carina complete, not joined to spiracular margin by a short carina. Fore tibia slightly flattened, with scattered spines on outer surface. Mid-leg with longer tibial spur 1.40× the length of the shorter. Hind leg with tarsomere IV

3.3× as long as broad; claws evenly down-curved on apical quarter (Figure 3C), with long stout pecten.




Figure 2. Holotype of *Enicospilus carmenae* (**A**,**C**,**E**) (photos by E. Palacio); holotype of *Enicospilus gamezi* (**B**,**D**,**F**) (photos by G. Broad): (**A**,**B**) head in frontal view; (**C**,**D**) mandibles; (**E**,**F**) fore wing; arrow points to the basal section of vein *Rs* + 2*r*.




Figure 3. *Enicospilus carmenae* (**A**–**C**) (photos by E. Palacio); *E. gamezi* (**D**) (photo by G. Broad from Holotype in NHMUK): (**A**) propodeum in dorsal view; (**B**) metasomal segment II in lateral view; (**C**) metatarsal claw of *E. carmenae;* (**D**) metatarsal claw of *E. gamezi*.

Wings: Fore wing with vein *Rs* + 2*r* strongly sinuate, not basally angulate below (Figure 2E) fenestra without distinct sclerites, basally not extending to level of base of pterostigma, its apex clearly separated from base of vein *Rs*; *cu-a* proximal to base of *RS* and *M* by about 0.27× its own length; marginal cell proximally evenly hirsute; first subdiscal cell with anterior 0.4 sparsely hirsute; second discal cell 3.29× as long as wide. Hind wing with eight hamuli on R1, first abscissa of *Rs* straight, second abscissa slightly curved.

Metasoma: Slender, tergite 2 in profile 4.4× as long as posteriorly deep, laterotegite not pendant, thyridia oval and separated from anterior margin of tergite by about 3.9× its own length (Figure 3B). Ovipositor slender, its sheath narrow.

Colour: Pale yellowish-brown, with basal antennomeres and metasomal segment V darker (Figure 4A); interocellar area with black markings bordering the ocelli. Pterostigma golden, wings hyaline.




Figure 4. Habitus of *Enicospilus:* (**A**) holotype *E. carmenae* (photo by E. Palacio); (**B**) holotype *Enicospilus gamezi* (photo by G. Broad from Holotype in NHMUK).

Male: Unknown.

3.2. Diagnosis

E. carmenae can be distinguished from other species of *Enicospilus* by the combination of the following characteristics: fore wing with vein *Rs* + 2r tubular, sinuous, not basally angulate below; fenestra without distinctive sclerites (Figure 3E); apex of fenestra detached from *Rs* vein base; second discal cell 3.29× longer than high; anterior transverse carina of propodeum complete, strong (Figure 4A).

Remarks: *E. carmenae* sp. nov. links to *E. gamezi* in couplet 20 of Gauld's [23] key to *Enicospilus* species. As in *E. gamezi* (Figure 4B), the new species has central flagellar segments 2 or more times as long as broad and a propodeum without posterior transverse carina, but differs in the more slender mandibles with a subtruncated upper tooth (Figure 3C) (thicker mandibles, with the upper tooth pointed in *E. gamezi*—Figure 2D), the clypeus 1.6× as broad as long (Figure 2A) (1.4–1.5× in *E. gamezi*—Figure 3B), the vein Rs + 2r of the fore wing not basally angulated below (Figure 2E) (clearly angulated below in *E. gamezi*—Figure 3F), and the hind leg tarsal claw evenly down-curved on its apical quarter (Figure 3C) (abruptly down-curved in *E. gamezi*—Figure 3D).

3.3. Etymology

It is named after Carmen Pozo, who has dedicated most of her life to the study of the taxonomy, biogeography, and ecology of Lepidoptera and the conservation of biodiversity in southeastern Mexico.

3.4. Distribution

Southern Mexico, Yucatan Peninsula.

3.5. Ecology

Habitat: Known in semi-evergreen tropical forests.

3.6. Biology

All species of Ophioninae are known as koinobiont endoparasitoids, and this is also the case for *E. carmenae*, which was parasitizing the *Zanola verago* larva while it was still actively feeding on *Piper* leaves. The exact moment at which the host caterpillar was parasitized is unknown, but it became a pupa eight days after being collected and the female parasitoid wasp emerged as an adult 18 days later.

3.7. DNA Barcode

A single 584 bp sequence was recovered from the holotype (AR730, Bold no. DFCM CH752B07, NCBI MZ960307). The closest related species was *E. gamezi,* with a variation p-distance of 1.5% (Figure 5), with differences in nucleotide positions 238G, 268A, 340C, 373G, 442C, 460C, 562C, and 629A. These differences are sufficient to separate many morphologically and ecologically distinctive species [40–42]. Interspecific variations with other related species (*E. cameronii*, *E. bozai*, *E. chiriquensis*) show p-distances of 6–8%.



Figure 5. Phenogram based on K2P distances for COI for *E. carmenae* sp. nov. (NCBI MZ960307) and related sequences of the genus *Enicospilus* downloaded from the BOLD system and International Nucleotide Sequences Database. The black circle indicates holotype.

4. Discussion

During the 40 years of intensive rearing of caterpillar parasitoids carried out by Janzen and Hallwachs and their team in Costa Rica, only one species of *Enicospilus*, *E. gamezi*, has been recorded as a parasitoid of apatelodid larvae [17]. It is striking that *E. gamezi* is the most closely related described species to *E. carmenae* according to the current study, suggesting

a consistent pattern of interactions between *Enicospilus* and Apatelodidae hosts. Similar caterpillar-rearing projects, carried out for over 35 years in Costa Rica [42] and 25 years in Ecuador [43], have not reported apatelodid caterpillars parasitized by *Enicospilus*, so these interactions are likely to be rare. Rare ecological interactions and rare species are key components of the functional and taxonomic diversity of tropical ecosystems, often contributing disproportionately to ecosystem stability and resilience [4,44]. These rare interactions may reflect specialized evolutionary histories or ecological constraints that are an important part of understanding the full complexity of tropical food webs [4]. We acknowledge that this description is based on a single specimen, and while the evidence supports its designation as a new species, further research on this genus is necessary, especially for exploring genetic variation among closely related species.

Taking into account that Lasiocampidae, Erebidae, and Saturniidae are the Lepidopteran families most commonly reported as hosts of Mesoamerican *Enicospilus* [16,45], the recording of Apatelodidae is not entirely unexpected. However, it is interesting that the host plant, *P. neesianum*, had no herbivore records to date [16,17] and that a single collection event yielded the rearing of a new species of *Enicospilus* from a host belonging to a family of Lepidoptera that had only been reported once as a host of Ophioninae [17]. There is much to be done in the study of food webs, especially with respect to the identification of tropical species involved in these poorly studied plant–herbivore–parasitoid systems [4,5], and much more research is required to understand patterns of specialization and speciation in Ophioninae [44,46].

5. Conclusions

In conclusion, this new plant–herbivore–parasitoid system contributes to the continued documentation of tri-trophic interaction diversity [1]. The fact that this is a singleton is relevant to questions about assessing specialization and interaction diversity in tri-trophic interaction webs that are dominated by rare species. It is necessary to continue documenting trophic interaction diversity [1] while contributing to resolving taxonomic and systematic issues for the taxa that comprise these interactions.

Author Contributions: Conceptualization, D.F.C.-M., E.P. and C.P.; Methodology, D.F.C.-M., E.P. and L.A.L.-P.; Formal analysis, D.F.C.-M. and E.P.; Investigation, D.F.C.-M.; Resources, L.A.D.; Data curation, D.F.C.-M.; Writing—original draft, D.F.C.-M. and E.P.; Writing—review & editing, D.F.C.-M., E.P., L.A.L.-P., J.B.W., C.P. and L.A.D. All authors have read and agreed to the published version of the manuscript.

Funding: Graduate scholarship awarded by CONAHCYT (CVU_824833) and the U.S. National Science Foundation (NSF DEB-1442103/DEB1542103/DEB1442075/EN 2133818).

Institutional Review Board Statement: Not applicable.

Data Availability Statement: Data are contained within the article.

Acknowledgments: We are grateful to INIFAP for access to the research area and for the technical assistance from Raziel Lucio (fieldwork), Noemí Salas (specimen processing), Humberto Bahena-Basave (photography), Alma García (Nodo BARCODE), and Blanca Prado (BOLD System Project) at ECOSUR, Chetumal, MX. We also thank Eric Tepe for the identification of *Piper neesianum* C. DC. (Department of Biological Sciences University of Cincinnati, Cincinnati, UE) and Gavin Broad (NHM, London, UK) for taking the layer photos of *E. gamezi*.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Dyer, L.A.; Smilanich, A.M.; Gompert, Z.; Forister, M.L. Insect conservation, technological traps, and the fading arts of natural history and field ecology. *Curr. Opin. Insect Sci.* **2024**, *66*, 101261. [CrossRef] [PubMed]
2. Wilson, E.O. Biodiversity research requires more boots on the ground. *Nat. Ecol. Evol.* **2017**, *1*, 1590–1591. [CrossRef]
3. Price, P.W. Resource-driven terrestrial interaction webs. *Ecol. Res.* **2002**, *17*, 241–247. [CrossRef]
4. Dyer, L.A.; Singer, M.S.; Lill, J.T.; Stireman, J.O.; Gentry, G.L.; Marquis, R.J.; Ricklefs, R.E.; Greeney, H.F.; Wagner, D.L.; Morais, H.C.; et al. Host specificity of Lepidoptera in tropical and temperate forests. *Nature* **2007**, *448*, 696–699. [CrossRef] [PubMed]
5. Forister, M.L.; Dyer, L.A.; Singer, M.S.; Stireman, J.O.; Lill, J.T. Revisiting the evolution of ecological specialization, with emphasis on insect–plant interactions. *Ecology* **2012**, *93*, 981–991. [CrossRef]
6. Singer, M.S.; Stireman, J.O. The tri-trophic niche concept and adaptive radiation of phytophagous insects. *Ecol. Lett.* **2005**, *8*, 1247–1255. [CrossRef]
7. Dyer, L.A.; Forister, M.L. Challenges and advances in the study of latitudinal gradients in multitrophic interactions, with a focus on consumer specialization. *Curr. Opin. Insect Sci.* **2019**, *32*, 68–76. [CrossRef]
8. Toro-Delgado, E.; Hernández-Roldán, J.; Dincă, V.; Vicente, J.C.; Shaw, M.R.; Quicke, D.L.; Vodă, R.; Albrecht, M.; Fernández-Triana, J.; Vidiella, B.; et al. Butterfly–parasitoid–hostplant interactions in Western Palaearctic Hesperiidae: A DNA barcoding reference library. *Zool. J. Linn. Soc.* **2022**, *196*, 757–774. [CrossRef]
9. Zhang, Y.M.; Ridenbaugh, R.D.; Sharanowski, B.J. Integrative taxonomy improves understanding of native beneficial fauna: Revision of the Nearctic *Peristenus* pallipes complex (Hymenoptera: Braconidae) and implications for release of exotic biocontrol agents. *Syst. Entomol.* **2017**, *42*, 596–608. [CrossRef]
10. Poelman, E.H.; van Loon, J.J.; Dicke, M. Consequences of variation in plant defense for biodiversity at higher trophic levels. *Trends Plant Sci.* **2008**, *13*, 534–541. [CrossRef]
11. Thierry, M.; Hrček, J.; Lewis, O.T. Mechanisms structuring host–parasitoid networks in a global warming context: A review. *Ecol. Entomol.* **2019**, *44*, 581–592. [CrossRef]
12. Evans, D.M.; Kitson, J.J.; Lunt, D.H.; Straw, N.A.; Pocock, M.J. Merging DNA metabarcoding and ecological network analysis to understand and build resilient terrestrial ecosystems. *Funct. Ecol.* **2016**, *30*, 1904–1916. [CrossRef]
13. Quijano-Abril, M.A.; Callejas-Posada, R.; Miranda-Esquivel, D.R. Areas of endemism and distribution patterns for Neotropical *Piper* species (Piperaceae). *J. Biogeogr.* **2006**, *33*, 1266–1278. [CrossRef]
14. Bornstein, A.J. Taxonomic studies in the Piperaceae—I: The pedicellate pipers of Mexico and Central America. *J. Arnold Arbor.* **1989**, *70*, 1–55. [CrossRef]
15. Ulloa Ulloa, C.P.; Acevedo-Rodríguez, S.G.; Beck, M.J.; Belgrano, R.; Bernal, P.E.; Berry, L.; Brako, M.; Celis, G.; Davidse, S.R.; Gradstein, O.; et al. An integrated assessment of the vascular plant species of the Americas. *Science* **2018**, *358*, 1614–1617. [CrossRef] [PubMed]
16. HOSTS—A Database of the World's Lepidopteran Host Plants. 2010. Available online: https://data.nhm.ac.uk/dataset/hosts (accessed on 12 August 2020).
17. Janzen DHHW Philosophy, Navigation and Use of a Dynamic Database ("ACG Caterpillars SRNP") for An Inventory of the Macrocaterpillar Fauna, Its Food Plants and Parasitoids, of the Area de Conservación Guanacaste (ACG), Northwestern Costa Rica. 2012. Available online: https://www.gdfcf.org/content/acg-species-pages-over-1000-and-counting (accessed on 21 April 2025).
18. Kitching, I.; Rougerie, R.; Zwick, A.; Hamilton, C.; Laurent, R.S.; Naumann, S.; Mejia, L.B.; Kawahara, A. A global checklist of the Bombycoidea (Insecta: Lepidoptera). *Biodivers. Data J.* **2018**, *6*, e22236. [CrossRef]
19. Walker, F. List of the specimens of lepidopterous insects in the collection of the British Museum. In *The Turstees*; British Museum: London, UK, 1855; Volume 5.
20. Dyar, H.G. Descriptions of the larvae of some Lepidoptera from Mexico. *Proc. Entomol. Soc. Wash.* **1912**, *14*, 54–58.
21. Ricardo-Molina, J.; Murillo-Ramos, L.; Álvarez-Pérez, P. Larvas y plantas nutricias de Lepidoptera en fragmentos de bosque seco tropical del departamento de Sucre, Colombia (Insecta: Lepidoptera). *Rev. De Lepidopterol.* **2019**, *47*, 5–24.
22. Rabelo, R.S.; Dyer, L.A.; Lepesqueur, C.; Salcido, D.M.; da Silva, T.P.; Rodrigues, H.P.A.; Trindade, T.B.; Diniz, I.R.; Nascimento, A.R.; Tepe, E.J.; et al. Tritrophic interaction diversity in gallery forests: A biologically rich and understudied component of the Brazilian cerrado. *Arthropod-Plant Interact.* **2021**, *15*, 773–785. [CrossRef]
23. Gauld, I.D. A survey of the Ophioninae (Hymenoptera: Ichneumonidae) of tropical Mesoamerica with special reference to the fauna of Costa Rica. *Bull. Br. Mus. Nat. Hist. Entomol. Ser.* **1988**, *57*, 1–309.
24. Gauld, I.D. The species of the *Enicospilus americanus* complex (Hymenoptera: Ichneumonidae) in eastern North America. *Syst. Entomol.* **1988**, *13*, 31–53. [CrossRef]
25. Fernández-Triana, J.L. The taxonomy and biogeography of Cuban Ophioninae (Hymenoptera: Ichneumonidae). *Zootaxa* **2005**, *1007*, 1–60. [CrossRef]

26. Garcia-Garcia, A. Tres especies nuevas de *Enicospilus* (Ichneumonidae: Ophioninae) de Colombia. *Rev. Colomb. Entomol.* **2011**, *37*, 145–151. [CrossRef]
27. Lima, A.R.; Jacobi, C.M.; Kumagai, A.F. A key to the Neotropical species of the *Enicospilus ramidulus* species-group (Hymenoptera: Ichneumonidae: Ophioninae), with the description of a new Brazilian species. *Zootaxa* **2012**, *3409*, 63–68. [CrossRef]
28. Yu, D.S.; van Achterberg, C.V.; Horstmann, K. Taxapad 2016, Ichneumonoidea 2015. In *Database on Flash-Drive*; The Catalogue of Life: Nepean, ON, Canada, 2016.
29. Ruíz-Cancino, E. La familia Ichneumonidae (Hymenoptera) en México. *Entomol. Mex.* **2015**, *2*, 1–13.
30. Campos-Moreno, D.; Dyer, L.; Salcido, D.; Massad, T.J.; Pérez-Lachaud, G.; Tepe, E.; Whitfield, J.; Pozo, C. Importance of interaction rewiring in determining spatial and temporal turnover of tritrophic (*Piper*-caterpillar-parasitoid) metanetworks in the Yucatán Península, México. *Biotropica* **2021**, *53*, 1071–1081. [CrossRef]
31. Ivanova, N.V.; Dewaard, J.R.; Hebert, P.D. An inexpensive, automation friendly protocol for recovering high-quality DNA. *Mol. Ecol. Notes* **2006**, *6*, 998–1002. [CrossRef]
32. Folmer, O.; Black, M.; Hoeh, W.; Lutz, R.; Vrijenhoek, R. DNA primers for amplification of mitochondrial cytochrome C oxidase subunit I from diverse metazoan invertebrates. *Mol. Mar. Biol. Biotechnol.* **1994**, *3*, 294–299. [PubMed]
33. Stigenberg, J.; Ronquist, F. Revision of the Western Palearctic Meteorini (Hymenoptera, Braconidae), with a molecular characterization of hidden Fennoscandian species diversity. *Zootaxa* **2011**, *3084*, 1–95. [CrossRef]
34. Ceccarelli, F.S.; Sharkey, M.J.; Zaldívar-Riverón, A. Species identification in the taxonomically neglected, highly diverse, neotropical parasitoid wasp genus *Notiospathius* (Braconidae: Doryctinae) based on an integrative molecular and morphological approach. *Mol. Phylogenetics Evol.* **2012**, *62*, 485–495. [CrossRef]
35. Ratnasingham, S.; Hebert, P.D. The Barcode of Life Data System. *Mol. Ecol. Notes* **2007**, *7*, 355–364. [CrossRef] [PubMed]
36. Ratnasingham, S.; Hebert, P.D. A DNA-based registry for all animal species: The Barcode Index Number (BIN) system. *PLoS ONE* **2013**, *8*, e66213. [CrossRef] [PubMed]
37. Kimura, M. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J. Mol. Evol.* **1980**, *16*, 111–120. [CrossRef]
38. Tamura, K.; Stecher, G.; Peterson, D.; Filipski, A.; Kumar, S. MEGA6: Molecular evolutionary genetics analysis version 6.0. *Mol. Biol. Evol.* **2013**, *30*, 2725–2729. [CrossRef]
39. Saitou, N.; Nei, M. The neighbor-joining method: A new method for reconstructing evolutionary trees. *Mol. Biol. Evol.* **1987**, *4*, 406–425.
40. Burns, J.; Janzen, D.H.; Hajibabaei, M.; Hallwachs, W.; Hebert, P.D. DNA barcodes of closely related (but morphologically and ecologically distinct) species of skipper butterflies (Hesperiidae) can differ by only one to three nucleotides. *J. Lepid. Soc.* **2007**, *61*, 138–153.
41. Janzen, D.H.; Burns, J.M.; Cong, Q.; Hallwachs, W.; Dapkey, T.; Manjunath, R.; Hajibabaei, M.; Hebert, P.D.N.; Grishin, N.V. Nuclear genomes distinguish cryptic species suggested by their DNA barcodes and ecology. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, 8313–8318. [CrossRef] [PubMed]
42. Janzen, D.W.H. DNA barcoding the Lepidoptera inventory of a large complex tropical conserved wildland, Area de Conservacion Guanacaste, northwestern Costa Rica. *Genome* **2016**, *59*, 641–660. [CrossRef]
43. Dyer, L.A.; Miller, J.S.; Rab Green, S.B.; Gentry, G.L.; Greeney, H.F.; Walla, T.W. Caterpillars and parasitoids of the Eastern Andes in Ecuador. Available online: http://www.caterpillars.org (accessed on 8 February 2021).
44. Säterberg, T.; Jonsson, T.; Yearsley, J.; Berg, S.; Ebenman, B. A potential role for rare species in ecosystem dynamics. *Sci. Rep.* **2019**, *9*, 11107. [CrossRef]
45. Janzen, D.H. Ecological characterization of a Costa Rican dry forest caterpillar fauna. *Biotropica* **1988**, *20*, 120–135. [CrossRef]
46. Janzen, D.H.; Gauld, I.D. Caterpillars (Lepidoptera: Saturnidae and Sphingidae) by Ichneumonid parasitoids (Hymenoptera) in Costa Rican dry forest. In *Forests and Insects*; Watt, A.D., Stork, N.E., Hunter, M.D., Eds.; Springer Science & Business Media: Berlin/Heidelberg, Germany, 1997; Volume 251.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.