

Supporting Information

Ultra-Fast Polarity Switching, Non-Radioactive Drift Tube for the Miniaturization of Drift-Time Ion Mobility Spectrometer

Lingfeng Li, Hao Gu, Yanzhen Lv, Yunjing Zhang, Xingli He and Peng Li *

School of Electronic and Information Engineering, Soochow University, Suzhou 215006, China; lingfengli@suda.edu.cn (L.L.); hguguhao@stu.suda.edu.cn (H.G.); lvyanzhen110@gmail.com (Y.L.);

yjzhang1223@suda.edu.cn (Y.Z.); hexingli@suda.edu.cn (X.H.)

* Correspondence: lipengsuda@suda.edu.cn; Tel.: +86-136-562-498-81


Contents

Figure S1. Photos of the drift tube device.

Figure S2. Example IMS spectra of 5 narcotics: (a) 10 ng cocaine; (b) 50 ng methamphetamine; (c) 50 ng heroin; (d) 50 ng morphine; (f) 50 ng marijuana.

Figure S3. Example IMS spectra of 6 explosives: (a) 50 ng black powder; (b) 50 ng PETN; (c) 100 ng HMX; (d) 50 ng RDX; (f) 100 ng HMTD; (g) 100 ng TATP.

(a)



(b)




Figure S1. Photos of the drift tube device.




Figure S2. Example IMS spectra of 5 narcotics: (a) 10 ng cocaine; (b) 50 ng methamphetamine; (c) 50 ng heroin; (d) 50 ng morphine; (f) 50 ng marijuana.




Figure S3. Example IMS spectra of 6 explosives: (a) 50 ng black powder; (b) 50 ng PETN; (c) 100 ng HMX; (d) 50 ng RDX; (e) 100 ng HMTD; (f) 100 ng TATP.