Table S1. Characteristics of the patients participating in the study

Anthropometric Data		Total $n = 175$ (100%)	GA Group $n = 35$ (20%)	M Group $n = 35$ (20%)	P Group $n = 35$ (20%)	PBB Group $n = 35$ (20%)	T Group $n = 35$ (20%)	p-Value
The results of the one-way analysis of variance (ANOVA)/the Kruskal–Wallis test by ranks								
Age X ± S M (Rk)	[years]	64.5 ± 11.7 66 (13)	65.1 ± 10.8 67 (9)	61.9 ± 11.9 63 (14)	66.1 ± 9.9 67 (8)	66.8 ± 12.1 69 (13)	62.7 ± 13.3 65 (14)	p = 0.25 [a] NS
High X ± S M (Rk)	[cm]	165.8 ± 8.7 165 (12)	166.9 ± 8.6 168 (14)	168 ± 7.4 170 (14)	163.4 ± 8.7 160 (12)	165.9 ± 8.3 164 (12)	164.7 ± 10.3 164 (18)	p = 0.18 [a] NS
Weight X ± S M (Rk)	[kg]	77.6 ± 15.9 75.5 (17)	83.4 ± 19.8 82 (20)	74.7 ± 14.9 74 (19)	74.1 ± 13.3 74 (22)	78.8 ± 16 75 (11)	77.1 ± 13.7 80 (21)	p = 0.19 [a] NS
BMI X ± S M (Rk)	[kg/m^2]	28.3 ± 5.4 27.5 (6.4)	29.9 ± 6.6 28.4 (5.3)	26.4 ± 4.6 25.3 (5.4)	27.9 ± 5.3 27.6 (7.7)	28.6 ± 5.1 27.1 (4.4)	28.5 ± 4.9 28.4 (7.3)	p = 0.05 [a]
The results of the χ^2 test of independence								
Gender n (%)	Female	97 (55.4)	18 (51.4)	15 (42.9)	24 (68.6)	21 (60)	19 (54.3)	p = 0.26 [b] NS
	Male	78 (44.6)	17 (48.6)	20 (57.1)	11 (31.4)	14 (40)	16 (45.7)	
Diabetes Mellitus	Insulin-dependent	53 (30.3)	11 (31.4)	6 (17.1)	12 (34.3)	12 (34.3)	12 (34.3)	p = 0.45 [b] NS
	Insulin-independent	45 (25.7)	10 (28.6)	3 (8.6)	12 (34.3)	9 (25.7)	11 (31.4)	p = 0.11 [b] NS
The results of the multiple proportions test								
BMI n (%)	Norm	50 (28.7)	5 (14.3)	15 (42.9)	14 (40)	7 (20.6)	9 (25.7)	p < 0.05 [c]
	Overweight	72 (41.4)	18 (51.4)	13 (37.1)	9 (25.7)	19 (55.9)	13 (37.1)	p = 0.09 [c] NS
	Obesity	52 (29.9)	12 (34.3)	7 (20)	12 (34.3)	8 (23.5)	13 (37.1)	p = 0.41 [c] NS

Results are presented as means ± standard deviations and medians (interquartile ranges). a One-way analysis of variance (ANOVA)/the Kruskal–Wallis test by ranks. Nominal data are presented as numbers (percentages). b χ^2 test of independence. c Multiple proportions test. BMI, body mass index. MAP—Mean Arterial Pressure; SAP—Systolic Arterial Pressure; DAP—Diastolic Arterial Pressure; PBB—preprocedural peribulbar block group; M group—metamizole group; P—Paracetam