

Correction

Correction: Benington et al. In Vitro Assessment of Wound-Healing Efficacy of Stabilized Basic Fibroblast Growth Factor (FGF-2) Solutions. *Pharmaceuticals* 2024, 17, 247

Leah Benington ¹ , Jingxin Mo ^{2,3}, Mingxin Li ^{2,4}, Gunesh Rajan ^{5,6} , Cornelia Locher ¹  and Lee Yong Lim ^{1,*} 

- ¹ Division of Pharmacy, School of Allied Health, University of Western Australia, Perth, WA 6009, Australia; leah.benington@uwa.edu.au (L.B.); connie.locher@uwa.edu.au (C.L.)
- ² Neuroscience Laboratory, The Affiliated Hospital of Guilin Medical University, Guilin 541001, China; jingxin.mo@hotmail.com (J.M.); mingxinli822@163.com (M.L.)
- ³ Graduate School of Biomedical Engineering, University of New South Wales, Sydney, NSW 2052, Australia
- ⁴ Department of Pharmacy, Tangshan Central Hospital, Tangshan 063000, China
- ⁵ Otolaryngology, Head & Neck Surgery, Division of Surgery, Medical School, University of Western Australia, Perth, WA 6009, Australia; gunesh.rajan@luks.ch
- ⁶ Department of Otolaryngology, Head & Neck Surgery, Luzerner Kantonsspital, 6000 Luzern, Switzerland
- * Correspondence: lee.lim@uwa.edu.au

In the original publication [1], there was a mistake in Figure 5. which was published. The image for the FGF-2 solution in vehicle 5 was mislabeled and required a replacement. A corrected version of Figure 5 appears below. The authors confirm that the scientific conclusions are unaffected. This correction was approved by the Academic Editor. The original publication has also been updated.



	Vehicle	Blank Vehicle	FGF-2 Solution
1			
2			

Figure 5. Cont.



Citation: Benington, L.; Mo, J.; Li, M.; Rajan, G.; Locher, C.; Lim, L.Y. Correction: Benington et al. In Vitro Assessment of Wound-Healing Efficacy of Stabilized Basic Fibroblast Growth Factor (FGF-2) Solutions. *Pharmaceuticals* 2024, 17, 247.

Pharmaceuticals **2024**, *17*, 892. <https://doi.org/10.3390/ph17070892>

Received: 28 June 2024

Accepted: 1 July 2024

Published: 5 July 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).




Figure 5. Fluorescence micrographs showing chemotactic migration of human dermal fibroblast cells to the basal surface of a transwell membrane in response to FGF-2. Cells seeded on the apical transwell membrane were exposed to FGF-2 solutions or blank vehicles added to the basolateral chamber. Samples comprise blank vehicles or FGF-2 (50 ng/mL in the vehicle) in the vehicles: 1 (water only), 2 (methylcellulose (MC) 0.05% *w/v*), 3 (alanine 20 mM), 4 (human serum albumin (HSA) 1 mg/mL), 5 (MC 0.05% *w/v* and alanine 20 mM) and 6 (MC 0.05% *w/v* and HSA 1 mg/mL). Magnification: 200 $\times$, scale bar = 100 μ m.

Reference

1. Benington, L.; Mo, J.; Li, M.; Rajan, G.; Locher, C.; Lim, L.Y. In Vitro Assessment of Wound-Healing Efficacy of Stabilized Basic Fibroblast Growth Factor (FGF-2) Solutions. *Pharmaceuticals* **2024**, *17*, 247. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.