

Article from:

Omics in Plant Disease Resistance

Current Issues in Molecular Biology. Volume 19 (2016). Focus Issue

DOI: <http://dx.doi.org/10.21775/9781910190357>

Edited by:

Vijai Bhaduria

Crop Development Centre/Dept. of Plant Sciences
51 Campus Drive
University of Saskatchewan
Saskatoon, SK S7N 5A8 Canada.
Tel: (306) 966-8380 (Office), (306) 716-9863 (Cell)
Email: vijai.bhaduria@usask.ca

Copyright © 2016

Caister Academic Press, U.K. www.caister.com

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the publisher. No claim to original government works.

Current publications of interest

Microalgae

Current Research and Applications

Edited by: MN Tsaloglou

152 pp, January 2016

Book: ISBN 978-1-910190-27-2 £129/\$259

Ebook: ISBN 978-1-910190-28-9 £129/\$259

The latest research and newest approaches to the study of microalgae.

Bacteria-Plant Interactions

Advanced Research and Future Trends

Edited by: J Murillo, BA Vinatzer, RW Jackson, et al.

x + 228 pp, March 2015

Book: ISBN 978-1-908230-58-4 £159/\$319

Ebook: ISBN 978-1-910190-00-5 £159/\$319

"an up-to-date overview" (Ringgold)

Microarrays

Current Technology, Innovations and Applications

Edited by: Z He

x + 246 pp, August 2014

Book: ISBN 978-1-908230-49-2 £159/\$319

Ebook: ISBN 978-1-908230-59-1 £159/\$319

"a valuable and useful source ...

recommended" (Biotechnol. Agron. Soc. Environ.)

Metagenomics of the Microbial Nitrogen Cycle

Theory, Methods and Applications

Edited by: D Marco

xiv + 268 pp, September 2014

Book: ISBN 978-1-908230-48-5 £159/\$319

Ebook: ISBN 978-1-908230-60-7 £159/\$319

"a strong overview" (Ringgold)

Proteomics

Targeted Technology, Innovations and Applications

Edited by: M Fuentes, J LaBaer

x + 186 pp, September 2014

Book: ISBN 978-1-908230-46-1 £159/\$319

Ebook: ISBN 978-1-908230-62-1 £159/\$319

"many excellent chapters" (Doodys)

Applied RNAi

From Fundamental Research to Therapeutic Applications

Edited by: P Arbuthnot, MS Weinberg

x + 252 pp, June 2014

Book: ISBN 978-1-908230-43-0 £159/\$319

Ebook: ISBN 978-1-908230-67-6 £159/\$319

"Essential reading" (Biotechnol Agron Soc Environ);

"recommended" (Fungal Diversity); "an excellent

resource" (Doodys)

Next-generation Sequencing

Current Technologies and Applications

Edited by: J Xu

xii + 160 pp, March 2014

Book: ISBN 978-1-908230-33-1 £120/\$240

Ebook: ISBN 978-1-908230-95-9 £120/\$240

"written in an accessible style" (Zentralblatt Math);

"recommend this book to all

investigators" (ChemMedChem)

Omics in Soil Science

Edited by: P Nannipieri, G Pietramellara, G Renella

x + 198 pp, January 2014

Book: ISBN 978-1-908230-32-4 £159/\$319

Ebook: ISBN 978-1-908230-94-2 £159/\$319

"a recommended reference" (Biotechnol. Agron. Soc.

Environ.); "a must for Soil scientists" (Fungal

Diversity)

Genome Analysis

Current Procedures and Applications

Edited by: MS Poptsova

xiv + 374 pp, January 2014

Book: ISBN 978-1-908230-29-4 £159/\$319

Ebook: ISBN 978-1-908230-68-3 £159/\$319

An up-to-date and comprehensive overview of next-generation sequencing data analysis, highlighting problems and limitations, applications and developing trends in various fields of genome research.

RNA Editing

Current Research and Future Trends

Edited by: S Maas

viii + 240 pp, June 2013

Book: ISBN 978-1-908230-23-2 £159/\$319

Ebook: ISBN 978-1-908230-88-1 £159/\$319

"an essential book" (Doodys)

Real-Time PCR

Advanced Technologies and Applications

Edited by: NA Saunders, MA Lee

viii + 284 pp, July 2013

Book: ISBN 978-1-908230-22-5 £159/\$319

Ebook: ISBN 978-1-908230-87-4 £159/\$319

"an invaluable reference" (Doodys); "wide range of real

time PCR technologies" (Food Sci Technol Abs); "I was

impressed by this text" Aus J Med Sci

Bionanotechnology

Biological Self-assembly and its Applications

Edited by: BHA Rehm

x + 310 pp, February 2013

Book: ISBN 978-1-908230-16-4 £159/\$319

Ebook: ISBN 978-1-908230-81-2 £159/\$319

"the most striking and successful approaches" Book

News

Linking Biomarker and Comparative Omics to Pathogens in Legumes

Marwan Diapari

Southern Crop Protection and Food Research Centre, Agriculture and Agri-Food Canada / Agriculture et Agroalimentaire Canada, 1391 Sandford St., London, ON, N5V 4T3 Canada. Email Marwan.Diapari@agr.gc.ca
DOI: <http://dx.doi.org/10.21775/9781910190357.13>

Abstract

It is envisioned that a more precise study of the association between the traits and biomarkers will dramatically decrease the time and costs required to bring new improved disease resistance lines to market. The field of omics has an enormous potential to assess diseases more precise, including the identification and understanding of pathogenic mechanisms in legume crops, and have been exemplified by a relatively large number of studies. Recently, molecular genetic studies have accumulated a huge amount of genotypic data, through a more affordable next generation sequencing (NGS) technology, causing the omics approaches to fall behind. In this paper I provide an overview of genomics and proteomics and their use in legume crops, including the use of comparative genomics to identify homologous markers within legume crops.

Introduction

Legumes are economically important, rich source of quality protein for humans and animals, and enrich the soil by producing their own nitrogen in symbiosis with nitrogen-fixing bacteria (Singh et al., 2007). A better knowledge of its physiological characteristics and behaviour within its own environment is imperative in order to increase legume competitiveness and to adapt its production to the new constraints arising from the climatic changes and the occurrence of new diseases (Adam-Blondon, 2007).

Molecular approaches (OMICS) that involve mRNA (transcriptomics), proteins (proteomics), metabolites (metabolomics), are indispensable in the era of functional genomics, and have been exemplified by a relatively large number of studies, such as gene expression, genetic diversity, phylogeny, comparative genomics and epigenetics (Ma et al., 2013; Sabir et al., 2014). OMICS has brought the possibilities for genetic improvement, conservation of biodiversity, and imparting disease resistance in crops for further targeted breeding (Rubiales et al., 2015). Maintaining genetic diversity and capturing genetic improvement are often through structuring populations into a genetic hierarchy, setting population sizes, and managing pedigrees and inbreeding.

Plant diseases can drastically abate the crop yields as the degree of disease outbreak is getting severe around the world. Major diseases are rusts, powdery and downy

mildews, ascochyta blights, botrytis gray molds, anthracnose, damping-off, root rots, collar rots, vascular wilts and white molds (Gururani et al., 2012; Rubiales et al., 2015). New disease occurrence can involve gene duplication, transfer or recombination (Hollomon and Brent, 2009). The negative impact of diseases and the genetic mechanisms and gene interactions involved in the resistance response are still poorly understood (Cantu et al., 2013). Comprehensive body of knowledge and understanding of the cellular functions will aid the development of novel practical approaches to enhance disease resistance (Choi and Hwang, 2014).

Legume crops have been exploited by breeding program (Robertson et al., 1996), including for disease purpose breeding program. It is believed that wild relatives are considered as important source of genetic diversity for agriculture, although their gene repertoire remains largely unexplored (Li et al., 2014). Proposed draft legume genomes allow the prediction of the full complement of genes, phylogenetic relationships and chromosomal locations will provide breeders and scientists with a useful tool to identify novel disease resistance traits (Andolfo et al., 2014). It is expected the array will accelerate hypothesis generation and gene discovery in disease defense pathways, responses to abiotic stresses, development, and evolutionary diversity in plants (Close et al., 2004). Directed molecular evolution can be used to develop enabling technologies for plant biologists; how genes can be optimized to generate improved input traits such as those conferring insect tolerance, disease control and herbicide tolerance; and how plant quality can be altered to improve yield (Lassner and McElroy, 2002). Molecular genetics provide an unparalleled opportunity to understand how genes and genetic changes interact with environmental stimuli to either preserve health or cause disease (Schwartz et al., 2004).

Plant breeders strive to stay ahead of the evolving pathogens by releasing cultivars with new resistance genes or gene combinations (Webb and Fellers, 2006). Breeding for disease resistance in crops has mainly been accomplished by incorporating single resistance gene. There are advantages to quantitative resistance in terms of durability. New technologies in genomics and proteomics are providing insights into disease-resistance pathways (Jordan et al., 2006). Genetic improvement of the crop is being actively pursued and numerous functional genomics studies aimed at characterizing gene controlling key agronomic characteristics and providing some insight into the basis of disease resistances (Hulbert and Pumphrey, 2014; Mellor et al., 2012).

The fields of environmental genetics and environmental genomics has enormous potential to affect our ability to accurately assess the risk of developing disease, identify and understand basic pathogenic mechanisms that are critical to disease progression (Schwartz et al., 2004). Genetic studies pathogens during the past decade have made it an excellent system for investigating pathogen-host interactions (Xu and Xue, 2002).

Genetic variation has an important role in determining biologic phenotypes and disease resistance (Sladek and Hudson, 2006). Inter-genomic comparisons identified lineage-specific genes, some of which show evidence of positive selection and may contribute to variation of agronomic traits such as disease resistance (Li et al., 2014). Scarcity of genetic polymorphism makes crops especially vulnerable to a wide variety of pathogens (Ratnaparkhe et al., 2011). Sequence analysis of plant disease resistance genes shows similarity among themselves (Ramachandra et al., 2011) and are present in abundance and majority of these encode proteins with a nucleotide-binding site (NBS) (Basak et al., 2007; Ribas et al., 2011), comprising up to 2% of all genes (Mun et al., 2009).

NBS encoding resistance genes are key plant disease-resistance genes (R-genes) represent the major class of disease resistance genes (Mun et al., 2009; Richly et al., 2002) also play an important role in offering resistance to pathogens (Yu et al., 2014). Numerous plant R-genes have been used with varying degree of success in crop improvement (Gururani et al., 2012). A number of studies have focused on NBS-encoding genes in disease resistant breeding programs for diverse plants (Wei et al., 2013). Plant possesses a large number of R-genes with diverse recognition specificities which are activated in response to a variety of microbial pathogens (Sharma et al., 2014) and have the ability to detect a pathogen attack and facilitate a counter attack against the pathogen (Gururani et al., 2012; Martin et al., 2003) also often show significant similarity amongst themselves in terms of both their DNA sequences and structural motifs present in their protein products (López et al., 2003).

The majority of verified plant disease resistance genes isolated to date are of the NBS-LRR class (Madsen et al., 2003). The NBS-LRR genes are the largest class of disease resistance genes in plants (Wei et al., 2013; Xu et al., 2011). NBS profiling, in particular, targets conserved nucleotide binding site-encoding sequences of resistance gene analogs (RGAs), and is widely used to identify molecular markers for R genes (Vossen et al., 2013).

Reactive oxygen species (ROS) signaling network mediating disease resistance in plants information became available, thanks to the complete Arabidopsis (*Arabidopsis thaliana*) and rice (*Oryza sativa*) genome sequences (Kotchoni and Gachomo, 2006). ROS play multiple roles in interactions between plants and microbes (Daub et al., 2013) and in the plant hypersensitive disease resistance response (Cecconi et al., 2009), including induced systemic resistances (Deepak et al., 2008).

Genomic and Proteomics

Recent advances in metabolomics, genomics, transcriptomic and metabolic modelling offer new opportunities to address this question and generate a system-level understanding of metabolic interactions at the host-pathogen interface (Rico et al., 2011). Functional genomics represents a systematic approach to elucidating the function of the novel genes revealed by complete genome sequences. The different approaches are grouped into four domains: genome, transcriptome, proteome, and metabolome (Oliver, 2002). Genomics is a new and promising area, which can broadly be defined as the application of high throughput genomics and functional genomic technologies to the study of pathology in plants (Brown and van der Ouderaa, 2007). Access to complete genomic sequences, coupled with rapidly accumulating data related to RNA and protein expression patterns, has made it possible to determine comprehensively how genes contribute to complex phenotypes (Close et al., 2004).

The huge amount of genomic data now becoming available offers both opportunities and challenges for breeding program (Garrett et al., 2006). Innovative approaches may be required to sequence, assemble, annotate and analyse host and pathogen genomes. Overcoming these challenges will enable scientists to investigate the genes and genome organisation may ultimately lead to new solutions for control diseases legume (Van Zee et al., 2007). The methods utilised provide new opportunities for studying the nature and role of defence mechanisms in plants (Collinge et al., 2008).

Knowledge of structural and functional genomics and genes and comparative genomics with model species has increased our understanding genomic diversity through integration of genetical, evolutionary and structural data (Heslop-Harrison and Schwarzacher, 2007). We will also gain a better understanding of the genes responsible for disease resistance. With a collection of desirable genes in mind, we can again use genomics as a diagnostic tool to search for these genes in the wide variety and follow their transfer by classical breeding (Regan et al., 2006). Following largescale source of sequencing of libraries and bioinformatics searches in public databases pave the way for the design and conduct of large-scale experiments that will help to a better understanding the complex biology of disease resistance (Bernier, 2004) and facilitating the isolation of novel genes, as well as by helping to identify targets for chemical control (Kamoun et al., 2002).

DNA sequencing is now complete for some hosts and several pathogens to form predictions about epidemic features and outcomes and for understanding host resistance and pathogen evolution (Garrett et al., 2006). The use of large-scale complementary DNA library constructions and genome-wide transcript profiles of plants exposed to biotic stress provide the data required to drive hypotheses concerning the function of newly identified genes (Collinge et al., 2008). For example, an extensive programme of functional genomics is being undertaken through the systematic analysis of insertional mutants in Arabidopsis (Delseny and Pelletier, 2001).

Proteins are involved in disease resistance and defense (Mooney et al., 2006). In recent years, proteomics has been used to gain in-depth understanding of many aspects of the host defense against pathogens to identify differentially accumulated proteins in a mutant and a wild type (Chen et al., 2013; Zimaro et al., 2011). Fueled by ever growing DNA sequence information, proteomics is providing insight into the mechanisms of biological processes in a high-throughput mode (Sharma et al., 2008). Proteomic approach was employed to investigate the cold stress-responsive proteins in citrus (Long et al., 2012). Various proteomic techniques for isolation and purification of proteins have been developed; each tailored to preserve protein properties relevant to study of a particular disease type (Vo and Palsson, 2007). The application of combined proteomic and RNA interfering analyses is an efficient strategy to identify genes required disease resistance and probably other biological processes in plants (Xu et al., 2012).

The recent development of large scale phenotyping, genome sequencing and analysis of gene, protein and metabolite expressions can be of great help to further decipher plant-pathogen interactions and identify key resistance components that may be introgressed into crop plants through breeding (Rubiales et al., 2015). With the sequencing of entire genomes it has become technically feasible to study transcription on a global scale (Barnett and Fisher, 2006). In 2001, The Medicago Genome Initiative (MGI) is a database of EST sequences of the model legume *Medicago truncatula* (<http://www.noble.org>), which is taking a global approach in studying the genetic and biochemical events associated with the growth, development and environmental interactions of this model legume (Bell et al., 2001). Genome sequencing of the model legumes, *Medicago truncatula* and *Lotus japonicus* provides an opportunity for large-scale sequence-based comparison of two genomes in the same plant family (Cannon et al., 2006). Recent genome sequencing enables mega-base scale comparisons between related genomes. In 2013, chickpea draft sequence genome using next-generation sequencing platforms, bacterial artificial chromosome end sequences and a genetic map has been proposed. The draft genome sequence is expected to facilitate genetic enhancement and breeding to develop improved chickpea varieties (Jain et al., 2013). In the same year whole genome shotgun sequence of CDC Frontier, a kabuli chickpea variety, this contains an estimated 28,269 genes published. Candidate genes for disease resistance and agronomic traits are highlighted, including traits that distinguish the two main market classes of cultivated chickpea—desi and kabuli. These data comprise a resource for chickpea improvement through molecular breeding and provide insights into both genome diversity and domestication (Varshney et al., 2013). The genome of *V. faba* is exceedingly large (ca. 13 Gb) provides a valuable tool for future transcript profiling and also enriches the genetic resources available for this important legume crop species (Arun-Chinnappa and McCurdy, 2015). Development of bacterial artificial chromosome (BAC) library in White clover, sought a large-scale BAC-end

sequencing strategy has the potential to anchor a significant proportion of the genome of white clover onto the gene-space sequence of *M. truncatula* (Febrer et al., 2007). The first draft of the genome sequence of pigeonpea, and development of markers will be useful for fingerprinting and diversity analysis of pigeonpea germplasm and molecular breeding applications (Dubey et al., 2011; Singh et al., 2012). In 2012, Illumina next-generation sequencing platform to generate 237.2 Gb of sequence applied in pigeonpea (Varshney et al., 2012).

In the whole genome sequencing, genetic map provides an essential framework for accurate and efficient genome assembly and validation. The analysis revealed an overall conserved marker order, although some localized inversions between draft genome assembly and the genetic map were detected (Deokar et al., 2014). Many vectors, especially artificial chromosome vectors, have been developed for genome-scale mapping and sequencing. These data showed that this genomic library was reliable for further molecular research in *Lotus japonicus* (Guo Xi-ZhiGuo et al., 2004).

Comparative Genomics

Within the legume plant family, glimpses of synteny have also been observed. Characterizing syntenic relationships in legumes is important in transferring knowledge from model legumes to crops that are important sources of protein, fixed nitrogen, and health-promoting compounds (Mudge et al., 2005). The *Lotus* and *Medicago* genomes share a minimum of 10 large-scale synteny blocks, each with substantial collinearity and frequently extending the length of whole chromosome arms (Cannon et al., 2006). 56 conserved synteny blocks were identified between eggplant and tomato (Hirakawa et al., 2014). The December 22 2011 issue of *Nature* published the result of syntenic relationships between *Medicago*, *Glycine*, *Lotus* and *Vitis* illustrated in Circos diagram (Young et al., 2011) (Figure 1). Phylogenetic interpretation of DNA sequence data from multiple genomic regions has become the gold standard for species delimitation and population genetics (Kohn, 2004).

The interaction between genomics and quantitative genetics has been a two-way street. Genomics contributed genetic markers and genetic maps making it possible to study quantitative trait loci (QTLs), and quantitative genetics contributed new theories and computational techniques to deal with the data generated by QTL studies (van Buijtenen, 2001). It will be possible to integrate-based genomic tools into approaches as genetic approaches aiming at developing efficient marker-assisted selection strategies for the development of resistant varieties for example, or for designing models for predicting the behaviour of the plant in the field for greater resistance (Adam-Blondon, 2007).

Gene discovery holds enormous promise, but depends heavily on comparative genomics, capitalizing on genomic information (Burdon and Wilcox, 2007). Comparative genomics approach are often used to investigate crops with narrow genetic base of cultivars, and to evaluate the




Figure 1. Homologous gene pairs were identified for all pairwise comparisons between *M. truncatula* (Mt), *G. max* (Gm), *L. japonicus* (Lj) and *V. vinifera* (Vv) genomes, syntenic regions are associated with the ancestral whole-genome duplication (WGD) occurred approximately 58 Myr ago (Young et al., 2011).

evolution of a complex gene cluster that is associated with several disease resistance (R) genes of known function (Ashfield et al., 2012; Bertoli et al., 2003; Sharma et al., 2013). Comparative genomics provides a tool to utilize the exponentially increasing sequence information from model plants to clone agronomically important genes from less studied crop species (Huang et al., 2005). Comparative genomics also plays a very important role in identification of genes/QTLs linked to the resistance genes (Babu et al., 2014). With access to sequences various model organisms and many important microbial pathogens, structural biology is on the verge of a dramatic transformation. Structural genomics will yield a large number of experimental protein structures (tens of thousands) and an even larger number of calculated comparative protein structure models (millions) (Burley, 2000).

The emergence of "omic" technologies and the establishment of model legume plants, such as *Medicago truncatula* and *Lotus japonicus* are promising strategies for understanding the molecular genetic basis of stress

resistance (Dita et al., 2006). The legume species *M. truncatula* is a model plant to help in the identification of agronomically important genes in crop legumes, e.g. disease resistance (Journet et al., 2001).

QTL mapping and synteny analysis identified genomic loci harboring resistance factors has offered considerable promise and experimental power with varying degrees of success (Colwell, 2002; Muchero et al., 2011). The potential of GWAS approaches likely bearing resistance alleles of significant effect useful in breeding programs (Debibakas et al., 2014). Genome-wide analyses of gene function and gene expression are beginning to yield valuable information in many areas of biological research, and these genomic tools are now being applied to crop pest and disease research (Keon et al., 2003), thus strategy of applying a sequence-to-phenotype (or functional genomics) paradigm applied to the discovery of R-genes (Kamoun et al., 2002).

Conclusions

Current endeavours to find disease-causing genes have received a growing appreciation of the extent of conservation of genes, the extent to which synteny will be a useful tool throughout all legume species. The impact of the discovery of conserved synteny is also most important in the consideration of genomics programs for the major legume genomes. Genomics, through deciphering allelic effects on phenotypes and identifying patterns of adaptive variation at the landscape level, will in the future constitute a useful tool, if cost-effective, to design conservation strategies for plants (Gonzalez-Martinez et al., 2006), including disease resistance breeding program. However, the discovery still needs to be validated in a much larger sample size, where a high hurdle that many putative biomarkers fail to make. Poste (Poste, 2011) explained that although over 150,000 papers claimed to have found disease biomarkers in human, only 100 biomarkers were routinely used in clinic. Perhaps the same case may occurred in legumes, and future research on linking biomarkers comparative omics will help to fulfill this challenge.

References

- Adam-Blondon, A.-F. (2007). What benefits can viticulture possibly expect from genomics? Quels benefices la genomique peut-elle apporter a la viticulture? [French]. *Cahiers Agricultures* 16, 87-92.
- Andolfo, G., Jupe, F., Witek, K., Etherington, G.J., Ercolano, M.R., and Jones, J.D.G. (2014). Defining the full tomato NB-LRR resistance gene repertoire using genomic and cDNA RenSeq. *BMC Plant Biol* 14.
- Arun-Chinnappa, K.S., and McCurdy, D.W. (2015). De novo assembly of a genome-wide transcriptome map of *Vicia faba* (L.) for transfer cell research. *Front Plant Sci* 6, 217.
- Ashfield, T., Egan, A.N., Pfeil, B.E., Chen, N.W.G., Podicheti, R., Ratnaparkhe, M.B., Ameline-Torregrosa, C., Denny, R., Cannon, S., Doyle, J.J., et al. (2012). Evolution of a complex disease resistance gene cluster in diploid phaseolus and tetraploid glycine. *Plant Physiol* 159, 336-354.
- Babu, B.K., Dinesh, P., Agrawal, P.K., Sood, S., Chandrashekar, C., Bhatt, J.C., and Kumar, A. (2014). Comparative genomics and association mapping approaches for blast resistant genes in finger millet using SSRs. *PLoS One* 9.
- Barnett, M.J., and Fisher, R.F. (2006). Global gene expression in the rhizobial-legume symbiosis. *Symbiosis* 42, 1-24.
- Basak, J., Kundu, S., and Pal, A. (2007). Phylogenetic analysis of divergent structural organization of nucleotide binding domain encoded by resistance genes and gene homologs in the family Fabaceae. *Ind J Biotechnol* 6, 9-17.
- Bell, C.J., Dixon, R.A., Farmer, A.D., Flores, R., Inman, J., Gonzales, R.A., Harrison, M.J., Paiva, N.L., Scott, A.D., Weller, J.W., et al. (2001). The Medicago Genome Initiative: A model legume database. *Nucleic Acids Res* 29, 114-117.
- Bernier, L. (2004). Genomics of ophiostomatoid fungi: the first two thousand genes. *Phytoprotection* 85, 39-43.
- Bertioli, D.J., Leal-Bertioli, S.C.M., Lion, M.B., Santos, V.L., Pappas Jr, G., Cannon, S.B., and Guimarães, P.M. (2003). A large scale analysis of resistance gene homologues in *Arachis*. *Mol Genet Genomics* 270, 34-45.
- Brown, L., and van der Ouderaa, F. (2007). Nutritional genomics: food industry applications from farm to fork. *Br J Nutr* 97, 1027-1035.
- Burdon, R.D., and Wilcox, P.L. (2007). Population management: potential impacts of advances in genomics. *New Forests* 34, 187-206.
- Burley, S.K. (2000). An overview of structural genomics. *Nat Struct Biol* 7, 932-934.
- Cannon, S.B., Sterck, L., Rombauts, S., Sato, S., Cheung, F., Gouzy, J., Wang, X., Mudge, J., Vasdewani, J., Scheix, T., et al. (2006). Legume genome evolution viewed through the *Medicago truncatula* and *Lotus japonicus* genomes. *Proc Natl Acad Sci U S A* 103, 14959-14964.
- Cantu, D., Yang, B., Ruan, R., Li, K., Menzo, V., Fu, D., Chern, M., Ronald, P.C., and Dubcovsky, J. (2013). Comparative analysis of protein-protein interactions in the defense response of rice and wheat. *BMC Genomics* 14.
- Cecconi, D., Orzetti, S., Vandelle, E., Rinalducci, S., Zolla, L., and Delledonne, M. (2009). Protein nitration during defense response in *Arabidopsis thaliana*. *Electrophoresis* 30, 2460-2468.
- Chen, C., Chen, H., Shan, J.X., Zhu, M.Z., Shi, M., Gao, J.P., and Lin, H.X. (2013). Genetic and physiological analysis of a novel type of interspecific hybrid weakness in rice. *Mol Plant* 6, 716-728.
- Choi, H.W., and Hwang, B.K. (2014). Molecular and cellular control of cell death and defense signaling in pepper. *Planta* 241.
- Close, T.J., Wanamaker, S.I., Caldo, R.A., Turner, S.M., Ashlock, D.A., Dickerson, J.A., Wing, R.A., Muehlbauer, G.J., Kleinhofs, A., and Wise, R.P. (2004). A new resource for cereal genomics: 22K Barley GeneChip comes of age. *Plant Physiol* 134, 960-968.
- Collinge, D.B., Jensen, M.K., Lyngkjaer, M.F., and Rung, J. (2008). How can we exploit functional genomics approaches for understanding the nature of plant defences? Barley as a case study. *Eur J Plant Pathol* 121, 257-266.
- Colwell, R.R. (2002). Fulfilling the promise of biotechnology. *Biotechnol Adv* 20, 215-227.
- Daub, M.E., Herrero, S., and Chung, K.R. (2013). Reactive oxygen species in plant pathogenesis: The role of perylenequinone photosensitizers. *Antioxid Redox Signal* 19, 970-989.
- Debikakas, S., Rocher, S., Garsmeur, O., Toubi, L., Roques, D., D'Hont, A., Hoarau, J.Y., and Daugrois, J.H. (2014). Prospecting sugarcane resistance to Sugarcane yellow leaf virus by genome-wide association. *Theor Appl Genet* 127, 1719-1732.
- Deepak, S.A., Shibato, J., Ishii, H., Ogawa, Y., Yoshida, Y., Iwahashi, H., Masuo, Y., Agrawal, G.K., and Rakwal, R. (2008). Proteomics approach for investigating the disease resistance using cucumber as model plant. *Am J Biochem Biotechnol* 4, 231-238.

- Delseny, M., and Pelletier, G. (2001). From Arabidopsis to rice genomics: A survey of French programmes. *Comptes Rendus de l'Académie des Sciences Serie III Sciences de la Vie* 324, 1103-1110.
- Deokar, A.A., Ramsay, L., Sharpe, A.G., Diapari, M., Sindhu, A., Bett, K., Warkentin, T.D., and Tar'an, B. (2014). Genome wide SNP identification in chickpea for use in development of a high density genetic map and improvement of chickpea reference genome assembly. *BMC Genomics* 15, 708.
- Dita, M.A., Rispail, N., Prats, E., Rubiales, D., and Singh, K.B. (2006). Biotechnology approaches to overcome biotic and abiotic stress constraints in legumes. *Euphytica* 147, 1-24.
- Dubey, A., Farmer, A., Schlueter, J., Cannon, S.B., Abernathy, B., Tuteja, R., Woodward, J., Shah, T., Mulasmanovic, B., Kudapa, H., *et al.* (2011). Defining the transcriptome assembly and its use for genome dynamics and transcriptome profiling studies in pigeonpea (*Cajanus cajan* L.). *DNA research : an international journal for rapid publication of reports on genes and genomes* 18, 153-164.
- Febrer, M., Cheung, F., Town, C.D., Cannon, S.B., Young, N.D., Abberton, M.T., Jenkins, G., and Milbourne, D. (2007). Construction, characterization, and preliminary BAC-end sequencing analysis of a bacterial artificial chromosome library of white clover (*Trifolium repens* L.). *Genome* 50, 412-421.
- Garrett, K.A., Hulbert, S.H., Leach, J.E., and Travers, S.E. (2006). Ecological genomics and epidemiology. *Eur J Plant Pathol* 115, 35-51.
- Gonzalez-Martinez, S.C., Krutovsky, K.V., and Neale, D.B. (2006). Forest-tree population genomics and adaptive evolution. *New Phytol* 170, 227-238.
- Guo Xi-ZhiGuo, X.-Z., Liu Yao-GuangLiu, Y.-G., and Luo DaLuo, D. (2004). The construction and screening of genomic library based on transformable artificial chromosome (TAC) vector in *Lotus japonicus*. *Zhiwu Shengli yu Fenzi Shengwuxue Xuebao* 30, 234-238.
- Gururani, M.A., Venkatesh, J., Upadhyaya, C.P., Nookaraju, A., Pandey, S.K., and Park, S.W. (2012). Plant disease resistance genes: Current status and future directions. *Physiol Mol Plant Pathol* 78, 51-65.
- Heslop-Harrison, J.S., and Schwarzacher, T. (2007). Domestication, genomics and the future for banana. *Ann Bot* 100, 1073-1084.
- Hirakawa, H., Shirasawa, K., Miyatake, K., Nunome, T., Negoro, S., Ohyama, A., Yamaguchi, H., Sato, S., Isobe, S., Tabata, S., *et al.* (2014). Draft genome sequence of eggplant (*Solanum melongena* L.): the representative solanum species indigenous to the old world. *DNA research : an international journal for rapid publication of reports on genes and genomes* 21, 649-660.
- Hollomon, D.W., and Brent, K.J. (2009). Combating plant diseases - The Darwin connection. *Pest Manage Sci* 65, 1156-1163.
- Huang, S., Van Der Vossen, E.A.G., Kuang, H., Vleeshouwers, V.G.A.A., Zhang, N., Borm, T.J.A., Van Eck, H.J., Baker, B., Jacobsen, E., and Visser, R.G.F. (2005). Comparative genomics enabled the isolation of the R3a late blight resistance gene in potato. *Plant J* 42, 251-261.
- Hulbert, S., and Pumphrey, M. (2014). A time for more booms and fewer busts? unraveling cereal-rust interactions. *Mol Plant-Microbe Interact* 27, 207-214.
- Jain, M., Misra, G., Patel, R.K., Priya, P., Jhanwar, S., Khan, A.W., Shah, N., Singh, V.K., Garg, R., Jeena, G., *et al.* (2013). A draft genome sequence of the pulse crop chickpea (*Cicer arietinum* L.). *The Plant journal : for cell and molecular biology* 74, 715-729.
- Jordan, M.C., Cloutier, S., Somers, D., Procunier, D., Rampitsch, C., and Xing, T. (2006). Beyond R genes: dissecting disease-resistance pathways using genomics and proteomics. *Can J Plant Pathol* 28, S228-S232.
- Journet, E.P., Carreau, V., Gouzy, J., Thoquet, P., Rosenberg, C., Barker, D., Huguet, T., Dénarié, J., and Gamas, P. (2001). Génomique de la légumineuse modèle *medicago truncatula* : État des lieux et perspectives. *OCL OI Corps Lipides* 8, 478-484.
- Kamoun, S., Dong, S., Hamada, W., Huitema, E., Kinney, D., Morgan, W.R., Styer, A., Testa, A., and Torto, T.A. (2002). From sequence to phenotype: Functional genomics of *Phytophthora*. *Can J Plant Pathol* 24, 6-9.
- Keon, J., Curtis, R., Cabrera, H., and Hargreaves, J. (2003). A genomics approach to crop pest and disease research. *Pest Manage Sci* 59, 143-148.
- Kohn, L.M. (2004). Applying comparative genomics to plant disease epidemiology. *Phytoprotection* 85, 45-48.
- Kotchoni, S.O., and Gachomo, E.W. (2006). The reactive oxygen species network pathways: An essential prerequisite for perception of pathogen attack and the acquired disease resistance in plants. *J Biosci* 31, 389-404.
- Lassner, M.W., and McElroy, D. (2002). Directed molecular evolution: Bridging the gap between genomics leads and commercial products. *OMICS A Journal of Integrative Biology* 6, 153-162.
- Li, Y.H., Zhou, G., Ma, J., Jiang, W., Jin, L.G., Zhang, Z., Guo, Y., Zhang, J., Sui, Y., Zheng, L., *et al.* (2014). De novo assembly of soybean wild relatives for pan-genome analysis of diversity and agronomic traits. *Nat Biotechnol* 32, 1045-1052.
- Long, G., Song, J., Deng, Z., Liu, J., and Rao, L. (2012). Ptcp gene induced by cold stress was identified by proteomic analysis in leaves of *poncirus trifoliata* (L.) Raf. *Mol Biol Rep* 39, 5859-5866.
- López, C.E., Zuluaga, A.P., Cooke, R., Delseny, M., Tohme, J., and Verdier, V. (2003). Isolation of resistance gene candidates (RGCs) and characterization of an RGC cluster in cassava. *Mol Genet Genomics* 269, 658-671.
- Ma, N.L., Rahmat, Z., and Lam, S.S. (2013). A Review of the Omics Approach to Biomarkers of Oxidative Stress in *Oryza sativa*. *Int J Mol Sci* 14, 7515-7541.
- Madsen, L.H., Collins, N.C., Rakwalska, M., Backes, G., Sandal, N., Krusell, L., Jensen, J., Waterman, E.H., Jahoor, A., Ayliffe, M., *et al.* (2003). Barley disease resistance gene analogs of the NBS-LRR class: Identification and mapping. *Mol Genet Genomics* 269, 150-161.
- Martin, G.B., Bogdanove, A.J., and Sessa, G. (2003). Understanding the Functions of Plant Disease Resistance Proteins. In *Annu Rev Plant Biol*, pp. 23-61.
- Mellor, K.E., Hoffman, A.M., and Timko, M.P. (2012). Use of *ex vitro* composite plants to study the interaction of

- cowpea (*Vigna unguiculata* L.) with the root parasitic angiosperm *Striga gesnerioides*. *Plant Methods* 8.
- Mooney, B.P., Miernyk, J.A., Greenlief, C.M., and Thelen, J.J. (2006). Using quantitative proteomics of *Arabidopsis* roots and leaves to predict metabolic activity. *Physiol Plant* 128, 237-250.
- Muchero, W., Ehlers, J.D., Close, T.J., and Roberts, P.A. (2011). Genic SNP markers and legume synteny reveal candidate genes underlying QTL for *Macrophomina phaseolina* resistance and maturity in cowpea [*Vigna unguiculata* (L) Walp.]. *BMC Genomics* 12.
- Mudge, J., Cannon, S.B., Kalo, P., Oldroyd, G.E., Roe, B.A., Town, C.D., and Young, N.D. (2005). Highly syntenic regions in the genomes of soybean, *Medicago truncatula*, and *Arabidopsis thaliana*. *BMC Plant Biol*, 15.
- Mun, J.H., Yu, H.J., Park, S., and Park, B.S. (2009). Genome-wide identification of NBS-encoding resistance genes in *Brassica rapa*. *Mol Genet Genomics* 282, 617-631.
- Oliver, S.G. (2002). Functional genomics: Lessons from yeast. *Philosophical Transactions of the Royal Society of London B Biological Sciences* 357, 17-23.
- Poste, G. (2011). Bring on the biomarkers. *Nature* 469, 156-157.
- Ramachandra, S.B., Sathyanarayana, N.R., Subramonium, S., and Shetty, S.H. (2011). Isolation, cloning and characterization of resistance gene analogues in pearl millet based on conserved nucleotide-binding sites. *J Phytopathol* 159, 382-389.
- Ratnaparkhe, M.B., Wang, X., Li, J., Compton, R.O., Rainville, L.K., Lemke, C., Kim, C., Tang, H., and Paterson, A.H. (2011). Comparative analysis of peanut NBS-LRR gene clusters suggests evolutionary innovation among duplicated domains and erosion of gene microsynteny. *New Phytol* 192, 164-178.
- Regan, S., Gustafson, V., Mallubhotla, S., Chakravarty, B., Bagchi, M., Siahbazi, M., Rothwell, C., Sardana, R., Goyer, C., Audy, P., *et al.* (2006). Finding the perfect potato: using functional genomics to improve disease resistance and tuber quality traits. *Can J Plant Pathol* 28, S247-S255.
- Ribas, A.F., Cenci, A., Combes, M.C., Etienne, H., and Lashermes, P. (2011). Organization and molecular evolution of a disease-resistance gene cluster in coffee trees. *BMC Genomics* 12.
- Richly, E., Kurth, J., and Leister, D. (2002). Mode of amplification and reorganization of resistance genes during recent *Arabidopsis thaliana* evolution. *Mol Biol Evol* 19, 76-84.
- Rico, A., McCraw, S.L., and Preston, G.M. (2011). The metabolic interface between *Pseudomonas syringae* and plant cells. *Curr Opin Microbiol* 14, 31-38.
- Robertson, L.D., Singh, K.B., Erskine, W., and El Moneim, A.M.A. (1996). Useful genetic diversity in germplasm collections of food and forage legumes from West Asia and North Africa. *Genetic Resources and Crop Evolution* 43, 447-460.
- Rubiales, D., Fondevilla, S., Chen, W., Gentzbittel, L., Higgins, T.J.V., Castillejo, M.A., Singh, K.B., and Rispail, N. (2015). Achievements and Challenges in Legume Breeding for Pest and Disease Resistance. *Crit Rev Plant Sci* 34, 195-236.
- Sabir, J., Mutwakil, M., El-Hanafy, A., Al-Hejin, A., Abdel Sadek, M., Abou-Alsoud, M., Qureshi, M., Saini, K., and Ahmed, M. (2014). Applying molecular tools for improving livestock performance: From DNA markers to next generation sequencing technologies. *J Food, Agric Environ* 12, 541-553.
- Schwartz, D.A., Freedman, J.H., and Linney, E.A. (2004). Environmental genomics: a key to understanding biology, pathophysiology and disease. *Hum Mol Genet* 13, R217-R224.
- Sharma, R., Chhokar, V., Jha, S., and Rana, J.S. (2008). Proteomics: Current approaches and future perspectives. *Journal of Applied Bioscience* 34, 1-16.
- Sharma, S., Upadhyaya, H.D., Varshney, R.K., and Gowda, C.L.L. (2013). Pre-breeding for diversification of primary gene pool and genetic enhancement of grain legumes. *Front Plant Sci* 4.
- Sharma, T.R., Das, A., Thakur, S., and Jalali, B.L. (2014). Recent understanding on structure, function and evolution of plant disease resistance genes. *Proc Indian Natl Sci Acad* 80, 83-93.
- Singh, N.K., Gupta, D.K., Jayaswal, P.K., Mahato, A.K., Dutta, S., Singh, S., Bhutani, S., Dogra, V., Singh, B.P., Kumawat, G., *et al.* (2012). The first draft of the pigeonpea genome sequence. *J Plant Biochem Biotechnol* 21, 98-112.
- Singh, R.J., Chung, G.H., and Nelson, R.L. (2007). Landmark research in legumes. *Genome* 50, 525-537.
- Sladek, R., and Hudson, T.J. (2006). Elucidating cis- and trans-regulatory variation using genetical genomics. *Trends Genet* 22, 245-250.
- van Buijtenen, J.P. (2001). Genomics and quantitative genetics. *Can J For Res* 31, 617-622.
- Van Zee, J.P., Geraci, N.S., Guerrero, F.D., Wikel, S.K., Stuart, J.J., Nene, V.M., and Hill, C.A. (2007). Tick genomics: The *Ixodes* genome project and beyond. *Int J Parasitol* 37, 1297-1305.
- Varshney, R.K., Chen, W., Li, Y., Bharti, A.K., Saxena, R.K., Schlueter, J.A., Donoghue, M.T., Azam, S., Fan, G., Whaley, A.M., *et al.* (2012). Draft genome sequence of pigeonpea (*Cajanus cajan*), an orphan legume crop of resource-poor farmers. *Nat Biotechnol* 30, 83-89.
- Varshney, R.K., Song, C., Saxena, R.K., Azam, S., Yu, S., Sharpe, A.G., Cannon, S., Baek, J., Rosen, B.D., Tar'an, B., *et al.* (2013). Draft genome sequence of chickpea (*Cicer arietinum*) provides a resource for trait improvement. *Nat Biotechnol* 31, 240-246.
- Vo, T.D., and Palsson, B.O. (2007). Building the power house: recent advances in mitochondrial studies through proteomics and systems biology. *American Journal of Physiology Cell Physiology* 292, C164-C177.
- Vossen, J.H., Dezhsetan, S., Esselink, D., Arens, M., Sanz, M.J., Verweij, W., Verzaux, E., and van der Linden, C.G. (2013). Novel applications of motif-directed profiling to identify disease resistance genes in plants. *Plant Methods* 9.
- Webb, C.A., and Fellers, J.P. (2006). Cereal rust fungi genomics and the pursuit of virulence and avirulence factors. *FEMS Microbiol Lett* 264, 1-7.
- Wei, H., Li, W., Sun, X., Zhu, S., and Zhu, J. (2013). Systematic Analysis and Comparison of Nucleotide-

- Binding Site Disease Resistance Genes in a Diploid Cotton *Gossypium raimondii*. *PLoS One* 8.
- Xu, J.-R., and Xue, C. (2002). Time for a blast: Genomics of *Magnaporthe grisea*. *Mol Plant Pathol* 3, 173-176.
- Xu, Q.F., Cheng, W.S., Li, S.S., Li, W., Zhang, Z.X., Xu, Y.P., Zhou, X.P., and Cai, X.Z. (2012). Identification of genes required for Cf-dependent hypersensitive cell death by combined proteomic and RNA interfering analyses. *J Exp Bot* 63, 2421-2435.
- Xu, W., Yu, Y., Zhou, Q., Ding, J., Dai, L., Xie, X., Xu, Y., Zhang, C., and Wang, Y. (2011). Expression pattern, genomic structure, and promoter analysis of the gene encoding stilbene synthase from Chinese wild *Vitis pseudoreticulata*. *J Exp Bot* 62, 2745-2761.
- Young, N.D., Debelle, F., Oldroyd, G.E., Geurts, R., Cannon, S.B., Udvardi, M.K., Benedito, V.A., Mayer, K.F., Gouzy, J., Schoof, H., *et al.* (2011). The *Medicago* genome provides insight into the evolution of rhizobial symbioses. *Nature* 480, 520-524.
- Yu, J., Tehrim, S., Zhang, F., Tong, C., Huang, J., Cheng, X., Dong, C., Zhou, Y., Qin, R., Hua, W., *et al.* (2014). Genome-wide comparative analysis of NBS-encoding genes between *Brassica* species and *Arabidopsis thaliana*. *BMC Genomics* 15.
- Zimaro, T., Gottig, N., Garavaglia, B.S., Gehring, C., and Ottado, J. (2011). Unraveling plant responses to bacterial pathogens through proteomics. *J Biomed Biotechnol* 2011.




Other Publications of Interest

- Xu, J.-P. (2014). Next-generation Sequencing: Current Technologies and Applications. Caister Academic Press, U.K. ISBN: 978-1-908230-33-1
- He, Z. (2014). Microarrays: Current Technology, Innovations and Applications. Caister Academic Press, U.K. ISBN: 978-1-908230-49-2
- Poptsova, M.S. (2014). Genome Analysis: Current Procedures and Applications. Caister Academic Press, U.K. ISBN: 978-1-908230-29-4
- Fuentes, M. and LaBaer, J. (2014). Proteomics: Targeted Technology, Innovations and Applications. Caister Academic Press, U.K. ISBN: 978-1-908230-46-1
- Murillo, J., Vinatzer, B.A., Jackson, R.W. and Arnold, D.L. (2015). Bacteria-Plant Interactions: Advanced Research and Future Trends. Caister Academic Press, U.K. ISBN: 978-1-908230-58-4
- Caranta, C., Aranda, M.A., Tepfer, M. and Lopez-Moya, J.J. (2011). Recent Advances in Plant Virology. Caister Academic Press, U.K. ISBN: 978-1-904455-75-2
- Nannipieri, P., Pietramellara, G. and Renella, G. (2014). Omics in Soil Science. Caister Academic Press, U.K. ISBN: 978-1-908230-32-4
- Herold, K.E. and Rasooly, A. (2009). Lab-on-a-Chip Technology: Fabrication and Microfluidics. Caister Academic Press, U.K. ISBN: 978-1-904455-46-2



Further Reading

Caister Academic Press is a leading academic publisher of advanced texts in microbiology, molecular biology and medical research. Full details of all our publications at [caister.com](http://www.caister.com)

- **MALDI-TOF Mass Spectrometry in Microbiology**
Edited by: M Kostrzewa, S Schubert (2016)
www.caister.com/malдитof
- **Aspergillus and Penicillium in the Post-genomic Era**
Edited by: RP Vries, IB Gelber, MR Andersen (2016)
www.caister.com/aspergillus2
- **The Bacteriocins: Current Knowledge and Future Prospects**
Edited by: RL Dorit, SM Roy, MA Riley (2016)
www.caister.com/bacteriocins
- **Omics in Plant Disease Resistance**
Edited by: V Bhadauria (2016)
www.caister.com/opdr
- **Acidophiles: Life in Extremely Acidic Environments**
Edited by: R Quatrini, DB Johnson (2016)
www.caister.com/acidophiles
- **Climate Change and Microbial Ecology: Current Research and Future Trends**
Edited by: J Marxsen (2016)
www.caister.com/climate
- **Biofilms in Bioremediation: Current Research and Emerging Technologies**
Edited by: G Lear (2016)
www.caister.com/biorem
- **Microalgae: Current Research and Applications**
Edited by: MN Tsaloglou (2016)
www.caister.com/microalgae
- **Gas Plasma Sterilization in Microbiology: Theory, Applications, Pitfalls and New Perspectives**
Edited by: H Shintani, A Sakudo (2016)
www.caister.com/gasplasma
- **Virus Evolution: Current Research and Future Directions**
Edited by: SC Weaver, M Denison, M Roossinck, et al. (2016)
www.caister.com/virusevol
- **Arboviruses: Molecular Biology, Evolution and Control**
Edited by: N Vasilakis, DJ Gubler (2016)
www.caister.com/arbo
- **Shigella: Molecular and Cellular Biology**
Edited by: WD Picking, WL Picking (2016)
www.caister.com/shigella
- **Aquatic Biofilms: Ecology, Water Quality and Wastewater Treatment**
Edited by: AM Romani, H Guasch, MD Balaguer (2016)
www.caister.com/aquaticbiofilms
- **Alphaviruses: Current Biology**
Edited by: S Mahalingam, L Herrero, B Herring (2016)
www.caister.com/alpha
- **Thermophilic Microorganisms**
Edited by: F Li (2015)
www.caister.com/thermophile



- **Flow Cytometry in Microbiology: Technology and Applications**
Edited by: MG Wilkinson (2015)
www.caister.com/flow
- **Probiotics and Prebiotics: Current Research and Future Trends**
Edited by: K Venema, AP Carmo (2015)
www.caister.com/probiotics
- **Epigenetics: Current Research and Emerging Trends**
Edited by: BP Chadwick (2015)
www.caister.com/epigenetics2015
- **Corynebacterium glutamicum: From Systems Biology to Biotechnological Applications**
Edited by: A Burkovski (2015)
www.caister.com/cory2
- **Advanced Vaccine Research Methods for the Decade of Vaccines**
Edited by: F Bagnoli, R Rappuoli (2015)
www.caister.com/vaccines
- **Antifungals: From Genomics to Resistance and the Development of Novel Agents**
Edited by: AT Coste, P Vandeputte (2015)
www.caister.com/antifungals
- **Bacteria-Plant Interactions: Advanced Research and Future Trends**
Edited by: J Murillo, BA Vinatzer, RW Jackson, et al. (2015)
www.caister.com/bacteria-plant
- **Aeromonas**
Edited by: J Graf (2015)
www.caister.com/aeromonas
- **Antibiotics: Current Innovations and Future Trends**
Edited by: S Sánchez, AL Demain (2015)
www.caister.com/antibiotics
- **Leishmania: Current Biology and Control**
Edited by: S Adak, R Datta (2015)
www.caister.com/leish2
- **Acanthamoeba: Biology and Pathogenesis (2nd edition)**
Author: NA Khan (2015)
www.caister.com/acanthamoeba2
- **Microarrays: Current Technology, Innovations and Applications**
Edited by: Z He (2014)
www.caister.com/microarrays2
- **Metagenomics of the Microbial Nitrogen Cycle: Theory, Methods and Applications**
Edited by: D Marco (2014)
www.caister.com/n2



Order from caister.com/order