

***In vitro* Anti-inflammatory Study of Limonoids Isolated from *Chisocheton* Plants**

Erina Hilmayanti^{1,3}, Xuhao Huang¹, Supriatno Salam⁵, Nurlelasari³, Unang Supratman^{3,4}, Kazuya Kabayama^{1,2*}, and Koichi Fukase^{1,2,*}

¹ Department of Chemistry, Graduate School of Science, Osaka University, Toyonaka 560-0043, Japan

² Project Research Center for Fundamental Sciences, Osaka University, Toyonaka 560-0043, Japan

³ Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Padjadjaran, Jatinangor 45363, Indonesia

⁴ Central Laboratory, Universitas Padjadjaran, Jatinangor 45363, Indonesia

⁵ Faculty of Pharmacy, Universitas Mulawarman, Samarinda, Kalimantan Timur 75123, Indonesia

* Correspondence: koichi@chem.sci.osaka-u.ac.jp; Tel.: +81-6-6850-5388 (K.F.); kaba@chem.sci.osaka-u.ac.jp; Tel.: +81-6-6850-5192 (K.K.)

Contents

Figure S1. Agonist effects of 17 Limonoid compounds (**1-17**) at a concentration of 20 μ M on the NF- κ B activation in HEK-BlueTM hTLR4 Cells. All data from four independent experiments are expressed as mean \pm SD. ##p<0.0001 vs. culture medium-only control group; *p<0.01, **p<0.001, ***p<0.0001 vs. culture medium-only control group. Analyzed by one-way ANOVA, followed by Dunnett's test using GraphPad Prism 9.

Figure S2. Effects of dexamethasone, and compounds **3-5**, **9**, **11-12**, and **15-17** in concentrations of 1, 5, 10, and 20 μ M on the viability of THP-1 Cells presented as a heatmap.

Figure S3. Western blot result of compounds **3**, **4**, **5**, **9**, **12**, **15**, and **16** on MAPK activation (A: phosphorylated p38 MAPK; B: total p38 MAPK) in LPS-stimulated THP-1 cells.

Table S1. Cytotoxic effects of 17 tested compounds against RAW264.7 macrophage.




Figure S1. Agonist effects of 17 Limonoid compounds (1–17) at a concentration of 20 μ M on the NF- κ B activation in HEK-Blue™ hTLR4 Cells. All data from four independent experiments are expressed as mean \pm SD. ## p <0.0001 vs. culture medium-only control group; * p <0.01, ** p <0.001, *** p <0.0001 vs. culture medium-only control group. Analyzed by one-way ANOVA, followed by Dunnett's test using GraphPad Prism 9.




Figure S2. Effects of dexamethasone, and compounds 3-5, 9, 11-12, and 15-17 in concentrations of 1, 5, 10, and 20 μM on the viability of THP-1 Cells presented as a heatmap.

Table S1. Cytotoxic effects of 17 tested compounds against RAW264.7 macrophage.

Samples	Viability cell (%)		
	RAW 264.7		
	5 μ M	10 μ M	20 μ M
Control*	>99.90	>99.90	>99.90
Etoposide	67.53 \pm 2.36	54.79 \pm 2.58	38.85 \pm 3.19
1	>99.90	>99.90	>99.90
2	96.25 \pm 3.67	>99.90	>99.90
3	90.27 \pm 5.24	91.44 \pm 8.26	70.21 \pm 2.29
4	85.04 \pm 5.66	85.92 \pm 5.34	82.41 \pm 1.15
5	83.58 \pm 8.91	84.29 \pm 0.28	3.19 \pm 1.24
6	103.06 \pm 6.05	>99.90	96.66 \pm 2.14
7	>99.90	>99.90	95.88 \pm 2.19
8	89.47 \pm 9.38	94.75 \pm 0.32	87.46 \pm 3.33
9	93.76 \pm 1.86	89.94 \pm 2.44	79.41 \pm 10.16
10	93.19 \pm 4.20	82.06 \pm 1.31	86.14 \pm 4.07
11	90.10 \pm 7.88	79.02 \pm 1.58	78.88 \pm 2.21
12	>99.90	>99.90	89.39 \pm 6.90
13	>99.90	>99.90	91.89 \pm 4.37
14	>99.90	88.19 \pm 3.86	91.33 \pm 4.19
15	92.82 \pm 1.79	91.90 \pm 3.59	90.91 \pm 4.95
16	91.79 \pm 3.18	81.32 \pm 7.05	89.42 \pm 7.08
17	77.49 \pm 10.22	72.24 \pm 4.68	73.09 \pm 2.02

*Control: 1% EtOH in medium