current issues in molecular biology

MDPI

Article

Loss of Function of the *Zxdb* Gene Leads to a Decrease in the Decidualization Rate and Number of Pups Born in Mice by Affecting the Expression of the Cell Adhesion Molecules

Yafei Tian [1], Yang Zhang [1], Mengru Li [1], Rui Yin [1], Pingping Ding [1], Letong Liang [1], Bowen Chen [1], Rui Xu [1], Hongyan Chen [1], Chenming Xu [1], Songchang Chen [2,*] and Daru Lu [1,3,*]

[1] State Key Laboratory of Genetic Engineering, School of Life Sciences, Fudan University, Shanghai 200438, China; tianyf21@m.fudan.edu.cn (Y.T.); yang_zhang23@m.fudan.edu.cn (Y.Z.); mrli23@m.fudan.edu.cn (M.L.); ryin23@m.fudan.edu.cn (R.Y.); 20110700035@fudan.edu.cn (P.D.); 21110700032@m.fudan.edu.cn (L.L.); 22110700004@m.fudan.edu.cn (B.C.); 24110700107@m.fudan.edu.cn (R.X.); chenhy@fudan.edu.cn (H.C.); chenming_xu2006@163.com (C.X.)

[2] Shanghai Key Laboratory of Reproduction and Development, Institute of Reproduction and Development, Obstetrics and Gynecology Hospital, Fudan University, Shanghai 200011, China

[3] State Key Laboratory of Genetics and Development of Complex Phenotypes, School of Life Sciences and Institute of Medical Genetics and Genomics, Fudan University, Shanghai 200438, China

* Correspondence: chensongch@fudan.edu.cn (S.C.); drlu@163.com (D.L.)

Abstract

The Zinc Finger X-Linked Duplicate B (*ZXDB*) gene is one of a pair of replicated zinc finger genes on chromosome Xp11.21. The homologous gene of *ZXDB* in mice is *Zxdb*. Recent studies have found that *Zxdb* plays a role in the spermatogenic process of mice; however, its impact on the female reproductive system has not yet been explored. In our study, we found, for the first time, that the loss of function of *Zxdb* leads to reduced decidualization rates and a decrease in litter size in female mice. Secondly, we found that maternal loss of *Zxdb* is the determinant of these phenotypes. Thirdly, the transcriptional and proteomic differential expression genes in the uterine tissues of wild-type (WT) and *Zxdb* knockout (*Zxdb*-KO) mice were significantly enriched in signaling pathways such as adhesion molecules. Finally, we demonstrated that the disorder of expression and uneven distribution of adhesion molecules in mouse uterine tissue may be the main reason for the decline in embryo implantation rate. In conclusion, we have established for the first time a link between the *Zxdb* gene and reduced female fertility. This study will help provide guidance and genetic counseling for future common clinical complications such as Recurrent Spontaneous Abortion (RSA) or Recurrent Implantation Failure (RIF).

Keywords: *Zxdb*; decidualization; adhesion molecules; embryo implantation

Check for updates

Academic Editor: Cristina Manuela Dragoi

Received: 4 January 2026
Revised: 24 January 2026
Accepted: 26 January 2026
Published: 28 January 2026

Copyright: © 2026 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license.

1. Introduction

Embryo implantation is an early pregnancy event and a pivotal step in the mammalian reproductive process [1,2]. It refers to the process by which the blastocyst and the mother recognize each other, locate each other in the uterus, and adhere to and invade specific parts of the mother's uterus [3,4]. It is estimated that 50% of pregnancies result in miscarriage due to biochemical loss or implantation failure in the preclinical stage [5,6], and another 20% are clinically confirmed as pregnancy failure, mainly occurring in the first three months of pregnancy [7]. Successful implantation needs to occur during the WOI (window of implantation), so that the normally developing blastocyst can interact with the endometrium

in a receptive state for a short period of time [8]. At this time, endometrial epithelial cells express specific adhesion molecules to provide support for embryo attachment. If this WOI is missed, the embryo will not be able to implant.

Decidualization occurs during the secretory stage of the menstrual cycle and is controlled by progesterone and other ovarian hormones [9]. It is a prerequisite for successful embryo implantation [10,11] and results from complex interactions between hormones, transcription factors, morphogenetic factors, cytokines, cell cycle regulators, and signaling pathways [12]. In mice, there is a slight difference from humans; the decidualization process begins only after the embryo enters the uterus [9,13]. Decidualized cells change in cell morphology from slender fibroblast-like endometrial stromal cells to round or polygonal decidua cells. Decidua cells produce a large number of secreted proteins, including some marker proteins [14–16]. Among them, CDH1 (E-cadherin) and CDH2 (N-cadherin) are members of the cadherin family and play a crucial role in cell adhesion and signal transduction, especially during epithelial–mesenchymal transition (EMT) and mesenchymal–epithelial transition (MET) [15,17]. The ratio of CDH1/CDH2 is often used as a marker for MET/EMT, and the dynamic balance of MET/EMT is crucial for the success of pregnancy. Abnormal ratios may lead to implantation failure and recurrent miscarriage [18,19]. The localization and subsequent decidualization of endometrial vascular permeability responses suggest that they occur in response to interactions between the embryo and the endometrium [20,21]. Subsequently, the embryos begin to invade the endometrial tissue and are successfully implanted.

The *ZXDB* gene is one of a pair of replicated zinc finger genes on chromosome Xp11.21, the other being *ZXDA*. *ZXDB* has only one exon, with a GC content of approximately 70%, and encodes 803 amino acids. It is highly conserved in many species [22]. The *ZXDB* refers to the human gene; *Zxdb* is the murine ortholog. The predicted ZXDB protein contains 10 zinc finger tandem motifs that bind to DNA, and the protein's 576-703 amino acid sequence also has a transcriptional activation domain for gene expression. RT-PCR and transcriptome sequencing detected the expression of *ZXDB* in most human tissues, suggesting that it may play an important fundamental role in the development of organisms. Previous studies have found that *ZXDB* can affect the expression of MHCI and MHCII molecules [23,24].

In the mouse genome, there is a frameshift mutation in the *Zxda* gene, so the *Zxdb* gene is the only functional mouse homolog. A recent study has found that the FBXO38-dependent degradation of ZXDB is a key factor for achieving chromatin dynamic regulation, especially the centromere protein CENP-B. The ZXDA/B protein binds to the centromere protein CENP-B. Functionally, the ZXDA/B factor controls the level of CENP-B proteins associated with the chromatin. In testes, the *Fbxo38* gene is specifically expressed in Sertoli cells, while Sertoli cells lacking *Fbxo38* exhibit stable ZXDB protein and increased chromatin in the centromere region. After KO of the *Fbxo38* gene, the mice grow slowly and affect multiple organs, including the male reproductive system. Detailed analysis of the mutant testes revealed pathological changes in the spermatogenic tubules, along with a significant reduction in sperm production and decreased fertility [25].

Research in human cell lines or samples is currently very limited. Previous studies regarded *ZXDB* and *ZXDA* as similar proteins for investigation. Currently, there is insufficient research specifically targeting the *ZXDB* gene, and its important role as a transcription factor, especially in the reproductive system, remains to be explored.

Here we have, for the first time, discovered that knockdown or knockout of the *Zxdb* gene can lead to decidualization and a decrease in the litter size. Sequencing analysis of pre-implantation uterine tissues, conducted through transcriptomics and proteomics, revealed that significantly enriched pathways included those related to adhesion molecules,

immune responses, and other relevant pathways. WB experiments confirmed that the expression of adhesion molecules such as CDH1(E-cadherin), CDH2(N-cadherin), and BMP2 was disrupted. Immunofluorescence experiments revealed abnormal expression and distribution of CDH1 in the uterus, resulting in a decreased implantation rate. This study is conducive to deepening the analysis of RSA factors and genetic counseling by scientific researchers and clinicians.

2. Materials and Methods

2.1. Cell Line and Cell Culture

The Neuro-2a(N2a) cell line was purchased from the Cell Bank of the Chinese Academy of Sciences. Cell culture was performed using DMEM (GIBCO, 10566016, Waltham, MA, USA) medium supplemented with 10% high-quality fetal bovine serum and 1% penicillin–streptomycin solution (Beyotime, C0222, Shanghai, China). Cells were cultured in an incubator at 5% CO_2 and 37 °C. The siRNA transfection process was carried out according to the instructions provided in the Lipofectamine RNAi MAX kit (Invitrogen, 13778075, Thermo Fisher Scientific, Waltham, MA, USA). siRNA (*Zxdb*) sense sequence: 5′-GAGCUUCACCACCGUCUACAA-3′; siRNA (*Zxdb*) antisense sequence: 5′-GUAGACGGUGGUGAAGCUCUU-3′. The use of 2′-O-methyl (2′-OMe) modification for siRNA is to enhance its stability. All siRNAs were ordered from Shanghai GENEray Bioengineering Co., LTD. (Shanghai, China).

2.2. Mouse Breeding and Ethics

Both C57BL/6J and ICR mice involved in this laboratory were purchased from Shanghai JieSiJie Laboratory Animal Co., Ltd. (Shanghai, China). The female and male ICR mice were 6 to 8 weeks old, the C57BL/6J female mice were 4 to 6 weeks old, and the C57BL/6J male mice were over 8 weeks old. All the mice in this experiment were of SPF grade. The experimental process strictly adhered to the regulations of the Animal Ethics Committee of the School of Life Sciences, Fudan University. The content of the animal experiments has obtained ethical approval, with the approval number: No. 2022JS071.

2.3. Embryo Acquisition and Embryo Culture

In this study, fertilized eggs were obtained through in vivo/in vitro fertilization. Each female mouse was injected with 5 to 10 IU of PMSG (Nanjing Aibei Biotechnology Co., Ltd., M2620, Nanjing, China). After 48 h, 5 to 10 IU of HCG (Nanjing Aibei Biotechnology Co., Ltd., M2530, Nanjing, China) was injected. The mice were mated in one cage, and the vaginal plugs were checked the next morning. For mice with vaginal plugs, 14 h after HCG injection, they were sacrificed. The fallopian tubes at the junction of the uterus and ovary were removed through an opening on the back. Under a stereomicroscope, oocytes were drawn out in M2 buffer (Nanjing Aibei Biotechnology Co., Ltd., M1250, Nanjing, China) with a 1 mL syringe, and the zona pellucida was removed using hyaluronidase (Nanjing Aibei Biotechnology Co., Ltd., M2215, Nanjing, China). Transfer clean eggs or embryos to KSOM medium (MERCK, MR-101, Darmstadt, Germany). The culture medium needs to be prepared in advance and balanced overnight. The morphological observation of the embryo is carried out according to the changes in the division time. Generally, it is checked once a day until it reaches the blastocyst stage.

2.4. siRNA Injection Embryo and Uterine Horn Interference Experiment

siRNA embryo injection was carried out by microinjection. When operating, first place the obtained Cumulus Oocyte Complexes (COCs) in hyaluronidase (Nanjing Aibei Biotechnology Co., Ltd., M2215, Nanjing, China) for digestion to remove the outer granular cells.

Embryos with granulosa cells removed were washed and transferred to KSOM medium for further culture. Microinjection was performed when the embryos were cultured to the prokaryotic stage. Subsequently, a 20 μM siRNA solution that had been pre-centrifuged was directly injected into the cytoplasm or prokaryote of the fertilized egg using a microscopic operating system and a glass capillary needle. The injection volume was approximately 2 to 5 pl per fertilized egg. After the injection was completed, the embryos were immediately transferred into KSOM culture medium and cultured at 37 °C and 5% CO_2. For the RNA interference experiments on uterine tissue, the abdominal incision exposed the uterine horn after the female mice were anesthetized. A fine needle was used to directly inject the siRNA solution into the uterine cavity, avoiding penetration of the uterine wall. After injection, the abdominal cavity was sutured, and samples were collected 7~8 days after recovery.

2.5. Construction of the Zxdb-KO Mouse Model

The *Zxdb*-KO mouse model was constructed by our research team using a dual sgRNA protocol. Guide RNA sequence1: 5′-ACGCCGAGAUGGAAAUCCCG-3′; Guide RNA sequence2: 5′-GGGACCGCAGACAGACAGCG-3′; sgRNAs were prepared by in vitro transcription using the T7 High Yield RNA Transcription kit (Novoprotein, E131, Suzhou, China). Then, superovulation was performed on the female mice, and they mated with the male mice to obtain fertilized eggs. The Ribonucleoprotein complex (200 ng/μLCas9 protein + 100 ng/μL sgRNA) was injected into the prokaryote or cytoplasm of the fertilized egg by microinjection. After injection, the embryo was transplanted into the uterus of the surrogate mother mouse. After the birth of the pups, genotypes were identified through PCR, sequencing, and other methods to screen for positive F0 generation (Founder mouse). Finally, the positive mice were expanded, and a stable genetic model was obtained.

2.6. Mouse Genotype Identification

Seven days after the birth of the mice, their nails were cut to extract genomic DNA for PCR genotype identification. The first genotype identification experiment identified the specific genotype of the mutant through PCR. There is a 528 bp deletion between sgRNA1 and sgRNA2 target sites. Subsequently, primers were designed across breakpoints, and the genotype was directly determined based on the size of the electrophoresis bands. Primer 1F: 5′-GCCTCACCGACGCCGAGATGGAAATCTGTCT-3′; Primer 2F: 5′-CTTGACCCGGTGGGTGGCGATGTAGAAACC-3′. Universal primers Primer1 and 2R: 5′-GTGGTCAGGGTGAGGACGCCGTTCTCGAA-3′. The breakpoint product has a KO band of 187 bp and a WT band size of 308 bp. Both the homozygous and WT mice have only one band, while the heterozygous has two bands.

2.7. Transcriptome and Proteomics Sequencing Analysis

Uterine tissue of WT and *Zxdb*-KO mice was collected with vaginal plugs on the third day. Transcriptome and proteome sequencing were carried out in Tsingke Biotechnology Co., Ltd. (Beijing, China). For the sequencing results of transcriptome data, firstly, conduct quality control and filtering on the original sequencing data; subsequently, high-quality sequences are aligned to the reference genome or transcriptome. Next, quantitative analysis of gene expression levels (such as counting reads) is carried out, and on this basis, differential expression analysis is conducted to screen for genes with significant changes among different groups. Finally, enrichment analysis of the differentially expressed genes was conducted in combination with functional annotation databases to reveal the biological functions, pathways, or regulatory networks they are involved in. The DESeq2 (1.45.0) was used for significance analysis of differences in the project. A p-value of less than 0.05 and a ratio of expression multiples greater than or equal to 2 ($\log2FC \geq 1$) were taken as the significance criteria for differences. Proteomics employs Ultra-Fast technology. This

analytical method is based on high-throughput mass spectrometry technology and combines ultrafast liquid chromatography with high-resolution mass spectrometry platforms (such as Orbitrap Astral) to achieve rapid separation and detection of complex protein samples. Through efficient database search (such as MaxQuant, FragPipe) and bioinformatics analysis, the system systematically analyzes differentially expressed proteins and their functions.

2.8. Trypan Blue Dyeing

One percent trypan blue dye (Beyotime, C1313M, Shanghai, China) was injected into the tail vein of WT and *Zxdb*-KO mice on the 7th day of pregnancy for in vivo assessment of the rate of decidualization.

2.9. Hematoxylin–Eosin Staining

Hematoxylin–Eosin (H&E) staining was carried out in accordance with the H&E staining kit instructions (Beyotime, C0105M, Shanghai, China). Paraffin sections of uterine tissue were subjected to gradient dewaxing with xylene and anhydrous ethanol, followed by water washing. The nuclei were stained with hematoxylin for 4 to 10 min and differentiated to turn blue, and the cytoplasm was stained with eosin for 3 to 5 min. Then, the sections were dehydrated with gradient ethanol, made transparent with xylene, sealed with neutral gum, and finally photographed under the microscope (Olympus, IX73, Tokyo, Japan) for observation and analysis.

2.10. Immunofluorescence Staining

Mouse uterine tissue was embedded in OCT (SAKURA, 4583, Torrance, CA, USA), rapidly frozen in liquid nitrogen, and then frozen into sections with a thickness of 7~10 μm. The sections were washed with PBS, dried at 37 °C, and sealed with 5% BSA and goat serum (Solarbio, SL038, Beijing, China). Subsequently, the primary antibody was incubated overnight in a wet box at 4 °C. On the second day, the corresponding fluorescent secondary antibodies (Abcam Goat Anti-Rabbit IgG H&L Alexa Fluor® 594, ab150080, Shanghai, China) and DAPI (Beyotime, P0131, Shanghai, China) were incubated after the runs on the slides. After PBS washing, the plates were encapsulated with anti-quenching mounting medium and finally observed, and images were collected under a fluorescence microscope.

2.11. Quantitative PCR

RNA was extracted using the traditional TRIzol (Invitrogen, 15596018CN, Carlsbad, CA, USA) method. Reverse transcription of 1 μg of RNA. Reverse transcription and genome removal were performed using Novozyme HiScript III RT SuperMix for qPCR Kit (Vazyme, R323-01, Nanjing, China). Dilute the cDNA and perform quantitative PCR using tSYBR® Green Realtime PCR Master Mix (TOYOBO, QPK-201, Osaka, Japan).

2.12. Western Blot (WB)

The WB experiment was conducted according to the classic wet-film transfer method. The antibodies involved in this study include β-actin (Servicebio, GB11001-100, Wuhan, China), CDH1 (Zenbio, 340341, Chengdu, China), CDH2 (Shanbenbio, 1206, Hefei, China), and BMP2 (Shanbenbio, 52878, Hefei, China).

2.13. Statistical Analysis

All statistical analysis and data visualization were performed using Prism 9 software. Data are expressed as the mean ± standard deviation (mean ± SD). Analysis of variance (ANOVA) was used for inter-group comparisons. The *t*-test is used to compare the differences in the means between two groups. The drawing software uses Biorender and

CorelDRAW 2025. ns, no significant difference; statistical significance was expressed as * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$.

3. Results

3.1. Expression Analysis of Zxdb in the Reproductive System of Female Mice

To explore the expression of *Zxdb* in the early stage of the embryo, we re-analyzed the transcriptome and proteome data of mouse embryos from Shaorong Gao's Data [26]. The results from Figure 1A show that the mRNA level of *Zxdb* was not present in all embryos during the single-cell zygote stage. No mRNA expression of *Zxdb* was detected at the two-cell, four-cell, or eight-cell stage. At the morula and blastocyst stages, only some embryos expressed *Zxdb* mRNA. Overall, *Zxdb* mRNA was at a very low or absent expression level. As an internal reference gene for the early embryonic development, the Pipa gene is often used to standardize embryonic RT-qPCR or RNA-seq data to ensure the accuracy and comparability of gene expression analysis at different development stages [27–29]. Figure 1B shows that as the embryo develops and the number of cells increases, the mRNA of the *Pipa* gene shows a good, gradual increase. After homogenization, it was found that *Zxdb* was barely detectable in the two-cell, four-cell, eight-cell, morula, and blastocyst stages. To better understand the expression of *Zxdb*, we re-analyzed the proteomic data of embryos. Figure 1D shows that PIPA protein expression level steadily increased as the embryo developed from zygote to blastocyst, while the ZXDB protein was not detected at any stage of embryonic development, which also suggests that *Zxdb* may not play a key role in early embryonic development.




Figure 1. The expression of *Zxdb* in embryos and uterine tissue at different time points during the early pregnancy period. (**A**–**C**) The mRNA expression of, respectively, *Zxdb*, *Pipa*, and *Zxdb/Pipa* in zygote, two-cell, four-cell, eight-cell, morula, and blastocyst stages, with a sample size of $n = 3$. Figure (**D**) represents the protein expression level of the *Zxdb* gene in fertilized eggs, two-cell, four-cell, eight-cell, morula, and blastocyst cells. $n = 3$. (**E**) The H&E staining results of the uterine tissue. (**F**) Expression changes of *Zxdb* in the uterine tissue of mice on E1.5, E3.5, E5.5, and E7.5 of pregnancy. $n = 3$. E1.5, E3.5, E5.5, and E7.5, respectively, represent the 1.5, 3.5, 5.5, and 7.5 days when female mice have vaginal plugs. The day there is a vaginal plug is called E0.5.

In addition to observing the expression of the *Zxdb* gene in embryos, we also investigated the expression of *Zxdb* in the uterine tissue of mice. The uterine tissues of E1.5, E3.5, E5.5, and E7.5 mice were selected. As can be seen from the H&E staining results (Figure 1E), with the secretion of HCG hormone and others, the uterine cavity of the mice began to close from E1.5 to E7.5. When the embryo entered the uterine cavity on day 4.5, there was still some space in the uterus. Once the embryo is implanted, the uterine cavity will close rapidly. The fluid is discharged, and then the embryo invades the mother's uterine tissue. Around the 7th day, placental blood vessels begin to form (Figure 1E). The quantitative PCR results (Figure 1F) show that the expression of *Zxdb* gradually increased with the increase in gestational age. In both humans and mice, the majority of embryo loss occurs in the early stage of pregnancy [30,31]. The fertilized egg of a mouse usually begins to enter the uterus and attach to the endometrium after fertilization. A receptive endometrium plays a crucial role in the embryonic development of a mouse. The continuously enhanced expression of *Zxdb* in the uterine tissue of mice after pregnancy may have a certain effect on the receptivity of the endometrium and maintain the stable implantation of the embryo.

3.2. Decreased Expression of Zxdb by Maternal Factors Leads to Reduced Mouse Embryo Implantation

To further study whether knockdown of *Zxdb* in uterine tissue in pregnant mice affected embryo implantation, we injected stably modified siRNA directly into the mouse uterine tissue before embryo implantation on day 3 of pregnancy, and the validation of knockdown efficiency was conducted in N2a cells, with an efficiency of approximately 70%. As shown in Figure 2A, mice aged about 6 weeks were treated with PMSG, injected with HCG 48 h later, and then they mated with male mice in cages. The next morning, the vaginal plug was checked. On the third day of embryonic development, the uterus of mice was pulled out, and the uterine horns of the mice were subjected to injection experiments. In the negative control group, the left side served as a self-control, and the right side was injected with normal saline; in the si-*Zxdb* group, a negative control was injected on the left side as a self-control, and si-*Zxdb* was injected on the right side (Figure 2B). When the embryo developed to about 7.5 days, the mice were sacrificed. As shown in Figure 2B,C, there was no significant difference in the number of embryos implanted in the left uterus and the right uterus in the negative control group. In the si-*Zxdb* group, there was a significant difference in the number of embryos implanted in the left uterus and the right uterus. This result shows that knocking down the expression of *Zxdb* in mouse endometrial tissue reduces the decidualization rate of the mice.

Although from the perspective of transcriptomics and proteomics, the expression of *Zxdb* in early embryos is very little or non-existent, the effect of this minimal *Zxdb* expression on embryonic development in mouse embryos remains unclear. To this end, we conducted siRNA knockdown *Zxdb* experiments on mouse embryos of two strains: C57BL/6J and ICR. Figure 2D shows the results of embryonic development in ICR mice. The results showed that compared with the mock group and negative control group, there were no significant differences in the two-cell, four-cell, eight-cell, morula, and blastocyst formation rates of the injected embryos in the si-*Zxdb* group (Figure 2E). The corresponding mean values of two-cell, four-cell, right-cell, morula, and blastocyst formation rates in the mock group were 98 ± 4.47% (n = 5), 86.8 ± 9.57% (n = 5), 82.8 ± 13.8% (n = 5), 84.8 ± 12.07% (n = 5), and 81.3 ± 11.23% (n = 5), respectively. The corresponding mean values for two-cell (n = 5), four-cell (n = 5), eight-cell (n = 5), morula (n = 5) and blastocyst (n = 5) formation rates in the negative group were 98.18 ± 2.58% (n = 5), 92.15 ± 7.72% (n = 5), 89.65 ± 6.41% (n = 5), 86.89 ± 7.12% (n = 5) and 79.96 ± 8.04% (n = 5), respectively. The corresponding mean values for two-cell, four-cell, eight-cell, morula, and blastocyst formation rates in the si-*Zxdb* group were 96.54 ± 5.23% (n = 5), 92.09 ± 6.81% (n = 5),

85.97 ± 4.60% (n = 5), 84.86 ± 5.86% (n = 5), and 82.35 ± 5.04% (n = 5), respectively. For pictures of different developmental stages, please refer to the Supplement Figure S1. The results of interference in C57BL/6J mouse embryos were consistent with those in ICR mouse embryos.




Figure 2. Schematic diagram of siRNA injection into the uterine tissue and siRNA injection into embryos. (**A**) Schematic diagram of uterine injection. (**B**) Embryos removed from the mouse uterus. (**C**) Statistical results of the number of implanted embryos in the two groups. Two-Way ANOVA, n = 4, ns means not significant; ** $p < 0.01$. (**D**) The picture shows the pronuclear-stage embryo on day 1 and the blastocyst stage on day 5 of ICR mouse embryos. (**E**) summarizes the success rates of embryo development at the two-cell, four-cell, eight-cell, morula, and blastocyst stages in the si-*Zxdb*, negative control, and MOCK groups. Two-Way ANOVA, n = 5, ns means not significant.

3.3. Construction and Identification of Zxdb-KO Mice

To further investigate the effect of *Zxdb*, our team constructed the *Zxdb*-KO mouse genetic model. Figure 3A shows the process of gene-edited mice. Figure 3B shows a schematic diagram of two sgRNA targeted editing targeting the *Zxdb* gene. In addition to retaining some embryos to develop into blastocysts for editing efficiency testing, all the remaining two-cell embryos were transplanted into surrogate mother mice through the opening at the front end of the fallopian tube. After multiple experiments, a total of four mice were produced (Figure 3C). After genotype identification, a 528 bp deletion *Zxdb*-KO

male mouse was screened out (Figure 3D). The first genotype identification experiment identified the specific genotype of the mutant through PCR. Subsequently, as shown in Figure 3E, the design diagram of the primers was presented. Primers were designed across breakpoints, and the genotype was directly determined based on the size of the electrophoresis bands. The breakpoint product has a KO band of 187 bp and a WT band size of 308 bp (Figure 3F). As CRISPR/Cas9 technology has a certain possibility of being off-target, in order to detect the off-target efficiency, PCR was performed on the sites that the software predicted might be off-target, and first-generation sequencing verification was carried out. The verification results show that none of the possible off-target sites were edited, and there was no off-target effect.




Figure 3. Double sgRNA knockout of the *Zxdb* gene and genotype identification. (**A**) Gene editing production process. The small picture shows pronucleus injection; (**B**) schematic diagram of double sgRNA knockout *Zxdb* gene design; (**C**) 528 bp deletion of *Zxdb* gene knocked out and first-generation sequencing results of WT mice; (**D**) interpretation of identification results; (**E**) primer design idea; (**F**) chimeric mice born to surrogate mother mice. KO means homozygous knockout, WT means wild-type, and Het means heterozygote.

3.4. The Loss-Function of the Zxdb Gene Influences Female Reproductive Phenotype in Mice

To examine whether the phenotypes of genotype knockout mice were consistent with the results of previous RNA interference models, we conducted experiments on homozygous *Zxdb*-KO mice and WT mice. Figure 4A shows the number of embryos implanted in the uterus of WT and *Zxdb*-KO mice. The results showed (Figure 4B) that the average number of *Zxdb*-KO embryos was 9 ($n = 9$), which was significantly lower than the number of embryo implants in WT mice, which was 15 ($n = 9$), $p < 0.0001$. In addition, we conducted a statistic on the number of newborns (Figure 4C). The statistics showed that the average number of offspring born in WT mice was 14 ($n = 7$), and that in the KO mouse group was 10 ($n = 7$), $p < 0.01$. In conclusion, both the number of embryo implants and the number of offspring were significantly reduced after *Zxdb* knockout.




Figure 4. Statistics on decidualization and pup numbers. (**A**) The number of embryo implants in WT and *Zxdb*-KO mice and the trypan blue staining results, and the position indicated by the arrow represent the embryo that has implanted. (**B**) statistics on the number of embryo implants; (**C**) statistics of the number of newborns. *t*-test; **** $p < 0.0001$; ** $p < 0.01$.

3.5. Characterizing Differentially Expressed Genes (DEGs) by Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) Pathway Analysis

To study the effects of *Zxdb* knockout on the reproduction of female mice, especially the overall impact on embryo implantation. We performed transcriptomic sequencing on WT and *Zxdb*-KO mice. The results of principal component analysis (PCA) showed that the *Zxdb*-KO group and the WT group clustered well, indicating that the expression patterns of genes in the samples of the same group were similar (Figure 5A). The volcano plot (Figure 5B) shows that compared with WT, the *Zxdb*-KO group had a total of 3305 DEGs, among which 1951 DEGs were upregulated and 1354 DEGs were downregulated. The knockout sequencing results showed that after knocking out *Zxdb*, the expression of *Zxdb* was lost (Figure 5C), which was consistent with the genotype identification results.




Figure 5. Transcriptome sequencing analysis and pathway enrichment analysis of mouse uterine tissue. (**A**) PCA analysis; (**B**) volcano map; (**C**) the expression of *Zxdb* in RNA-seq data, *t*-test; ****, $p < 0.0001$; (**D**) GO enrichment analysis; (**E**) KEGG enrichment analysis; (**F**) heat map of expression of decidualization and implantation-related adhesion molecules.

According to gene ontology (GO) analysis, DEGs in *Zxdb*-KO vs WT mouse transcriptome can be categorized into 14 major biological processes and cellular components (Figure 5D): extracellular region (18.2%), extracellular exosome (14.5%), cell adhesion (9.8%), extracellular space (8.7%), cell surface (6.2%), plasma membrane (5.9%), extracellular matrix (5.4%), collagen-containing extracellular matrix (4.8%), basement membrane (4.1%), apical plasma membrane (3.7%), extracellular matrix structural constituent (3.5%), heparin binding (3.2%), calcium ion binding (2.9%), signaling receptor binding (2.6%), and other processes (approximately 26% combined for remaining terms) (Figure 5D). Based on the hypergeometric test, 12 of these terms were significantly enriched for differentially expressed genes, including extracellular region ($p = 1.33 \times 10^{-42}$), extracellular space ($p = 1.33 \times 10^{-42}$), extracellular exosome ($p = 2.52 \times 10^{-32}$), cell adhesion ($p = 1.84 \times 10^{-35}$), heparin binding ($p = 1.80 \times 10^{-32}$), extracellular matrix ($p \approx 10^{-20}$), collagen-containing extracellular matrix ($p \approx 10^{-19}$), cell surface ($p \approx 10^{-16}$), plasma membrane ($p \approx 10^{-13}$), basement membrane ($p \approx 10^{-12}$), apical plasma membrane ($p \approx 10^{-10}$), and extracellular matrix structural constituent ($p \approx 10^{-8}$). Similar to the results of the GO analysis, the KEGG plot of *Zxdb*-KO mice showed that the adhesive-related pathways were also disordered (Figure 5E): the focal adhesion and cell adhesion molecules were downregulated, weakening the adhesion between the embryo and the uterus. Upregulation of ECM–receptor interaction may lead to abnormal ECM remodeling and hinder trophoblast invasion. Al-

though the upregulation of *PI3K-Akt*, *Rap1*, and calcium signaling promotes proliferation, the out-of-control timing may cause inflammation or decidual differentiation disorders, disrupt the receptivity of the implantation window, and ultimately lead to insufficient decidual support, increased embryo absorption, and decreased litter size.

The heat map of *Zxdb*-KO mice shows dysregulation of key genes for implantation (Figure 5F): downregulation of *Cdh1*, *Itga9*, *Cdh2*, *Col26a1*, and *Cxcl14* weakens embryonic adhesion and ECM remodeling. Downregulation of *Mmp2* inhibits invasion. Downregulation of *Notch1*, *Ihh*, and *Hand2* blocks decidualization and differentiation. Although the upregulation of *Bmp2*, *Ptgs2*, and *Igf1* compensates, the disordered sequence causes inflammation. Overall disruption of uterine receptivity leads to implantation failure, insufficient decidual support, increased embryo absorption, and a decrease in the birth rate. These data indicate that *Zxdb* knockout might invoke a variety of genes participating in a wide range of extracellular matrix and cell adhesion-related biological processes.

3.6. Integrative Transcriptome and Proteome Analysis of Zxdb-KO Mice

To gain a clearer understanding of the impact of *Zxdb* being knocked out on the reproduction of female mice, we also conducted proteomic analysis. The proteomics results of PCA analysis showed that the *Zxdb*-KO group and the WT group clustered well, indicating that the expression patterns of genes in the samples of the same group were similar (Figure 6A). The volcano diagram (Figure 6B) shows that there are 253 upregulated proteins (red dots), 297 downregulated proteins (green dots), and 11,110 proteins with no significant change (gray dots). It was indicated that after *Zxdb* knockout, the expression levels of approximately 550 proteins changed significantly, suggesting that *Zxdb* knockout has a considerable impact on protein expression in mouse cells/tissues. Figure 6C shows the HeatScatter Plot of the expression levels of genes jointly identified by each sample group at the transcriptional level (genes) and the translation level (proteins), which can visually reflect the density distribution and correlation results of protein and gene expression levels. Control group ($\rho = 0.33$): There was a moderate positive correlation between gene expression and protein expression. KO group ($\rho = 0.322$): After knocking out *Zxdb*, the correlation coefficient slightly decreased but did not change much, still showing a moderate positive correlation. The correlation coefficients were close and relatively low, indicating that the overall coupling of gene–protein expression was weak in both groups. After the *Zxdb* knockout, there was no significant change in the gene–protein correlation, indicating that the knockout did not significantly disrupt the overall correlation, but it might affect the transcription–translation coupling on specific genes. The Venn diagram (Figure 6D) shows the overlapping relationships of all genes, diff genes, diff proteins, and all proteins. The number of differentially expressed genes was far greater than that of differentially expressed proteins (5194 genes vs 1206 genes), indicating that knockout mainly affects the transcriptional level, and transcriptional changes are not completely conducted to the protein level (possibly due to translation regulation or protein stability). The small overlap (105 genes) indicates that many differentially expressed genes did not cause protein changes, suggesting that transcriptional changes were not fully conducted to the proteins, which may involve post-transcriptional regulation. The KEGG enrichment bubble plot (Figure 6E) shows that in *Zxdb*-KO mice, the proteomes of "focal adhesion" and "ECM–receptor interaction" are significantly enriched (low *p*-value, Rich Factor 0.2~0.3, and high count); "cell adhesion molecules" are mainly characterized by transcriptome changes. The proteome shows greater significance in the focal adhesion and ECM interaction pathways, suggesting that post-translational regulation after *Zxdb* knockout enhances the alteration of adhesion signals, which may change the proliferation and invasion of uterine stromal cells.




Figure 6. Comprehensive analysis of transcriptomics and proteomics. (**A**) PCA analysis. (**B**) volcano map. (**C**) HeatScatter Plot, with the WT group on the left and the *Zxdb*-KO group on the right. X-axis: Log2 (protein expression)—the logarithmic transformation of protein expression levels. Y-axis: Log2 (gene expression)—the logarithmic transformation of gene expression levels. Each point represents a gene/protein pair, and colors ranging from blue to red indicate density (blue for low density and red for high density). (**D**) Venn plot—overlap analysis of differentially expressed genes/proteins. (**E**) Combined enrichment of the transcriptome and proteome.

3.7. Expression Changes of Decidualization Marker Genes and Adhesion Factor-Related Genes

To observe differences in uterine tissue before embryo implantation, uterine tissue was obtained from WT and *Zxdb*-KO mice on day 3 of gestation. Comparison of H&E staining of uterine implantation sites showed that the decidua layer in the WT group was thick, uniform, and dense, with rich cells, forming a full and regular ring structure surrounding the embryonic cavity. In the *Zxdb*-KO group, the decidua became significantly thinner, sparse, and uneven, with loose structure, increased edge folds, and flat implantation sites (Figure 7A). These histological differences reflect that *Zxdb* knockout leads to decidualization defects, insufficient differentiation of maternal decidual cells, an inability to fully

support embryo implantation and development, and may ultimately show a significant reduction in litter rate. Based on the enrichment results of differentially expressed genes in the transcriptome and proteome, we conducted WB verification on some crucial adhesion molecules (CDH1, CDH2, and BMP2) of decidualization and embryo implantation. Compared to the WT group, the expression of CDH1 and CDH2 proteins decreased, and the expression of BMP2 protein was upregulated. The expressions of BMP2, CDH1, and CDH2 were disordered compared to the WT, which might lead to abnormalities during embryo implantation and decidualization, resulting in phenotypes such as miscarriage or low fertility. During embryo implantation, normal "cadherin switching" on the uterine side requires downregulation of CDH1 and upregulation of CDH2. The CDH1/CDH2 ratio increases in our result (Figure 7F); this means that the endometrial receptivity window was not open normally, or the embryo develops abnormally, which would destroy the key steps of embryo adhesion and invasion, thus having a significant negative impact on embryo implantation. Immunofluorescence showed that the expression intensity of CDH1 was lower compared with that in the WT (Figure 8). The enlarged image confirmed that the expression network was damaged and its continuity was disrupted. This spatiotemporal disorder eventually leads to the inability of embryos to attach and implant normally, resulting in a significant decrease in the implantation rate.




Figure 7. The H&E staining results and verification of WB results. (**A**) The H&E staining results of uterine tissue; (**B**) WB experimental results; (**C**–**E**) statistical results of BMP2/CDH1/CDH2 expression; (**F**) statistical results of CDH1 expression. Statistical significance was expressed as * $p < 0.05$; *** $p < 0.001$.




Figure 8. Immunofluorescence staining of CDH1 protein. Uterine tissues were immunostained with anti-CDH1 (red) and DAPI (blue). Merge represents the combined result of CDH1 and DAPI; Enlarged represents the staining result after magnification.

4. Discussion

The process of embryo implantation and decidualization involves the synergistic effect of a large number of genes. Previous studies, through genetic screening of families with recurrent spontaneous abortion, have identified some key genes. These genes related to recurrent miscarriage include *HOXA10* [32,33], *FOXO1* [34,35], *CDH1* [32,36], *LIF* [37], *MUC1* [38], and *BMP2* [39] genes. However, even so, a large number of unexplained recurrent spontaneous abortion (URSA) cases still have no identifiable cause [40]. Perhaps there are more genes involved, especially the transcription factors that regulate differentiation and development.

Progesterone receptor (PR) mediates progesterone signaling, promoting the differentiation of endometrial stromal cells into decidua cells. Transcription factors are crucial in this process [12]. The key transcription factors include *CEBPB*, *HOXA10*, *FOXO1*, and *ZEB1*, which regulate cell proliferation, differentiation, and the expression of markers, ensuring the establishment of uterine receptivity [14]. *CEBPB* promotes the differentiation of human embryonic stem cells (hESCs), and decidualization does not occur in the mouse uterus without *CEBPB* [41,42]. HCG improves endometrial receptivity by increasing the expression of *HOXA10* [43]. *FOXO1* competes with *PGR* to control the expression of a subset of genes involved in decidualization [44]. *ZEB1* affects embryo implantation by regulating decidualization [45]. Abnormalities in these genes can lead to decidualization defects, implantation failure, or pregnancy complications. *ZXDB* is a zinc finger transcription factor. Here, we explored the siRNA uterine injection model and the *Zxdb*-KO genetic model to study the role of *Zxdb* in the female reproductive system of mice. We found that it plays an important role in decidualization and embryo implantation.

The expression of the *Zxdb* gene in mouse embryos and mouse uterus tissues was observed in our study. siRNA interference experiments on embryos confirmed that knockdown of *Zxdb* did not affect embryo development in the early stages of embryonic development (zygotic stage to blastocyst stage). By knocking down *Zxdb* in uterine tissue, we confirmed that the reduction in expression of *Zxdb*, which is a maternal factor, affects embryo implantation. However, there may be a post-implantation time period, such as from post-implantation to the fetal heart beating stage. Whether it has an impact on the development of the embryo still requires further research. In this study, the use of inbred strain ICR mice to construct the genetic model was based on their strong reproductive ability, which is highly suitable for conducting research on reproductive-related topics.

However, in order to eliminate the influence of different strains, we also performed RNA interference of the *Zxdb* gene on the embryos of C57BL/6J mice. The results of early embryonic development of C57BL/6J embryos were consistent with those of ICR embryos, indicating that *Zxdb* has a very limited role in early embryonic development. At present, the majority of KO mouse models available on the market are based on the C57BL/6J strain. Both the disadvantages and the advantages are quite obvious. The ICR mouse model might be a more suitable choice, especially for research topics related to embryo implantation and reproductive development, and it is worth considering.

Transcriptome and proteome are good tools for studying the effect of KO mouse genes on mice as a whole and have been widely used [46,47]. In our study, we used the transcriptome and proteome together to analyze the overall effect of *Zxdb*-KO mice. The heat map of the *Zxdb*-KO mouse transcriptome shows that the expression of key genes related to cell adhesion, embryo implantation, and decidualization is severely dysregulated. The primary cause may be the defect of adhesion molecules: *Cdh1* and *Cdh2* are significantly downregulated, weakening the initial adhesion between the embryo and the uterine epithelium. The CDH1/CDH2 ratio is often used as a marker for MET/EMT, and the dynamic balance of MET/EMT is crucial for successful pregnancy [17–19]. The mouse uterine tissues selected in this study were from the pre-implantation stage, reflecting the changes in the endometrium before implantation, which can fully and directly reflect the environment before embryo implantation and provide a complete picture of the environment before implantation. However, it provides limited information on the decidualization state after implantation. Changes in the uterine tissues after implantation may require further research. In addition, other adhesion molecules were also affected. The downregulation of *Col26a1* and *Cxcl14* may further disrupt the structure and chemotactic balance of the extracellular matrix (ECM), hindering blastocyst attachment and trophoblast invasion. Downregulation of *Mmp2* may inhibit ECM remodeling, while upregulation of *Adamtsl5* may cause excessive proteolysis, leading to uncontrolled invasion or fibrosis. Although the upregulation of *Bmp2*, *Ptgs2*, *Igf1*, and *Fgf2* attempts to compensatorily promote proliferation, the disordered sequence is prone to cause inflammation or tissue disorders. The upregulation of *Tfrc* reflects metabolic stress and further aggravates the abnormality of the implantation microenvironment. In conclusion, *Zxdb* knockout may disrupt the adhesion receptivity and decidualization ability of the uterine "implantation window", leading to unstable embryo adhesion, invasion obstruction, and insufficient decidual support. Ultimately, it is manifested as increased embryo absorption and a significant decrease in the birth rate.

Furthermore, the differences between the transcriptome and the proteome are also worthy of attention. We found that after *Zxdb* was knocked out, it was more affected by the transcriptome and rarely transmitted to the proteome. On the one hand, it might be because *Zxdb* regulates the transcriptome level more, and on the other hand, it might be limited by the technical limitations of proteomics. Many trace amounts of protein will be submerged by high-abundance proteins during detection, making them undetectable. ZXDB protein is a good example. Here, ZXDB is a trace protein. In our proteomics data, the ZXDB protein was not detected in either the WT or *Zxdb*-KO groups. However, this problem does not exist in the transcriptome. By comparing the correlations between the expression of genes and proteins in the two omics, the potential regulatory relationship between genes and proteins can be quickly understood. Of course, a comprehensive analysis of transcriptomics and proteomics can better assess the overall impact on mice after gene knockout.

Some limitations of this study should also be noted in future research. Firstly, one point worth discussing is that *Zxdb* is a transcription factor. Generally speaking, it may mainly regulate the direction of differentiation and development by regulating gene expression or coordinating with other transcription factors. This research focuses on the relationship

between genes and phenotypes and explores the main pathways and DEGs affected by the deletion of *Zxdb*. However, the motifs or molecular interactions of its action require more in-depth research by subsequent investigators. Secondly, *Zxdb* is an X-linked gene. The impact of the loss of function of this gene on the phenotype of mice is influenced by X chromosome inactivation. Perhaps there will be some differences in the phenotypes of heterozygous mutant and homozygous KO mice, and more mice are needed for further research. Additionally, the lack of clinical family data is a regrettable aspect. If some loss-of-function mutations of the *ZXDB* gene were found in RSA or RIF patients, they would undoubtedly have more significant contributions. Finally, the homologous gene of *Zxdb* in mice, *Zxda*, is a long non-coding gene. The relationship between *Zxda* and *Zxdb* and their impact on the mouse phenotype have not been explored and discussed in this study. More experiments in the future may be needed for further investigation.

5. Conclusions

This study is the first to confirm that the loss of *Zxdb* function is associated with decidualization of the uterus and the reduction in the number of mouse offspring. Secondly, we demonstrated that the loss of *Zxdb* function caused by maternal factors is the cause of these phenotypes. Thirdly, comprehensive experiments including transcriptomics, proteomics, Western blotting, and immunofluorescence confirmed that the expression and distribution of adhesion molecules such as CDH1, CDH2, and BMP2 were abnormal in the endometrial tissue of the uterus before embryo implantation. In conclusion, this study reveals the association between *Zxdb* and the decline in female reproductive capacity in mice, providing a good model for studying embryo implantation and decidualization of the uterus and offering new insights and genetic counseling guidance for researchers and clinicians studying RSA or RIF.

Supplementary Materials: The following supporting information can be downloaded at: https://www.mdpi.com/article/10.3390/cimb48020144/s1.

Author Contributions: Conceptualization, D.L. and S.C.; methodology, Y.T.; software, P.D.; validation, Y.T., Y.Z., M.L., L.L., B.C. and R.Y.; formal analysis, P.D.; resources, S.C.; data curation, R.X., B.C. and P.D.; writing—original draft preparation, Y.T.; writing—review and editing. C.X., H.C., D.L. and S.C. funding acquisition, S.C. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the research grants from the sub-project of the National Key R&D Program (2023YFC2705600, 2023YFC2705602) and the Shanghai Municipal Commission of Science and Technology Program (23ZR1408000).

Institutional Review Board Statement: The experimental process strictly adhered to the regulations of the Animal Ethics Committee of the School of Life Sciences, Fudan University. The content of the animal experiments has obtained ethical approval, with the approval number: No. 2022JS071; approval date 15 March 2022.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available on request from the corresponding author due to pending patent applications.

Acknowledgments: We are grateful to Guoqiang Zhou and Jingmin Yang for their guidance and assistance with gene-edited mice.

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

The following abbreviations are used in this manuscript:

ZXDB	Zinc Finger X-Linked Duplicate B
RSA	Recurrent spontaneous abortion
URSA	Unexplained recurrent spontaneous abortion
RIF	Recurrent Implantation Failure
CDH1	E-cadherin
CDH2	N-cadherin
WOI	Window of implantation
MET	Epithelial–mesenchymal transition
EMT	Mesenchymal–epithelial transition
WT	Wild-type
Zxdb-KO	*Zxdb* knockout
PMSG	Pregnant mare serum gonadotrophin
HCG	Human chorionic gonadotropin
KSOM	K+ Simplex Optimized Medium
COCs	Cumulus Oocyte Complexes
H&E	Hematoxylin–Eosin
WB	Western blot
Neuro-2a	N2a
hESCs	Human embryonic stem cells
ECM	Extracellular matrix
PR	Progesterone receptor
GO	Gene ontology
KEGG	Kyoto Encyclopedia of Genes and Genomes
DEGs	Differentially expressed genes
PCA	Principal components analysis

References

1. Yu, M.; Peng, X.; Li, H.; Xu, Y.; Sun, X.; Chen, J. Gankyrin has a potential role in embryo implantation via activation of STAT3. *Reproduction* **2022**, *163*, 157–165. [CrossRef] [PubMed]
2. Ashary, N.; Tiwari, A.; Modi, D. Embryo Implantation: War in Times of Love. *Endocrinology* **2018**, *159*, 1188–1198. [CrossRef] [PubMed]
3. Carson, D.D.; Bagchi, I.; Dey, S.K.; Enders, A.C.; Fazleabas, A.T.; Lessey, B.A.; Yoshinaga, K. Embryo implantation. *Dev. Biol.* **2000**, *223*, 217–237. [CrossRef] [PubMed]
4. Ojosnegros, S.; Seriola, A.; Godeau, A.L.; Veiga, A. Embryo implantation in the laboratory: An update on current techniques. *Hum. Reprod. Update* **2021**, *27*, 501–530. [CrossRef]
5. Wilcox, A.J.; Weinberg, C.R.; O'Connor, J.F.; Baird, D.D.; Schlatterer, J.P.; Canfield, R.E.; Armstrong, E.G.; Nisula, B.C. Incidence of early loss of pregnancy. *N. Engl. J. Med.* **1988**, *319*, 189–194. [CrossRef]
6. Chard, T. Frequency of implantation and early pregnancy loss in natural cycles. *Baillieres Clin. Obstet. Gynaecol.* **1991**, *5*, 179–189. [CrossRef]
7. Ammon Avalos, L.; Galindo, C.; Li, D.K. A systematic review to calculate background miscarriage rates using life table analysis. *Birth Defects Res. A Clin. Mol. Teratol.* **2012**, *94*, 417–423. [CrossRef]
8. Aplin, J.D.; Ruane, P.T. Embryo-epithelium interactions during implantation at a glance. *J. Cell Sci.* **2017**, *130*, 15–22. [CrossRef]
9. Ramathal, C.Y.; Bagchi, I.C.; Taylor, R.N.; Bagchi, M.K. Endometrial decidualization: Of mice and men. *Semin. Reprod. Med.* **2010**, *28*, 17–26. [CrossRef]
10. Cai, X.; Xu, M.; Zhang, H.; Zhang, M.; Wang, J.; Mei, J.; Zhang, Y.; Zhou, J.; Zhen, X.; Kang, N.; et al. Endometrial stromal PRMT5 plays a crucial role in decidualization by regulating NF-kappaB signaling in endometriosis. *Cell Death Discov.* **2022**, *8*, 408. [CrossRef]
11. Bellver, J.; Simon, C. Implantation failure of endometrial origin: What is new? *Curr. Opin. Obstet. Gynecol.* **2018**, *30*, 229–236. [CrossRef] [PubMed]
12. Okada, H.; Tsuzuki, T.; Murata, H. Decidualization of the human endometrium. *Reprod. Med. Biol.* **2018**, *17*, 220–227. [CrossRef] [PubMed]

13. Wang, C.; Zhao, M.; Zhang, W.Q.; Huang, M.Y.; Zhu, C.; He, J.P.; Liu, J.L. Comparative Analysis of Mouse Decidualization Models at the Molecular Level. *Genes* **2020**, *11*, 935. [CrossRef] [PubMed]
14. Gellersen, B.; Brosens, J.J. Cyclic decidualization of the human endometrium in reproductive health and failure. *Endocr. Rev.* **2014**, *35*, 851–905. [CrossRef]
15. Oghbaei, F.; Zarezadeh, R.; Jafari-Gharabaghlou, D.; Ranjbar, M.; Nouri, M.; Fattahi, A.; Imakawa, K. Epithelial-mesenchymal transition process during embryo implantation. *Cell Tissue Res.* **2022**, *388*, 1–17. [CrossRef]
16. Liu, M.; Chen, X.; Chang, Q.X.; Hua, R.; Wei, Y.X.; Huang, L.P.; Liao, Y.X.; Yue, X.J.; Hu, H.Y.; Sun, F.; et al. Decidual small extracellular vesicles induce trophoblast invasion by upregulating N-cadherin. *Reproduction* **2020**, *159*, 171–180. [CrossRef]
17. Tiwari, A.; Ashary, N.; Singh, N.; Sharma, S.; Modi, D. Modulation of E-Cadherin and N-Cadherin by ovarian steroids and embryonic stimuli. *Tissue Cell* **2021**, *73*, 101670. [CrossRef]
18. Zhang, X.H.; Liang, X.; Liang, X.H.; Wang, T.S.; Qi, Q.R.; Deng, W.B.; Sha, A.G.; Yang, Z.M. The mesenchymal-epithelial transition during in vitro decidualization. *Reprod. Sci.* **2013**, *20*, 354–360. [CrossRef]
19. Richardson, L.S.; Taylor, R.N.; Menon, R. Reversible EMT and MET mediate amnion remodeling during pregnancy and labor. *Sci. Signal.* **2020**, *13*, eaay1486. [CrossRef]
20. Kennedy, T.G.; Gillio-Meina, C.; Phang, S.H. Prostaglandins and the initiation of blastocyst implantation and decidualization. *Reproduction* **2007**, *134*, 635–643. [CrossRef]
21. King, J.S., Jr. Handbook of Physiology, Section 7: Endocrinology, Volume II (in two parts): Female Reproductive System. R. O. Greep and E. B.; Astwood, Section Eds.; R. O.; Greep, Volume Ed.; S. R.; Geiger, Executive Ed. American Physiological Society, Washington, D.C., 1973, vii + 657 and 375 pp. $44.50 (part 1) and $25 (part 2). *Clin. Chem.* **1974**, *20*, 633. [CrossRef]
22. Greig, G.M.; Sharp, C.B.; Carrel, L.; Willard, H.F. Duplicated zinc finger protein genes on the proximal short arm of the human X chromosome: Isolation, characterization and X-inactivation studies. *Hum. Mol. Genet.* **1993**, *2*, 1611–1618. [CrossRef]
23. Al-Kandari, W.; Jambunathan, S.; Navalgund, V.; Koneni, R.; Freer, M.; Parimi, N.; Mudhasani, R.; Fontes, J.D. ZXDC, a novel zinc finger protein that binds CIITA and activates MHC gene transcription. *Mol. Immunol.* **2007**, *44*, 311–321. [CrossRef] [PubMed]
24. Al-Kandari, W.; Koneni, R.; Navalgund, V.; Aleksandrova, A.; Jambunathan, S.; Fontes, J.D. The zinc finger proteins ZXDA and ZXDC form a complex that binds CIITA and regulates MHC II gene transcription. *J. Mol. Biol.* **2007**, *369*, 1175–1187. [CrossRef] [PubMed]
25. Dibus, N.; Zobalova, E.; Monleon, M.A.M.; Korinek, V.; Filipp, D.; Petrusova, J.; Sedlacek, R.; Kasparek, P.; Cermak, L. FBXO38 Ubiquitin Ligase Controls Sertoli Cell Maturation. *Front. Cell Dev. Biol.* **2022**, *10*, 914053. [CrossRef] [PubMed]
26. Gao, Y.; Liu, X.; Tang, B.; Li, C.; Kou, Z.; Li, L.; Liu, W.; Wu, Y.; Kou, X.; Li, J.; et al. Protein Expression Landscape of Mouse Embryos during Pre-implantation Development. *Cell Rep.* **2017**, *21*, 3957–3969. [CrossRef]
27. Mamo, S.; Gal, A.B.; Bodo, S.; Dinnyes, A. Quantitative evaluation and selection of reference genes in mouse oocytes and embryos cultured in vivo and in vitro. *BMC Dev. Biol.* **2007**, *7*, 14. [CrossRef]
28. Ragni, E.; Vigano, M.; Rebulla, P.; Giordano, R.; Lazzari, L. What is beyond a qRT-PCR study on mesenchymal stem cell differentiation properties: How to choose the most reliable housekeeping genes. *J. Cell Mol. Med.* **2013**, *17*, 168–180. [CrossRef]
29. Gu, Y.; Shen, X.; Zhou, D.; Wang, Z.; Zhang, N.; Shan, Z.; Jin, L.; Lei, L. Selection and expression profiles of reference genes in mouse preimplantation embryos of different ploidies at various developmental stages. *PLoS ONE* **2014**, *9*, e98956. [CrossRef]
30. Macklon, N.S.; Geraedts, J.P.; Fauser, B.C. Conception to ongoing pregnancy: The 'black box' of early pregnancy loss. *Hum. Reprod. Update* **2002**, *8*, 333–343. [CrossRef]
31. Cockburn, K.; Rossant, J. Making the blastocyst: Lessons from the mouse. *J. Clin. Investig.* **2010**, *120*, 995–1003. [CrossRef]
32. Yang, Y.; Chen, X.; Saravelos, S.H.; Liu, Y.; Huang, J.; Zhang, J.; Li, T.C. HOXA-10 and E-cadherin expression in the endometrium of women with recurrent implantation failure and recurrent miscarriage. *Fertil. Steril.* **2017**, *107*, 136–143 e132. [CrossRef] [PubMed]
33. Jiang, R.; Ding, L.; Zhou, J.; Huang, C.; Zhang, Q.; Jiang, Y.; Liu, J.; Yan, Q.; Zhen, X.; Sun, J.; et al. Enhanced HOXA10 sumoylation inhibits embryo implantation in women with recurrent implantation failure. *Cell Death Discov.* **2017**, *3*, 17057. [CrossRef] [PubMed]
34. Vasquez, Y.M.; Wang, X.; Wetendorf, M.; Franco, H.L.; Mo, Q.; Wang, T.; Lanz, R.B.; Young, S.L.; Lessey, B.A.; Spencer, T.E.; et al. FOXO1 regulates uterine epithelial integrity and progesterone receptor expression critical for embryo implantation. *PLoS Genet.* **2018**, *14*, e1007787. [CrossRef] [PubMed]
35. Tur-Torres, M.H.; Garrido-Gimenez, C.; Alijotas-Reig, J. Genetics of recurrent miscarriage and fetal loss. *Best. Pract. Res. Clin. Obstet. Gynaecol.* **2017**, *42*, 11–25. [CrossRef]
36. Bi, Y.; Huang, W.; Yuan, L.; Chen, S.; Liao, S.; Fu, X.; Liu, B.; Yang, Y. HOXA10 improves endometrial receptivity by upregulating E-cadherindagger. *Biol. Reprod.* **2022**, *106*, 992–999. [CrossRef]
37. Xu, B.; Sun, X.; Li, L.; Wu, L.; Zhang, A.; Feng, Y. Pinopodes, leukemia inhibitory factor, integrin-beta3, and mucin-1 expression in the peri-implantation endometrium of women with unexplained recurrent pregnancy loss. *Fertil. Steril.* **2012**, *98*, 389–395. [CrossRef]

38. Wu, F.; Chen, X.; Liu, Y.; Liang, B.; Xu, H.; Li, T.C.; Wang, C.C. Decreased MUC1 in endometrium is an independent receptivity marker in recurrent implantation failure during implantation window. *Reprod. Biol. Endocrinol.* **2018**, *16*, 60. [CrossRef]
39. Luo, J.; Zhu, H.; Chang, H.M.; Lin, Y.M.; Yang, J.; Leung, P.C.K. The regulation of IGFBP3 by BMP2 has a role in human endometrial remodeling. *FASEB J.* **2020**, *34*, 15462–15479. [CrossRef]
40. Sonehara, K.; Yano, Y.; Naito, T.; Goto, S.; Yoshihara, H.; Otani, T.; Ozawa, F.; Kitaori, T.; Biobank Japan, P.; Matsuda, K.; et al. Common and rare genetic variants predisposing females to unexplained recurrent pregnancy loss. *Nat. Commun.* **2024**, *15*, 5744. [CrossRef]
41. Wang, W.; Taylor, R.N.; Bagchi, I.C.; Bagchi, M.K. Regulation of human endometrial stromal proliferation and differentiation by C/EBPbeta involves cyclin E-cdk2 and STAT3. *Mol. Endocrinol.* **2012**, *26*, 2016–2030. [CrossRef]
42. Mantena, S.R.; Kannan, A.; Cheon, Y.P.; Li, Q.; Johnson, P.F.; Bagchi, I.C.; Bagchi, M.K. C/EBPbeta is a critical mediator of steroid hormone-regulated cell proliferation and differentiation in the uterine epithelium and stroma. *Proc. Natl. Acad. Sci. USA* **2006**, *103*, 1870–1875. [CrossRef]
43. Zhu, M.; Yi, S.; Huang, X.; Meng, J.; Sun, H.; Zhou, J. Human chorionic gonadotropin improves endometrial receptivity by increasing the expression of homeobox A10. *Mol. Hum. Reprod.* **2020**, *26*, 413–424. [CrossRef]
44. Adiguzel, D.; Celik-Ozenci, C. FoxO1 is a cell-specific core transcription factor for endometrial remodeling and homeostasis during menstrual cycle and early pregnancy. *Hum. Reprod. Update* **2021**, *27*, 570–583. [CrossRef]
45. Xu, Z.; Yang, H.H.; Chen, H.Z.; Huang, B.Z.; Yang, M.; Liao, Z.H.; Xiao, B.Q.; Chen, H.Q.; Ran, J. ZEB1 Promotes Epithelial-Mesenchymal Transition of Endometrial Epithelial Cells and Plays a Critical Role in Embryo Implantation in Mice. *Reprod. Sci.* **2024**, *32*, 1331–1337. [CrossRef]
46. Ghazalpour, A.; Bennett, B.; Petyuk, V.A.; Orozco, L.; Hagopian, R.; Mungrue, I.N.; Farber, C.R.; Sinsheimer, J.; Kang, H.M.; Furlotte, N.; et al. Comparative analysis of proteome and transcriptome variation in mouse. *PLoS Genet.* **2011**, *7*, e1001393. [CrossRef]
47. West, D.B.; Engelhard, E.K.; Adkisson, M.; Nava, A.J.; Kirov, J.V.; Cipollone, A.; Willis, B.; Rapp, J.; de Jong, P.J.; Lloyd, K.C. Transcriptome Analysis of Targeted Mouse Mutations Reveals the Topography of Local Changes in Gene Expression. *PLoS Genet.* **2016**, *12*, e1005691. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.