current issues in molecular biology

MDPI

Article

Subfunctionalization of *CpSVP-Y* Paralog Contributes to Male Reproductive Fitness in Papaya

Julie Nguyen-Edquilang [1,2], Jingjing Yue [1] and Ray Ming [1,2,*]

[1] FAFU and UIUC-SIB Joint Center for Genomics and Biotechnology, Fujian Provincial Key Laboratory of Haixia Applied Plant Systems Biology, Key Laboratory of Genetics, Breeding and Multiple Utilization of Corps, Ministry of Education, Fujian Agriculture and Forestry University, Fuzhou 350002, China; jknguyn2@illinois.edu (J.N.-E.)

[2] Department of Plant Biology, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

* Correspondence: rayming@illinois.edu

Abstract

Papayas possess three sex genotypes: female XX, male XY, and hermaphrodite XY^h. Only male plants produce long peduncles with numerous flowers, a trait that improves reproductive success. The gene *SHORT VEGETATIVE PHASE* (*CpSVP*) is located on the Y chromosome, absent from the X chromosome, and disrupted in the Y^h chromosome, making it a key candidate gene for long peduncles in male plants. An autosomal *CpSVP* allele (*CpSVP-A*) was also annotated in a papaya genome. The overexpression of the male allele in *Arabidopsis* increases pedicel length, which supports its role in pedicel elongation. Unexpectedly, the autosomal allele *CpSVP-A* produced a similar phenotype as the male one, while the hermaphroditic allele (*CpSVP-*Y^h) did not cause any significant change in pedicel length. Additionally, only the male allele rescued early flowering in an *Atsvp* mutant, indicating that it regulates both pedicel length and flowering time. In contrast, the autosomal allele only affected pedicel elongation and had no effect on flowering time. Together, these results demonstrate that *CpSVP-Y* underwent subfunctionalization, retaining both peduncle elongation and flowering time functions, while *CpSVP-A* likely lost the latter. Thus, *CpSVP-Y* contributes to male reproductive fitness through enhanced peduncle development without affecting sex determination, potentially regulated by Y chromosome-specific cis-elements. This study provides new insights into the genetic basis of sexually dimorphic traits in papayas.

Keywords: papaya; *CpSVP*; sex chromosome; peduncle elongation; flowering time

Check for updates

Academic Editor: Abdullah Shalmani

Received: 2 January 2026
Revised: 16 January 2026
Accepted: 23 January 2026
Published: 1 February 2026

Copyright: © 2026 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license.

1. Introduction

Papayas (*Carica papaya* L.) are a tropical fruit consumed worldwide with high nutritional value. Papaya trees are tridioecious (males, hermaphrodites, and females), in which their sex types are controlled by a pair of sex chromosomes that are approximately 7 million years old [1]. There are two slight sequence differences: the Y chromosome determines males, and the Y^h chromosome determines hermaphrodites. These two Y chromosomes diverged about 4000 years ago, with only a 0.4% sequence difference between them [2,3], making the identification of sex determination gene(s) difficult due to high sequence similarity. Any combination of these Y chromosomes is inviable at the seed development stage (YY, YY^h, or Y^hY^h) due to the presence of homozygous lethal gene(s) [4]. The sex-determining region (SDR) of the X and Y/Y^h chromosome is 3.4 Mb and 8.1 Mb in length, respectively [1,3]. This relatively small region contains the sex determination

genes as well as genes responsible for sex-linked traits such as peduncle length. The long peduncle trait always co-segregates with the male plant; therefore, the gene responsible for peduncle length is a gene that is located on the male Y chromosome but not on the hermaphrodite Y^h or female X chromosome [5,6]. Meanwhile, the sequence differences in SHORT VEGETATIVE PHASE (*SVP*) between male and hermaphrodite papayas have led to the hypothesis that *SVP* is involved in male–hermaphrodite differentiation [7,8]..

SHORT VEGETATIVE PHASE (*SVP*), a MADS box transcription factor, has been well characterized in *Arabidopsis* [9]. It acts in a dosage-dependent manner to repress flowering by forming a complex with FLOWERING LOCUS C (FLC). *AtSVP,* along with *AGAMOUS-LIKE24* (*AGL24*), determines floral meristem identity in *Arabidopsis* [10]. During the vegetative phase, *AtSVP* forms a dimer with *FLC*, which binds to CArG boxes to repress *FLOWERING LOCUS T* (*FT*) in the leaves and *SUPPRESSOR OF OVEREXPRESSION OF CONSTANS1* (*SOC1*) in the shoot apical meristem [11]. How *SVP* is regulated remains unclear, its function may be modulated through interactions with other MADS-box proteins. MADS box transcription factors activate or repress transcription by binding directly to CArG boxes. While the role of *SVP* genes in flowering time and floral meristem identity seems to be conserved in some species, it has been shown that *SVP* genes can have diverse functions. For example, the overexpression of *Oryza sativa SVP*-like genes in rice implicates their involvement in brassinosteroid signaling and shoot development [12]. In addition, pedicels were elongated when overexpressing *Medicago truncatula SVP* in *Arabidopsis*, in addition to delayed flowering and a floral defect phenotype [13].

By comparing the male, hermaphrodite, and female papaya chromosome sequences, a *SHORT VEGETATIVE PHASE* (*SVP*) coding sequence was found absent on the X chromosome, an intact *SVP* sequence was annotated on the Y chromosome (*CpSVP-Y*), and a truncated *SVP* sequence without the first exon was located on the Y^h chromosome (*CpSVP-*Y^h). In addition, there is an autosomal copy of *SVP* (*CpSVP-A*) in the papaya genome [1]. Since the elongated peduncle trait is sex-linked to male papaya trees, it is thus hypothesized that the intact *CpSVP* genes located on the Y chromosome are responsible for this trait, while the truncated *CpSVP* on the Y^h chromosome and intact *CpSVP* on the autosome are not.

The objective of this study is to functionally characterize *CpSVP* alleles in *Arabidopsis* to infer their potential roles in peduncle elongation and flowering time regulation in papayas and to discuss implications for sex chromosome evolution. In this study, the functional characterization of *CpSVP-Y*, *CpSVP-*Y^h, and their paralog *CpSVP-A* was carried out in *Arabidopsis*. *CpSVP-Y* in transgenic *Arabidopsis* is capable of pedicel elongation with altered floral organs, including enlarged sepals and extra petals but no defects in stamen or carpel structure. *CpSVP-Y* can also rescue the mutant *svp* phenotype of short pedicels and early flowering. However, *CpSVP-A* was found to regulate pedicel elongation but not flowering time in *Arabidopsis*. These findings suggest that subfunctionalization has occurred between *CpSVP-Y* and *CpSVP-A,* with functional divergence likely mediated by promoter regulatory elements. This study provides preliminary insights into the regulatory mechanisms of male organ development in papayas and identifies potential targets for molecular breeding.

2. Materials and Methods

2.1. Plant Material

The experiments were carried out using *Arabidopsis thaliana* (Col-0). The *svp-31* mutant (SALK_026551) was obtained from the Arabidopsis Biological Resource Center. Plants were maintained in controlled-environment growth chambers programmed for 16 h of light at 23 °C and 8 h of darkness at 18 °C, with relative humidity held at 50%.

2.2. Flowering Time Measurements

To determine flowering time, the number of rosette leaves was recorded once the primary inflorescence extended to roughly 2 cm. Pedicel length analyses were performed using nine independent T4 lines per genotype. Sixteen plants were grown for each line, and measurements were taken from the uppermost and lowermost six pedicels on each plant.

2.3. Plasmid Construction

Promoter regions for *CpSVP-Y* and *CpSVP-Y^h* were identified by aligning their coding sequences to male and hermaphrodite papaya pseudomolecules. Based on these alignments, 2 kb (*CpSVP Y*) and 1 kb (*CpSVP-Y^h*) upstream regions were retrieved. Predicted cis-regulatory motifs were annotated using the PLACE database [14]. Promoter fragments were amplified with specific primers (*CpSVP-Y^h* (1kb) and *CpSVP-Y* (2kb); see Table S1) and inserted into pMDC162 GUS [15]. For overexpression constructs, the coding regions of *CpSVP-Y*, *CpSVP-Y^h*, and CpSVP-A were amplified using primers listed in Table S1 and cloned into pMDC32 [15]. All plasmids were sequence-verified by Sanger sequencing.

2.4. Plant Transformation and Analysis of Transgenic Lines

Constructs were introduced into *Agrobacterium tumefaciens* GV3101 and subsequently used to transform *Arabidopsis* via the floral dip method [16]. At least twelve independent T1 transformants were generated for each construct. Seeds were surface-sterilized, plated on half-strength MS medium supplemented with hygromycin and cefotaxime, stratified at 4 °C for two days, and then transferred to growth chambers. Seedlings with four leaves were transplanted to the soil. T2 progeny were screened for single-locus insertions based on a 3:1 segregation ratio on selective medium. Genotyping was performed using gene-specific primers and the Kapa3G Plant PCR Kit (Cat. KK7252).

2.5. Quantitative RT-PCR (qPCR)

Total RNA was isolated with Trizol reagent (Invitrogen, Waltham, MA, USA) according to the supplier's protocol. RNA concentration was determined using a NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA), and RNA quality was evaluated by electrophoresis on a 1% agarose gel. For cDNA synthesis, 1 µg of total RNA was used as input for the Maxima First-Strand cDNA Synthesis Kit with dsDNase (Cat. K1672, Thermo Fisher Scientific, Waltham, MA, USA). For quantitative PCR, each reaction consisted of 1.6 µL of cDNA template, primers at 0.5 µM, and 10 µL of PowerUp SYBR Green Master Mix (Cat. A25741, Applied Biosystems, Waltham, MA, USA), with nuclease-free water added to volume. Amplification was performed on a CFX96 Touch real-time PCR instrument (Bio-Rad, Hercules, CA, USA). Primer sets used in this study (Table S2) included both previously published sequences and additional primers designed using Primer3 software [17,18]. The cycling conditions are 50 °C for 2 min and 95 °C for 2 min, followed by 35 cycles of denaturation at 95 °C for 15 s and annealing/extension at 60 °C for 1 min, ending with a melt-curve analysis. For expression analysis, a minimum of three independent T3 lines were examined for each gene, with two technical replicates per line. Transcript levels were normalized to *ACTIN2* and *ELONGATION FACTOR 1-α* using the ΔΔCt calculation method [19].

3. Results

3.1. Identification of the CpSVP Gene in Papayas

SVP is annotated in the male and hermaphrodite genomes, as well as a paralogous autosomal copy [1]. *CpSVP-Y* and *CpSVP-A* share 94% similarity in their coding sequence and contain the MADS-box and K-box domain. *CpSVP-Y* and *CpSVP-A* have an open

reading frame of 669 bp, which encodes for 223 amino acids. *CpSVP-Y*^{h} is 483 bp and encodes a 161-amino-acid truncated protein missing the MADS-box domain at its N-terminal. *CpSVP-Y* and *CpSVP-Y*^{h} share 71% similarity in their coding sequence, and both contain the K-box domain.

3.2. Male and Autosomal Copies of CpSVP Regulate Pedicel Elongation

To characterize the function of different copies of *CpSVP* in regulating pedicel length, overexpression lines were generated in *Arabidopsis* for each allele containing the open reading frame driven under the constitutive cauliflower mosaic virus (CaMV) 35S promoter. Both *35S:CpSVP-Y* and *35S:CpSVP-A* transgenic plants had a mean pedicel length of 8.9 mm, an increase of 30.9% when compared to the wild-type pedicels (Figure 1). For *35S:CpSVP-Y*^{h}, there was no significant difference in pedicel length when compared to wild-type. Only the *CpSVP-Y* and *CpSVP-A* alleles resulted in pedicel elongation.




Figure 1. Pedicel length measured in 40-day-old overexpression lines; 12 pedicels from 16 individual plants were measured for each T4 line, and 9 independent lines were used in this study (n = 192). Values are represented as means ± standard error (SE). Results were compared to Col-0 and analyzed using one-way ANOVA followed by Dunnett's multiple comparison test (** $p < 0.01$).

However, the *Arabidopsis AtSVP* knockout line (*svp-31*) had shorter pedicels when compared to wild-type plants ($p < 0.05$, Student's t-test). The average pedicel length for *svp-31* was 5 mm, a 26.5% decrease from wild-type pedicels (Figure 2). The overexpression constructs of the three alleles (*35S:CpSVP-Y*, *35S:CpSVP-Y*^{h}, and *35S:CpSVP-A*) and two translational constructs (native promoter fused with genes, *CpSVP-Y*^{h} *(1kb):CpSVP-Y*^{h} and *CpSVP-Y (2kb):CpSVP-Y*) were introduced into this *svp-31* mutant background to test if any of the alleles could rescue the short pedicel mutant phenotype. *35S:CpSVP-Y svp-31* and *35S:CpSVP-A svp-31* are able to complement the mutant phenotype and have a mean pedicel length of 10.7 mm and 8.7 mm, respectively. *35S:CpSVP-Y*^{h} *svp-31* and *CpSVP-Y*^{h}*(1kb):CpSVP-Y*^{h} *svp-31* did not have any significant difference in pedicel length when compared to *svp-31*, indicating the reduced pedicel phenotype in the knockout mutant was not rescued by the *CpSVP-Y*^{h} gene, no matter which promoter sequence (CaMV35S or native promoter) was used (Figure 2). *CpSVP-Y(2kb):CpSVP-Y svp-31* could also rescue the mutant phenotype and had a mean pedicel length of 7.6 mm. These results indicate that *CpSVP-Y* and *CpSVP-A* play a role in pedicel elongation but not *CpSVP-Y*^{h}.




Figure 2. Pedicel length measured in 40-day-old lines for complementation study; 12 pedicels from 16 individual plants were measured for each T4 line, and 9 independent lines were used in this study (n = 192). Values are represented as means ± standard error (SE). Results were compared to *svp-31* and analyzed using one-way ANOVA followed by Dunnett's multiple comparison test (** $p < 0.01$).

The text continues here (Figures 1 and 2).

3.3. CpSVP Regulates Flowering Time

SVP has been demonstrated to be involved in regulating flowering time by binding to FT [11]. *SVP* acts as a flowering repressor, so knocking out *SVP* should result in an early flowering phenotype. Indeed, we observed that the knockout (*svp-31*) line flowers earlier than wild-type ($p < 0.01$, Student's *t*-test; see Table 1). We introduced the overexpression constructs of the three alleles (*35S:CpSVP-Y*, *35S:CpSVP-Y^h*, and *35S:CpSVP-A*) and two translational constructs (native promoter fused with genes—*CpSVP-Y^h_(1kb):CpSVP-Y^h* and *CpSVP-Y (2kb):CpSVP-Y*) into this *svp-31* mutant background, to determine if any of the alleles could rescue the early flowering phenotype. *35S:CpSVP-A svp-31*, *35S:CpSVP-Y^h svp-31*, and *CpSVP-Y^h(1kb):CpSVP-Y^h svp-31* had the same early flowering phenotype as *svp-31* (Table S1). Only *35S:CpSVP-Y svp-31* and *CpSVP-Y_(2kb):CpSVP-Y svp-31* were able to rescue the early flowering phenotype in the knockout mutant, restoring the flowering time that is comparable to wild-type Col-0 (Table S1). Interestingly, overexpression lines of *CpSVP-Y* in the Col-0 background flower later than wild-type ($p < 0.01$, Student's *t*-test; see Table S1). Besides flowering time, additional morphological floral defects not found in the *svp-31* or Col-0 background were observed. In addition to elongated pedicels, both *35S:CpSVP-Y* and *35S:CpSVP-A* plants in the *svp-31* and Col-0 backgrounds had morphological defects, including extra petals and enlarged sepals, but not in *CpSVP-Y_(2kb):CpSVP-Y* (Figure 3). These enlarged sepals did not dehisce and remained attached to the siliques throughout senescence (Figure 4).

Table 1. Flowering time of overexpression and complementation lines.

Genotype	Number of Rosette Leaves
WT (Col-0)	11.8 ± 0.16
35S:CpSVP-A	12 ± 0.39
35S:CpSVP-Y	13.1 ± 0.29 **
35S:CpSVP-Y^h	12.3 ± 0.24
svp-31	8.5 ± 0.5 **
35S:CpSVP-A svp-31	9 ± 0.18 **
35S:CpSVP-Y svp-31	12.4 ± 0.48
35S:CpSVP-Y^h svp-31	8.2 ± 0.19 **
CpSVP-Y^h(1kb):CpSVP-Y^h svp-31	7.4 ± 0.23 **
CpSVP-Y(2kb):CpSVP-Y svp-31	11.6 ± 0.2

Individual plants per genotype (n = 8) were measured. Values are means ± standard error (SE). Asterisks indicate significant differences as compared with WT. (** $p < 0.01$, Student's t-test).




Figure 3. *CpSVP-Y* and *CpSVP-A* complemented flower in *svp-31* background. These lines are 40-day-old plants. Arrows depict enlarged sepals and abnormal petal numbers. Scale bars represent 1 mm.




Figure 4. Phenotypic effects of CpSVP-Y on pedicel and sepal development. Shown are (**A**) *svp-31* mutant (mean pedicel length = 5.6 mm), (**B**) wild-type (Col-0) (mean pedicel length = 6.8 mm), and (**C**) *CpSVP-Y* complementation line in the *svp-31* background (mean pedicel length = 10.7 mm), all at 40 days after germination. Arrows point to elongated pedicels and enlarged, non-dehiscent sepals in (**C**). Scale bars represent 5 mm. Comparison of *svp-31* (mean pedicel length = 5.6 mm) (**A**), Col-0 (mean pedicel length = 6.8 mm) (**B**), and *CpSVP-Y svp-31* at 40 days after germination (mean pedicel length = 10.7 mm) (**C**). Arrows depict elongated pedicels and enlarged sepals. These lines are 40-day-old plants. Scale bars represent 5 mm.

3.4. Deciphering the Mechanism of Pedicel Elongation

A previous field experiment showed that exogenous application of GA3 to papaya flowers caused peduncles to elongate [7]. *AtGA20ox2* plays a key role in GA biosynthesis, encoding the rate-limiting enzyme that catalyzes the second-to-last step of bioactive GAs [20,21]. To understand the role of *CpSVP* in GA signaling, the transcript levels of *AtGA20ox2* in the overexpression and complemented lines were examined. In all three over-

expression lines of *CpSVP*, *GA20ox2* was downregulated in the junction tissue, especially for *CpSVP-Y*. However, *GA20ox2* was significantly upregulated in young and old pedicels for all overexpression lines (Figure 5). In the complemented lines, *GA20ox2* is significantly downregulated in the junction of *35S:CpSVP-Y svp-31* and *CpSVP-Y(2kb):CpSVP-Y svp-31* and expressed at very low levels in young pedicels (Figure 6). However, in complemented lines that express *CpSVP-A* and *CpSVP-Y*^{h}, the *GA20ox2* expression remained high in both junction and young pedicel. All lines showed high *GA20ox2* expression in old pedicels, suggesting that there is no expression correlation between *CpSVP* and *GA20ox2* in this specific tissue.




Figure 5. Relative expressions of *CpSVP* and *GA20ox2* in overexpression lines. The qPCR expression level of each construct was normalized to *ACTIN* and *EIF1* expression in wild-type Col-0 plants. Values were obtained from three biological replicates with two technical replicates each and are presented as means ± standard error (SE). Statistical significance was determined by one-way ANOVA followed by Tukey's honestly significant difference (HSD) test. Different lowercase letters (a, b, c) above bars indicate significant differences among different constructs within the same tissue ($p < 0.05$). Bars sharing the same letter are not significantly different. Tissue colors indicate junction (red), old pedicel (blue), and young pedicel (green).




Figure 6. Relative expression of *CpSVP* and *GA20ox2* in complementation lines. The qPCR expression level of each construct was normalized to *ACTIN* and *EIF1* expression in *svp-31* plants. Values were obtained from three biological replicates and two technical replicates and are presented as means ± standard error (SE). Statistical significance was determined by one-way ANOVA followed by Tukey's honestly significant difference (HSD) test. Different lowercase letters (a, b, c, d) above bars indicate significant differences among different constructs within the same tissue ($p < 0.05$). Bars sharing the same letter are not significantly different. Tissue colors indicate junction (red), old pedicel (blue), and young pedicel (green).

4. Discussion

4.1. Specific Alleles of CpSVP Are Involved in Peduncle Elongation

The long peduncle trait co-segregates only with the male plant; therefore, the gene responsible for peduncle length is likely located on the male Y chromosome but not on the hermaphrodite Y^h or female X chromosome [7]. *SVP* has been identified in the male and hermaphrodite genomes, as well as a paralogous autosomal copy in the autosome [1]. To infer their potential functions, these alleles were functionally characterized in *Arabidopsis*. *35S:CpSVP-A* (autosomal copy) showed elongated pedicels and altered floral morphology compared to wild-type. Furthermore, the elongated pedicels were also observed in transgenic *Arabidopsis* carrying *35S:CpSVP-Y*. *35S:CpSVP-Y* (male copy), *35S:CpSVP-A* (autosomal copy), and *CpSVP-Y(2kb):CpSVP-Y* were able to complement the *svp* mutant's short pedicel phenotype. These findings indicate that both male and autosomal copies of *CpSVP* have the potential to control pedicel length in the *Arabidopsis* system. Unexpectedly, the autosomal copy also exhibited the ability to regulate pedicel length in transgenic *Arabidopsis*. This copy is present in all three sex types of papayas, but only male plants have the long peduncle phenotype, suggesting that the peduncle elongation function of the autosomal al-

lele is not expressed in papayas. This could be achieved through the control of *cis*-elements or the suppression of expression at the transcription and/or translational level. Further experiments are required to confirm the role of the autosomal copy in papaya plants.

On the other hand, *35S:CpSVP-Y^h* (hermaphrodite copy) and *CpSVP-Y^h (1kb):CpSVP-Y^h* had no effect on pedicel length in transgenic *Arabidopsis*. These observations coincide with our hypothesis that *CpSVP-Y^h* is a non-functional copy in hermaphroditic papayas that cannot control peduncle length. Since the gene structure of *CpSVP-Y^h* was disrupted by a transposon insertion at its first exon site, it encodes a truncated protein without the MADS domain at its N-terminal, thus presumably affecting its protein function. The authors of [8] hypothesize that *SVP* is involved in male–hermaphrodite differentiation in papayas based on the *SVP* coding sequence difference between male and hermaphroditic papaya plants. Based on observations in all three *CpSVP* allelic overexpression lines, no defects in stamen or carpel were found, so *CpSVP* is unlikely to be the gene responsible for sex determination in papayas.

In summary, the functional data from *Arabidopsis* indicate that the *CpSVP-Y* and *CpSVP-A* alleles can affect peduncle elongation in heterologous systems. The ancestral peduncle elongation function is retained in both copies, but its phenotypic expression is restricted to male papayas, most likely mediated by differential expression regulation. This expression regulation represents a key hypothesis for explaining sex-specific traits and should be prioritized in our future research.

4.2. Conserved Role of CpSVP in Flowering Time

SVP has been shown to be involved in controlling flowering time [9,11]; for example, *35S:AtSVP* lines are late in flowering and convert inflorescences into shoot-like structures [22]. In this study, this conserved role of *SVP* was observed, particularly in a specific allele of *CpSVP*. Overexpression lines of *CpSVP-Y* in the Col-0 background flowered later than wild-type, and this male copy was able to rescue the early flowering phenotype of the *Arabidopsis* knockout line *svp-31* in the complementation experiment, indicating that *CpSVP-Y* is involved in flowering time regulation. However, the overexpression of *CpSVP-Y^h* and *CpSVP-A* in *Arabidopsis* had no effect on flowering time. In nature, male papaya plants flower earlier than their female and hermaphrodite counterparts. *CpSVP-A* does not affect flowering time in *Arabidopsis*, unlike its *Arabidopsis* homolog *AtSVP*, suggesting the autosomal copy lost the ability to regulate flowering time, but the male allele retained this conserved role. The shared role of male and autosomal *CpSVP* in pedicel elongation but difference in flowering time regulation indicates that there is a subfunctionalization of these two *CpSVP* alleles in papayas.

Arabidopsis thaliana and *Carica papaya* diverged from a common ancestor 72 million years ago [23], and since this divergence, there have been two whole-genome duplication events in *Arabidopsis* [24]. Since a heterologous system was used to test the function of *CpSVP-A*, the possibility that *CpSVP-A* is involved in flowering time regulation in papayas cannot be ruled out. Ectopic expression of *Medicago truncatula SVP* in *Arabidopsis* delayed flowering and had floral defects such as long pedicels, but only floral defects were observed when transformed into Medicago [13]. In addition, the overexpression of *CpSVP-Y* and *CpSVP-A* caused floral defects, including enlarged sepals and extra petals, suggesting the genetic redundancy of these genes in flower development. However, *CpSVP-Y(2kb):CpSVP-Y* did not have altered floral organs, unlike the construct driven under the CaMV 35S promoter, suggesting that the promoter specifies the location where *CpSVP-Y* is expressed to minimize its deleterious effect on floral organs. In summary, *CpSVP-Y* retains its dual ability to regulate flower stalk structure and flowering time through subfunctionalization, which provides a direct molecular basis for enhancing male reproductive fitness. The

specific realization of the trait of long pedicels in male plants is likely to still require the drive of Y-chromosome-specific cis-elements.

4.3. The Preliminary Research on the Hormone Control of the Pedicel Development

Exogenous applications of gibberellic acid 3 (GA3) on female, hermaphrodite, and male papaya flowers resulted in an increase in peduncle length for all sexes [7]. This suggests a potential interaction between GA and peduncle growth. We identified four GA-related cis-elements in 2kb male and hermaphrodite *CpSVP* promoter regions using PLACE [14]. GA negatively regulates *GA20ox2* in a feedback loop. Thus, we surveyed the transcript levels of *GA20ox2*, a key enzyme in GA biosynthesis, in our transgenic *Arabidopsis* to see if the three alleles of *CpSVP* had any effect on its expression. In overexpression lines, *GA20ox2* was upregulated in young and old pedicels and downregulated in the junction, no matter which allelic copy of *CpSVP* was expressed. There is a stronger reduction in *GA20ox2* expression (4-fold) in the young pedicels of plants expressing *35S:CpSVP-Y* than in the other two lines (1- to 2-fold). Since the endogenous *AtSVP* was expressed in these transgenic lines, the effect contributing from papaya *SVP* might be overturned or minimized. Therefore, we surveyed the expression in the knockout line *svp-31*. When *CpSVP-Y* was expressed in *svp-31* plants, *GA20ox2* was significantly downregulated in the junction and young pedicel. This suggests a correlative link between high *CpSVP-Y* expression and reduced *GA20ox2* levels in these specific tissues, potentially implicating GA metabolism in pedicel development. On the other hand, we observed a positive expression correlation between *CpSVP-A*/*CpSVP-Y*^{h} and *GA20ox2* in both junction and young pedicel tissue, an opposite trend compared to *CpSVP-Y*. In the old pedicels, the *GA20ox2* expression remained high in all constructs, suggesting no direct correlation between *CpSVP* and *GA20ox2*. The correlation between *CpSVP-Y* and *GA20ox2* provides preliminary evidence for a specific association between *CpSVP-Y* and gibberellin (GA) signaling, while the direct mechanism underlying its interaction with gibberellin signaling remains to be validated through genetic and biochemical approaches in subsequent studies.

Optimal inflorescence architecture plays a pivotal role in development by ultimately determining the number of flowers that can set fruit and seed [25]. Peduncle length contributes to an optimal inflorescence structure, which helps increase the competitiveness of plants [26]. Male papaya plants are the only sex type to bear a long peduncle with a large number of flowers situated on pedicels. Therefore, a long peduncle allows male plants to gain a reproductive advantage by providing more space for flower growth per tree and increasing the capacity of flower bearing. In addition, pollen from male flowers has a higher wind pollination success rate with the aid of a long peduncle [27]. *CpSVP-Y* plays a major role in peduncle elongation and allows males with their elongated peduncles to remain competitive against hermaphrodites. Simultaneously, the *CpSVP-Y* copy retained its early flowering property like its *Arabidopsis* homolog, which allows male papaya plants to flower before female and hermaphroditic plants. This increases the outcrossing pollination rate of the Y chromosome in the papaya population and maintains high genetic diversity, while self-pollinated hermaphroditic plants usually have lower genetic diversity and possibly accumulate deleterious alleles. Although the pedicel structures of *Arabidopsis thaliana* (an annual herb) and papaya (a woody tree) are not directly homologous, in this study, *Arabidopsis thaliana* was used as a model system with powerful genetic tools and known gene networks to reveal the regulatory principles that may be suitable for papaya peduncle elongation. The mechanism insights obtained from *Arabidopsis thaliana* provide key directions and candidate genes for targeted gene function verification and breeding applications in papayas.

Supplementary Materials: The following supporting information can be downloaded at https://www.mdpi.com/article/10.3390/cimb48020166/s1.

Author Contributions: Conceptualization, J.N.-E.; methodology, J.N.-E.; validation, J.N.-E., writing—original draft preparation, J.N.-E.; writing—review and editing, J.N.-E., R.M. and J.Y.; supervision, R.M. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the US National Science Foundation (NSF) Plant Genome Research Program Award DBI-1546890 to R.M., and by a startup fund from Fujian Agriculture and Forestry University to R.M.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The original contributions presented in this study are included in the article/Supplementary Materials. Further inquiries can be directed to the corresponding author.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Wang, J.; Na, J.-K.; Yu, Q.; Gschwend, A.R.; Han, J.; Zeng, F.; Aryal, R.; VanBuren, R.; Murray, J.E.; Zhang, W.; et al. Sequencing Papaya X and Y^h Chromosomes Reveals Molecular Basis of Incipient Sex Chromosome Evolution. *Proc. Natl. Acad. Sci. USA* **2012**, *109*, 13710–13715. [CrossRef]
2. VanBuren, R.; Zeng, F.; Chen, C.; Zhang, J.; Wai, C.M.; Han, J.; Aryal, R.; Gschwend, A.R.; Wang, J.; Na, J.-K.; et al. Origin and Domestication of Papaya Y^h Chromosome. *Genome Res.* **2015**, *25*, 524–533. [CrossRef]
3. Yue, J.; VanBuren, R.; Liu, J.; Fang, J.; Zhang, X.; Liao, Z.; Wai, C.M.; Xu, X.; Chen, S.; Zhang, S.; et al. SunUp and Sunset Genomes Revealed Impact of Particle Bombardment Mediated Transformation and Domestication History in Papaya. *Nat. Genet.* **2022**, *54*, 715–724. [CrossRef]
4. Xiang, T.; Zhu, Y.; Wang, Y.; Chen, X.; Zhang, Z.; Lai, J.; Zhou, P.; Ming, R.; Yue, J. The Dynamic Regulatory Network of Stamens and Pistils in Papaya. *BMC Plant Biol.* **2025**, *25*, 254. [CrossRef]
5. Liu, Z.; Moore, P.H.; Ma, H.; Ackerman, C.M.; Ragiba, M.; Yu, Q.; Pearl, H.M.; Kim, M.S.; Charlton, J.W.; Stiles, J.I.; et al. A Primitive Y Chromosome in Papaya Marks Incipient Sex Chromosome Evolution. *Nature* **2004**, *427*, 348–352. [CrossRef]
6. Zhou, Y.; Pang, Z.; Wang, W.; Ming, R. Identification and Molecular Analysis of the Y-Specific *CpMp* Gene Controlling Long Male Peduncles in Papaya. *Nat. Commun.* **2025**, *16*, 10638. [CrossRef] [PubMed]
7. Han, J.; Murray, J.E.; Yu, Q.; Moore, P.H.; Ming, R. The Effects of Gibberellic Acid on Sex Expression and Secondary Sexual Characteristics in Papaya. *HortScience* **2014**, *49*, 378–383. [CrossRef]
8. Ueno, H.; Urasaki, N.; Natsume, S.; Yoshida, K.; Tarora, K.; Shudo, A.; Terauchi, R.; Matsumura, H. Genome Sequence Comparison Reveals a Candidate Gene Involved in Male-Hermaphrodite Differentiation in Papaya (*Carica papaya*) Trees. *Mol. Genet. Genom.* **2014**, *290*, 661–670. [CrossRef]
9. Hartmann, U.; Höhmann, S.; Nettesheim, K.; Wisman, E.; Saedler, H.; Huijser, P. Molecular Cloning of *SVP*: A Negative Regulator of the Floral Transition in Arabidopsis. *Plant J.* **2000**, *21*, 351–360. [CrossRef] [PubMed]
10. Gregis, V.; Sessa, A.; Colombo, L.; Kater, M.M. AGAMOUS-LIKE24 and SHORT VEGETATIVE PHASE Determine Floral Meristem Identity in *Arabidopsis*. *Plant J.* **2008**, *56*, 891–902. [CrossRef]
11. Li, D.; Liu, C.; Shen, L.; Wu, Y.; Chen, H.; Robertson, M.; Helliwell, C.A.; Ito, T.; Meyerowitz, E.; Yu, H. A Repressor Complex Governs the Integration of Flowering Signals in *Arabidopsis*. *Dev. Cell* **2008**, *15*, 110–120. [CrossRef]
12. Lee, J.H.; Yoo, S.J.; Park, S.H.; Hwang, I.; Lee, J.S.; Ahn, J.H. Role of *SVP* in the Control of Flowering Time by Ambient Temperature in *Arabidopsis*. *Genes. Dev.* **2007**, *21*, 397–402. [CrossRef]
13. Jaudal, M.; Monash, J.; Zhang, L.; Wen, J.; Mysore, K.S.; Macknight, R.; Putterill, J. Overexpression of Medicago *SVP* Genes Causes Floral Defects and Delayed Flowering in Arabidopsis but Only Affects Floral Development in Medicago. *J. Exp. Bot.* **2013**, *65*, 429–442. [CrossRef] [PubMed]
14. Higo, K.; Ugawa, Y.; Iwamoto, M.; Korenaga, T. Plant Cis-Acting Regulatory DNA Elements (PLACE) Database: 1999. *Nucleic Acids Res.* **1999**, *27*, 297–300. [CrossRef] [PubMed]
15. Curtis, M.D.; Grossniklaus, U. A Gateway Cloning Vector Set for High-Throughput Functional Analysis of Genes in Planta. *Plant Physiol.* **2003**, *133*, 462–469. [CrossRef]
16. Clough, S.J.; Bent, A.F. Floral Dip: A Simplified Method for *Agrobacterium*-Mediated Transformation of *Arabidopsis thaliana*. *Plant J.* **1998**, *16*, 735–743. [CrossRef] [PubMed]

17. Koressaar, T.; Remm, M. Enhancements and Modifications of Primer Design Program Primer3. *Bioinformatics* **2007**, *23*, 1289–1291. [CrossRef]
18. Untergasser, A.; Cutcutache, I.; Koressaar, T.; Ye, J.; Faircloth, B.C.; Remm, M.; Rozen, S.G. Primer3—New Capabilities and Interfaces. *Nucleic Acids Res.* **2012**, *40*, e115. [CrossRef]
19. Livak, K.J.; Schmittgen, T.D. Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the $2^{-\Delta\Delta CT}$ Method. *Methods* **2001**, *25*, 402–408. [CrossRef]
20. Huang, S.; Raman, A.S.; Ream, J.E.; Fujiwara, H.; Cerny, R.E.; Brown, S.M. Overexpression of 20-Oxidase Confers a Gibberellin-Overproduction Phenotype in *Arabidopsis*. *Plant Physiol.* **1998**, *118*, 773–781. [CrossRef]
21. Rieu, I.; Ruiz-Rivero, O.; Fernandez-Garcia, N.; Griffiths, J.; Powers, S.J.; Gong, F.; Linhartova, T.; Eriksson, S.; Nilsson, O.; Thomas, S.G.; et al. The Gibberellin Biosynthetic Genes *AtGA20ox1* and *AtGA20ox2* Act, Partially Redundantly, to Promote Growth and Development throughout the *Arabidopsis* Life Cycle. *Plant J.* **2007**, *53*, 488–504. [CrossRef]
22. Liu, C.; Xi, W.; Shen, L.; Tan, C.; Yu, H. Regulation of Floral Patterning by Flowering Time Genes. *Dev. Cell* **2009**, *16*, 711–722. [CrossRef]
23. Wikström, N.; Savolainen, V.; Chase, M.W. Evolution of the Angiosperms: Calibrating the Family Tree. *Proc. R. Soc. B Biol. Sci.* **2001**, *268*, 2211–2220. [CrossRef]
24. Blanc, G.; Hokamp, K.; Wolfe, K.H. A Recent Polyploidy Superimposed on Older Large-Scale Duplications in the *Arabidopsis* Genome. *Genome Res.* **2003**, *13*, 137–144. [CrossRef] [PubMed]
25. Zhu, Y.; Wagner, D. Plant Inflorescence Architecture: The Formation, Activity, and Fate of Axillary Meristems. *Cold Spring Harb. Perspect. Biol.* **2020**, *12*, a034652. [CrossRef] [PubMed]
26. Evers, J.B.; van der Krol, A.R.; Vos, J.; Struik, P.C. Understanding Shoot Branching by Modelling Form and Function. *Trends Plant Sci.* **2011**, *16*, 464–467. [CrossRef] [PubMed]
27. Charlesworth, B.; Charlesworth, D. A Model for the Evolution of Dioecy and Gynodioecy. *Am. Nat.* **1978**, *112*, 975–997. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.