

***Tiliacora triandra* Leaf Powder Ethanolic Extract in Combination with Cisplatin or Gemcitabine Synergistically Inhibits the Growth of Cholangiocarcinoma Cells In Vitro and in Nude Mouse Xenograft Models**

Arunta Samankul ¹, Gulsiri Senawong ¹, Suppawit Utaiwat ¹, Jeerati Prompipak ¹, Khanutsanan Woranam ¹, Chanokbhorn Phaosiri ², Banchob Sripa ³ and Thanaset Senawong ^{1,*}

¹ Department of Biochemistry, Faculty of Science, Khon Kaen University, Khon Kaen 40002, Thailand; s_arunta@kkumail.com (A.S.); gulsiri@kku.ac.th (G.S.); u.suppawit@kkumail.com (S.U.); jeerati.ppk@kkumail.com (J.P.); khanutsanan_w@kkumail.com (K.W.)

² Department of Chemistry, Faculty of Science, Khon Kaen University, Khon Kaen 40002, Thailand; chapha@kku.ac.th

³ WHO Collaborating Centre for Research and Control of Opisthorchiasis (Southeast Asian Liver Fluke Disease), Tropical Disease Research Center, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002, Thailand; banchob@kku.ac.th

* Correspondence: sthanaset@kku.ac.th




Figure S1. Antiproliferative effects of Gemcitabine against CCA cells at exposure time of 24 h. KKU-M213B (a), KKU-100 (b), and H69 (c) cells were treated with Gem for 24 h at the indicated concentrations.