

Article

Effect of Carboxymethylation and Phosphorylation on the Properties of Polysaccharides from *Sepia esculenta* Ink: Antioxidation and Anticoagulation in Vitro

Huazhong Liu, Fangping Li and Ping Luo *

College of Chemistry & Environment, Guangdong Ocean University, Zhanjiang 524088, China; liuhzbs@163.com (H.L.) and 15709482571@163.com (F.L.)

* Correspondence: luopingna@163.com; Tel.: +86 759 2383300

Received: 15 October 2019; Accepted: date; Published: 1 November 2019

Abstract: To investigate the effect of carboxymethylation and phosphorylation modification on *Sepia esculenta* ink polysaccharide (SIP) properties, this study prepared carboxymethyl SIP (CSIP) with the chloroacetic acid method, and phosphorylated SIP (PSIP) with the sodium trimetaphosphate (STMP)/sodium tripolyphosphate (STPP) method, on the basis of an orthogonal experiment. The in vitro antioxidant and anticoagulant activities of the derivatives were determined by assessing the scavenging capacity of the 1,1-diphenyl-2-picrylhydrazyl (DPPH) and hydroxyl radicals, which activated the partial thromboplastin time (APTT), prothrombin time (PT), and thrombin time (TT). The results showed that SIP was modified successfully to be CSIP and PSIP, and degrees of substitution (DSs) of the two products were 0.9913 and 0.0828, respectively. Phosphorylation efficiently improved the antioxidant property of SIP, and the IC₅₀ values of PSIP on DPPH and hydroxyl radicals decreased by 63.25% and 13.77%, respectively. But carboxymethylation reduced antioxidant activity of the native polysaccharide, IC₅₀ values of CSIP on the DPPH and hydroxyl radicals increased by 16.74% and 6.89%, respectively. SIP significantly prolonged the APTT, PT, and TT in a dose-dependent fashion, suggesting that SIP played an anticoagulant action through intrinsic, extrinsic, and common coagulation pathways. CSIP and PSIP both possessed a stronger anticoagulant capacity than SIP via the same pathways; moreover, CSIP was observed to be more effective in prolonging APTT and PT than PSIP.

Keywords: *Sepia esculenta* ink polysaccharide; carboxymethylation; phosphorylation; antioxidation; anticoagulation

1. Introduction

Sepia esculenta ink polysaccharides (SIP) is a type of glycosaminoglycans derived from *Sepia esculenta* ink that has proven to be multifunctional in marine materials, displaying properties such as antioxidation, antitumor, immunoregulation, and chemoprevention [1,2]. Our previous work reported that SIP derived from *Sepia esculenta* ink is mainly composed of galactosamine and arabinose, as well as a small amount of fucose and other monosaccharides [3]. As SIP has been found to have properties of chemoprevention, antitumor, and chemosensitization, the natural polysaccharide was recognized as a potential adjuvant agent of chemotherapy [1]. It is well known that chemotherapy induces blood hypercoagulability, and the consequent thrombus formation is a critical cause of death in cancer cases [1]; therefore, whether SIP mediates blood coagulation is an unavoidable, crucial issue for developing the marine bioactive substance to be an adjuvant drug.

Presently, no direct evidence shows an anticoagulation or procoagulation property of SIP. Sulphated SIP, a type of derivative of SIP isolated from *Ommastrephes bartrami* ink prepared by Chen et al., was found to possess an efficient anticoagulant activity, but the native SIP failed to alter the blood clotting time in vitro [4], suggesting that sulphation is an effective modification method for improving the property of SIP.

Chemical modification is well-recognized to be an effective way to change the physiochemical properties and to improve the biological activities of natural polysaccharides, which result from the alternation of the molecular structure caused by the conjugated functional groups [5]. It is a widely accepted notion that polysaccharide derivatives have a superior water solubility and antioxidant activity to native polysaccharides. Many modification methods have been widely used to modify native polysaccharides for improving properties, such as sulfation, methylation, hydroxypropylation, carboxymethylation, acetylation, phosphorylation, alkylation, and sulfonylation [6].

Hitherto, sulphated SIP is the only derivative of SIP, the native polysaccharide is derived from *Ommastrephes bartrami* ink. The modified polysaccharide shows anticoagulation, anti-angiogenesis, and anti-metastasis activities of HepG2 in tumor cells [4,7]. Another sulfated SIP prepared by sulfation on SIP isolated from *Sepiella maindroni* ink is anti-angiogenesis, as well as the anti-metastasis activities of SKOV3 tumor cells [8–10]. At present, no other derivatives of SIP have been reported, besides the sulphated SIP. To investigate the effect of the chemical modification on the SIP property, this study prepared carboxymethylated SIP (CSIP) and phosphorylated SIP (PSIP) to assess the effect of carboxymethylation and phosphorylation on SIP, based on antioxidation and anticoagulation.

2. Results

2.1. Determination of Carboxymethylation Parameters and Molecular Characteristics of CSIP

The data presented in Figure 1 show that the four scheduled influence factors clearly affected the degree of substitution (DS) of the CSIP. The optimal factors were 3.0 mol/L (NaOH), 2.4 g (chloroacetic acid), 40 °C (reaction temperature), and 3 h (reaction time), respectively. The optimal combined parameters were 5.0 mol/L of NaOH solution, 2.84 g of chloroacetic acid, 50 °C, and 2.5 h, according to the values of *R* and DS (Table 1). The DS of the CSIP was calculated to be 0.9913. Different from the spectrum of the SIP, the CSIP spectrum presented two strong absorption peaks at 1605 and 1419 cm^{-1} , two characteristic absorption bands of carboxymethyl of CSIP, suggesting that the SIP was carboxymethylated successfully (Figure 2).




Figure 1. Effect of different reaction conditions on degree of substitution (DS) of carboxymethylated *Sepia esculenta* ink polysaccharides (CSIP). Under the scheduled different reaction conditions, the SIP was carboxymethylated by chloroacetic acid. The degree of substitution of CSIP was determined by the neutralization titration method and calculation according to the formula.

Table 1. Orthogonal experimental design and results of the SIP carboxymethylation.

Items	Chloroacetic acid (g)	NaOH (mol/L)	Temperature (°C)	Time (h)	DS
1	1.89	1.0	30	2.5	0.43
2	1.89	3.0	40	3.0	0.60
3	1.89	5.0	50	3.5	0.58
4	2.37	1.0	40	3.5	0.45
5	2.37	3.0	50	2.5	0.75
6	2.37	5.0	30	3.0	0.75
7	2.84	1.0	50	3.0	0.61
8	2.84	3.0	30	3.5	0.67
9	2.84	5.0	40	2.5	0.84
K_1	0.49	0.53	0.61	0.67	
K_2	0.67	0.65	0.63	0.65	
K_3	0.72	0.70	0.64	0.56	
R	0.23	0.17	0.03	0.11	




Figure 2. IR spectra of SIP and CSIP. IR spectra of the SIP and CSIP were recorded with KBr pellets on a Bruker Tensor 27 Fourier infrared spectrophotometer between 400–4000 cm^{-1} .

2.2. Determination of the Phosphorylation Parameters and Molecular Characteristics of PSIP

A single factor experiment revealed that the optimal ratio of the sodium tripolyphosphate (STPP)/sodium trimetaphosphate (STMP), reaction time, temperature, and pH value was determined to be 50/20 (Table 2), 3.0 h, 40 °C, and pH 6.0, respectively, which were subjected to the orthogonal experiment (Figure 3). The optimal combined parameters were determined to be 40/30 (STPP/STMP), 3.0 h, 55 °C, and pH 5.0, respectively, according to the values of *R* and *DS* (Table 3). The phosphate content of PSIP was 1.52%, and the *DS* was calculated to be 0.0828. Compared with the spectrum of SIP, the PSIP spectrum observed two distinct absorption peaks at 1019 cm^{-1} (P–OH bond) and 889 cm^{-1} (P–O–C bond), suggesting that SIP was phosphorylated successfully (Figure 4).

Table 2. Effect of the phosphorylating reagent on the content of phosphate. STPP—sodium tripolyphosphate; STMP—sodium trimetaphosphate.

Items	STPP (g/L)	STMP (g/L)	Phosphate radical content (%)
1	0	0	0.305
2	70	0	1.451
3	60	10	2.095
4	50	20	3.818
5	40	30	3.661
6	30	40	2.826
7	20	50	3.081
8	10	60	1.921
9	0	70	2.269





Figure 3. Effect of different reaction conditions on the content of phosphate in phosphorylated SIP (PSIP). Under the scheduled different reaction conditions, the SIP was phosphorylated with the STMP/STPP method. The content of phosphate in PSIP was determined according to the equation, which was subjected to calculating the degree of substitution of PSIP.


Table 3. Orthogonal experimental design and results of SIP phosphorylation.

Items	STPP/STMP (g/L)	Temperature (°C)	Time (h)	pH	Phosphate content (%)
1	40/30	55	3	5.0	1.52
2	40/30	65	4	6.0	0.33
3	40/30	75	5	7.0	0.63
4	50/20	55	4	7.0	0.20
5	50/20	65	5	5.0	0.76
6	50/20	75	3	6.0	0.39
7	60/10	55	5	6.0	0.52
8	60/10	65	3	7.0	0.55
9	60/10	75	4	5.0	0.25
K_1	0.82	0.74	0.82	0.84	
K_2	0.45	0.54	0.26	0.41	
K_3	0.44	0.42	0.63	0.46	
R	0.38	0.32	0.56	0.43	

**Figure 4.** IR spectra of SIP and PSIP. The IR spectra of SIP and PSIP were recorded with KBr pellets on a BRUKER TENSOR 27 Fourier infrared spectrophotometer between 400–4000 cm^{-1} .


2.3. Antioxidation Activity of CSIP and PSIP

Figure 5 presents that the scavenging activity of SIP on the 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical was stronger than CSIP, but weaker than PSIP, which was demonstrated by the modified IC_{50} values—the PSIP dropped by 63.25% and the CSIP increased by 16.74%. Similarly, with respect to the scavenging capacity of SIP on the hydroxyl radical, the CSIP was weaker and the PSIP was stronger, and the IC_{50} values of the CSIP and PSIP increased by 6.89% and decreased by 13.77%, respectively. The data suggest that phosphorylation promoted the *in vitro* antioxidation activity of SIP, but carboxymethylation inhibited the property of SIP.

**Figure 5.** *In vitro* antioxidant activity of SIP, CSIP, and PSIP. The antioxidant capacity of SIP and its derivatives was assessed by the scavenging activities of DPPH and hydroxyl radicals.

2.4. Anticoagulation Activity of CSIP and PSIP

The activated partial thromboplastin time (APTT), prothrombin time (PT), and thrombin time (TT), three indicators evaluating the in vitro anticoagulation activity of the anticoagulant, express intrinsic, extrinsic, and common pathways of blood coagulation, respectively. The data in Figure 6 show that SIP prolonged the coagulation time in a dose-dependent manner, including APTT, PT, and TT, indicating that SIP had an anticoagulation action via endogenous, exogenous, and common coagulation pathways. Meanwhile, CSIP and PSIP both were observed in the capacity of extending the APTT, PT, and TT in dose-dependent fashions. Importantly, compared with SIP, both CSIP and PSIP showed a significantly stronger anticoagulant activity that was demonstrated by an extended APTT, PT, and TT. Moreover, CSIP was observed to have a stronger anticoagulant capacity than PSIP, ranging from 6.25 to 25 µg/mL, which was indicated by APTT and PT. These data indicate that SIP and its derivatives CSIP and PSIP, performed an anticoagulation action via intrinsic, extrinsic, and common pathways of blood coagulation. Moreover, both carboxymethylation and phosphorylation promoted the activity of the native polysaccharide, and the promotion of carboxymethylation was slightly superior to the phosphorylation modification.






Figure 6. In vitro anticoagulant property of SIP, CSIP, and PSIP. The anticoagulant capacity of SIP and its derivatives was assessed by determining the clotting time, including the activated partial thromboplastin time (APTT), prothrombin time (PT), and thrombin time (TT). Different capital or lowercase letters express significant differences among various dosages (0, 6.25, 12.5, 25, 50, and 100) of polysaccharide (SIP, CSIP, or PSIP), $_{ABCDE} p < 0.01$ or $_{abcDEF} p < 0.05$. Asterisk, * or **, means $p < 0.05$ or $p < 0.01$ between the same dosage of SIP and modified SIP (CSIP or PSIP), respectively. # or ## represent the difference, $p < 0.05$ or $p < 0.01$, between the same dosage of CSIP and PSIP, respectively.

3. Discussion

Carboxymethylation and phosphorylation have been widely used to modify natural polysaccharides for improving the physicochemical and biological properties of the native macromolecules [11]. In the present study, SIP was modified with chloroacetic acid and STMP/STPP methods, respectively. Through the chemical modification methods, SIP was successfully modified to be CSIP and PSIP, and their DSs were determined to be 0.9913 and 0.0828, respectively. DS is an important influencing factor of the biological activity of the polysaccharide derivative, indicating that CSIP and PSIP may possess different properties from SIP.

Currently, three types of SIPs have been reported, including polysaccharides from *Ommastrephes bartrami* ink, *Illex argentinus* ink, *Sepiella maindroni* ink, and *Sepia esculenta* ink. The first two SIPs share the same primary structure [1]. No evidence indicated a group of carboxymethyl or phosphate in SIP. The SIP used in this study was derived from *Sepia esculenta* ink, and was mainly composed of galactosamine and arabinose (molar ratio of 1:1), accounting for more than 81% of monosaccharides. In addition, fucose, xylose, glucuronic acid, galacturonic acid, glucosamine, and mannose were about 9.00%, 4.32%, 1.98%, 1.53%, 1.35%, and 0.09%, respectively [3]. Both the carboxymethylation and phosphorylation reactions occur on the hydroxyl group of polysaccharides, therefore, it is still unable to confirm which sites are substituted by carboxymethyl or phosphate until the primary structures of SIP, CSIP, and PSIP have been characterized.

The carboxymethyl group is an electron-withdrawing group that can influence the antioxidant capacity of heterocyclic compounds, by increasing or decreasing [12,13]. The opposite effect of carboxymethylation on natural polysaccharides has been proven by many reports [11,12,14–18]. This work exhibited a negative effect of carboxymethylation on the SIP antioxidant property. As the molecular structure of the SIP used in this study is still unknown, the involved mechanism of carboxymethylation on the antioxidant ability of SIP is unable to be explained.

Phosphorylation improving the antioxidant activity of polysaccharides result from the conjugated phosphate groups via ester linkage [19], which is partly attributed to phosphate DS, a critical influencing factor [20]. Phosphoryl polysaccharide generally possesses a low DS, and a stronger bioactivity than the sulfated derivative [21–23]. Our data revealed that the DS of PSIP was 0.0828. The DS of phosphorylated polysaccharide has a close relationship with the phosphorylation methods. Many methods have been developed in order to prepare phosphorylated polysaccharides [24,25]. The present work prepared PSIP with the STMP method. The mechanism involved in the

synthesis of PSIP is still unclear, but it can probably be deduced according to the findings [25]. STMP is a cross-linking agent that triggers a reaction between the hydroxyl groups of SIPs and the metaphosphate groups in STMP, resulting in phosphate ester (O–P–O) linkages between two polysaccharide moieties.

Three negative charges of the phosphate radical result in an increase of electronegativity and activity alternation of the polysaccharide. The phosphorylation modification improved the in vitro antioxidant activity of the SIP, demonstrated by promoted the scavenging capacity of the DPPH and hydroxyl radicals. Similar results were found in garlic polysaccharide [6], *Enteromorpha linza* polysaccharide [26], *Dictyophora indusiata* polysaccharide [24], *Portulaca oleracea* L. polysaccharide [20], pumpkin polysaccharide [27], and native ginseng polysaccharide [28].

The SIP from *Ommastrephes bartrami* ink is ineffective at changing the blood coagulation time in vitro [7]. However, the SIP from *Sepia esculenta* ink was proven to have an obvious anticoagulant activity in this work. This type of SIP is a kind of sulfate polysaccharide with a 2.25% sulfate content, but the sulfate was not detected in the SIP from *Ommastrephes bartrami* ink. Generally, the anticoagulant property of the sulfate polysaccharide is related to the sulfate content [29]. Moreover, the sulfated SIP (molar ratio of monosaccharide/sulfate group: 3/2) markedly extended the APTT and PT via inhibiting the blood coagulation factors IIa and Xa [7]. So, one important cause of the anticoagulant property should be the existence of sulfate in SIP used in this study. Another crucial cause may be the molecular structure—the two types of SIPs possess entirely different primary structures.

It is widely accepted that chemical modification effectively improves the physicochemical and biological properties of natural polysaccharides. This work testified that carboxymethylation and phosphorylation high-effectively improved the anticoagulant activity of SIP, indicated by prolonged APTT, PT, and TT. Interestingly, carboxymethylation was more effective in improving the anticoagulant capacity of SIP than phosphorylation.

Various phosphorylated polyanions play an anticoagulant action through enhancing the complexation of heparin cofactor II and thrombin [30], the binding is mainly related to the negative charges and the polymeric nature [31]. The phosphorylation increasing anticoagulant property has been found in oat spelt's xylan [29], which were related to the molecular weight (positively) and phosphate content (inversely) [31], as well as the highly polymeric nature, and phosphodiester or diphosphodiester bonds [29].

Currently, there is little understanding of the molecular structure of the novel SIP and its derivatives, CSIP and PSIP, and the structure-activity relationship needs to be further investigated in order to explain the involved regulatory mechanisms of carboxymethylation, phosphorylation on antioxidation, and the anticoagulation of SIP.

4. Materials and Methods

4.1. Preparation of SIP

According to the reported method of Gu et al. [32], the procedure is briefly described as follows. Fresh *Sepia esculenta* ink stored at $-70\text{ }^{\circ}\text{C}$ was thawed at $4\text{ }^{\circ}\text{C}$, diluted in phosphate buffer solution, and then ultrasonicated. After storage at $4\text{ }^{\circ}\text{C}$ for more than 8 h and then centrifugation (8000 rpm) at $4\text{ }^{\circ}\text{C}$, the supernatant was hydrolyzed with papain for 90 min, and then heated for 1 h in a boiling water bath. Following centrifugation, four volumes of ethanol were used to precipitate the crude polysaccharides that were then subjected to DEAE-52 cellulose column chromatography. The first fraction containing SIP was collected and dialyzed, desiccated in a vacuum freeze drier, and stored at $-20\text{ }^{\circ}\text{C}$.

4.2. Preparation of CSIP

An orthogonal experimental design was used to prepare the CSIP. The SIP (100 mg) was dissolved in 20 mL of NaOH solution (1.0, 3.0, 5.0, 7.0, and 9.0 mol/L), and stirred for 30 min at room temperature. Chloroacetic acid (0.95, 1.42, 1.89, 2.37 and 2.84 g) was dissolved in a SIP solution and

heated in a water bath (20, 30, 40, 50, and 60 °C) for a scheduled time (2.0, 2.5, 3.0, 3.5, and 4.0 h). The reaction system was cooled to room temperature and neutralized to pH 7.0 with glacial acetic acid. After dialysis against distilled water for 72 h, the solution was dried at 75 °C in a drier to harvest the CSIP.

The DS was determined according to the method of neutralization titration, with slight modification [33]. The CSIP sample (10 mg) dissolved in 3 ml of methanol (70%) was added in 15 ml of NaOH solution (0.167 mol/L). Immediately, the mixture was titrated with an HCl solution (0.1 mol/L). The DS was calculated by the following formulas:

$$A = (V_0M_0 - VM) / W,$$

$$DS = 0.162 A / (1 - 0.058A),$$

where $V_0 = 5$ mL; $M_0 = 0.5$ mol/L; $M = 0.1$ mol/L; and V and W represent the consumed volume of the hydrochloric acid (mL) and sample weight (g), respectively.

4.3. Preparation of PSIP

An orthogonal experimental design was used to prepare the PSIP. The SIP solution (10 mL) in distilled water (10 mg/mL) was added in 4.285 mL of a mixed solution (70 mg/mL) containing STPP and STMP, and 0.2 mL of NaSO₄ solution (50 mg/mL). The contents of STPP and STMP in the mixed solution are presented in Table 2. The reaction system was neutralized to scheduled pH values (5.0, 6.0, 7.0, 8.0, and 9.0) with HCl (1.0 mol/L), and cooled to scheduled temperatures (25, 45, 65, 85, and 100 °C). After reaction for a scheduled time (2.0, 3.0, 4.0, 5.0, and 6.0 h), three volumes of ethanol were used to precipitate the product for 24 h at room temperature. The precipitate was dried at 75 °C in a drier, and then dissolved in 15 mL of distilled water at 50 °C in a water bath. The conductivity determination of the solution occurred in the process of dialysis against distilled water. The dialysis was stopped when the conductivity declined to be 160 µs/cm. The solution was dried at 75 °C in a drier so as to harvest PSIP.

The phosphate content was determined according to the reported method, with slight modification; the standard curve of the phosphate radical was prepared according to the procedure [34]. Different volumes (0, 0.5, 1.0, 1.5, 2.0, 2.5, 3.0, 3.5, 4.0, 4.5, and 5.0 mL) of a phosphate standard solution (0.1 mg/mL) were mixed with distilled water, respectively, and the total volume reached 5.0 mL. Tris buffer solution (3.0 mL, 0.1 mol/L, pH 7.0) was added into 3 ml of phosphate quantitative reagent, reacting for 30 min at a temperature of 45 °C. Then, the absorbance was read at a wavelength of 580 nm. The standard curve of the phosphate radical content was established with the absorbance (ordinate) and phosphate radical content (abscissa). The phosphate content in the PSIP was determined according to the method in the literature [35]. The dried PSIP (10 mg) was ashed for 2 h at 600 °C, and then dissolved in 0.5 mL of HCl solution (0.5 mol/L), and diluted with distilled water to 10 mL. The ashed compound solution (1.0 mL) was put into a 25 mL cuvette, and the absorbance was read according to the abovementioned standard curve method. Based on the standard curve, phosphate radical content in the PSIP was calculated.

4.4. Infrared Spectral Analysis

The IR spectra of the SIP, CSIP, and PSIP were recorded with KBr pellets on a Bruker Tensor 27 Fourier infrared spectrophotometer (Bruker, Karlsruhe, Germany) between 400–4000 cm⁻¹.

4.5. Scavenging Activity of Hydroxyl Radical

The scavenging activity of the hydroxyl radical was assessed according to the method in the literature, with slight modification [33]. A sample (SIP, CSIP, or PSIP) was dissolved in distilled water, to be different concentrations, namely, 0.5, 1.0, 2.0, 4.0, and 8.0 g/L. The sample solution (1.0 mL) was diluted in 2.0 mL of a phosphate buffer solution (0.02 mol/L, pH 7.4), and was mixed with 1,10-phenanthroline (1.5 mL, 1.0 mmol/L), FeSO₄ (1.0 mL, 1.5 mmol/L), H₂O₂ (1.0 mL, 1.0%), and

distilled water (3.5 mL). Following incubation for 60 min at 37 °C, the absorbance was read at a wavelength of 536 nm. The scavenging activity was calculated according to the following formula:

$$\text{Scavenging activity (\%)} = \frac{\text{OD}_2 - \text{OD}_1}{\text{OD}_0 - \text{OD}_1} \times 100\%$$

where the sample was replaced by H₂O in the OD₁ associated reaction system, H₂O replaced the sample and H₂O₂ in the OD₀ associated reaction system, and OD₂ associated with the reaction system contained the sample.

4.6. Scavenging Activity of DPPH

The scavenging activity of the DPPH radical was determined according to the method from the literature, with slight modification [33]. A DPPH solution (2.0 mL, 0.1 mmol/L) was mixed with 0.5 mL of different concentrations of SIP, CSIP, or PSIP (0.5, 1.0, 2.0, 4.0, and 8.0 g/L), and 1.5 mL of distilled water, and was then kept away from light for 30 min at room temperature. The absorbance was read at a wavelength of 517 nm. The scavenging activity was calculated as the following formula.

$$\text{Scavenging activity (\%)} = \frac{1 - (\text{OD}_2 - \text{OD}_1)}{\text{OD}_0} \times 100\%$$

where OD₀ represents the negative control that is composed by 2.0 mL DPPH and 2.0 mL ethanol. OD₁ is a mixture of 2.0 mL ethanol, 0.5 mL of the sample solution, and 1.5 mL distilled water. OD₂ consists of 2.0 mL of a DPPH–ethanol solution, 0.5 mL of a sample solution, and 1.5 mL distilled water.

4.7. Determination of the Anticoagulation Activity

Male Kunming mice (20–25 g) purchased from Guangdong Medical Animal Experiment Center (SCKY-2018-0002) were habituated for 1 week and were randomly allocated to six groups, control group and five treated groups, ten mice in each group. Mice were sacrificed to harvest blood that was then prepared serum immediately. Each group of murine sera was treated with distilled water or different dosages (6.25, 12.5, 25, 50 and 100 µg/mL) of SIP, CSIP or PSIP solution in distilled water and then subjected to determine APTT, PT and TT.

The activated partial thromboplastin time (APTT), prothrombin time (PT) and thrombin time (TT) of the SIP, CSIP, and PSIP were assessed with kits, purchased from Nanjing Jiancheng Bioengineering Institute (Nanjing, China), according to the manufacturer's protocols.

4.8. Statistical Analysis

The data were expressed as mean ± the standard error, and were analyzed using SPSS statistical software. One-way analysis of variance (ANOVA) of the data was performed. *p* < 0.05 was considered to be a significant level.

5. Conclusions

In summary, the SIP was successfully modified to be CSIP with the chloracetic acid method, and to be PSIP with the STMP/STPP method, on the basis of the orthogonal experiment. The DSs of the two products were 0.9913 and 0.0828, respectively. The results revealed that the phosphorylation efficiently improved the antioxidant property of the SIP, but carboxymethylation reduced the activity of the native polysaccharide. Additionally, we found a strong in vitro anticoagulant action of SIP from *Sepia esculenta* ink via intrinsic, extrinsic, and common coagulation pathways, which was different from the SIP from *Ommastrephes bartrami* ink that did not observe an anticoagulant activity [4]. Moreover, the phosphorylation and carboxymethylation both could enhance the anticoagulant capacity of SIP, and CSIP showed a more effective anticoagulation action than PSIP in intrinsic and extrinsic pathways, but no obvious difference between them was observed in the common pathway.

Author Contributions: All of the authors were involved in this research, approved the final version to be

published and had full access to all of the data in the study, and took responsibility for the integrity of the data and the accuracy of the data analysis. H.L. and P.L. conceived and designed this study, and drafted or revised the article. H.L. and F.L. acquired and interpreted the data.

Funding: This work was supported by the Natural Science Foundation of Guangdong Province, China (2016A030313753), and the Science and Technology Project on Special Fund for Public Welfare Research and Ability Construction of Guangdong Province, China (2017A010105010).

Conflicts of interest: The authors declare no competing interest.

References

- Li, F.P.; Luo, P.; Liu, H.Z. A potential adjuvant agent of chemotherapy: sepia ink polysaccharide. *Mar. Drugs* **2018**, *16*, 106.
- Liu, H.Z.; Zhang, Y.B.; Li, M.W.; Luo, P. Beneficial effect of *Sepia esculenta* ink polysaccharide on cyclophosphamide-induced immunosuppression and ovarian failure in mice. *Int. J. Biol. Macromol.* **2019**, *140*, 1098–1105.
- Liu, H.Z.; Tao, Y.X.; Luo, P.; Deng, C.M.; Gu, Y.P.; Yang, L.; Zhong, J.P. Preventive effects of a novel polysaccharide from *Sepia esculenta* ink on ovarian failure and its action mechanisms in cyclophosphamide-treated mice. *J. Agric. Food Chem.* **2016**, *64*, 5759–5766.
- Chen, S.G.; Li, Z.G.; Wang, Y.M.; Li, G.Y.; Sun, B.B.; Xue, C.H. Sulfation of a squid ink polysaccharide and its anticoagulant activities. *Chem. J. Chin. Univ.* **2010**, *31*, 2407–2412.
- Bohn, J.A.; Be Miller, J.N. (1 → 3)-β-D-Glucan as biological response modifiers: A review of structure–functional activity relationships. *Carbohydr. Polym.* **1995**, *28*, 3–14.
- Chen, J.; Huang, G. Antioxidant activities of garlic polysaccharide and its phosphorylated derivative. *Int. J. Biol. Macromol.* **2019**, *125*, 432–435.
- Chen, S.G.; Wang, J.F.; Xue, C.H.; Li, H.; Sun, B.B.; Xue, Y.; Chai, W.G. Sulfation of a squid ink polysaccharide and its inhibitory effect on tumor cell metastasis. *Carbohydr. Polym.* **2010**, *81*, 560–566.
- Wang, S.; Cheng, Y.; Wang, F.; Sun, L.; Liu, C.; Chen, G.; Li, Y.; Ward, S.G.; Qu, X. Inhibition activity of sulfated polysaccharide of *Sepiella maindroni* ink on matrix metalloproteinase (MMP)-2. *Biomed. Pharmacother.* **2008**, *62*, 297–302.
- Zong, A.Z.; Zhao, T.; Zhang, Y.; Song, X.L.; Shi, Y.K.; Cao, H.Z.; Liu, C.H.; Cheng, Y.N.; Qu, X.J.; Cao, J.C.; Wang, F.S. Anti-metastatic and anti-angiogenic activities of sulfated polysaccharide of *Sepiella maindroni* ink. *Carbohydr. Polym.* **2013**, *91*, 403–409.
- Zong, A.Z.; Liu, Y.H.; Zhang, Y.; Song, X.; Shi, Y.K.; Cao, H.Z.; Liu, C.H.; Cheng, Y.; Jiang, W.J.; Du, F.L.; Wang, F.S. Anti-tumor activity and the mechanism of SIP-S: a sulfated polysaccharide with anti-metastatic effect. *Carbohydr. Polym.* **2015**, *129*, 50–54.
- Wang, Z.J.; Xie, J.H.; Shen, M.Y.; Tang, W.; Wang, H.; Nie, S.P.; Xie, M.Y. Carboxymethylation of polysaccharide from *Cyclocarya paliurus* and their characterization and antioxidant properties evaluation. *Carbohydr. Polym.* **2016**, *136*, 988–994.
- Kagimura, F.Y.; da Cunha, M.A.A.; Barbosa, A.M.; Dekker, R.F.H.; Malfatti, C.R.M. Biological activities of derivatized D-glucans: a review. *Int. J. Biol. Macromol.* **2015**, *72*, 588–598.
- Yanagimoto, K.; Lee, K.G.; Ochi, H.; Shibamoto, T. Antioxidative activity of heterocyclic compounds found in coffee volatiles produced by Maillard reaction. *J. Agric. Food Chem.* **2002**, *50*, 5480–5484.
- Xu, J.; Liu, W.; Yao, W.; Pang, X.; Yin, D.; Gao, X. Carboxymethylation of a polysaccharide extracted from *Ganoderma lucidum* enhances its antioxidant activities *in vitro*. *Carbohydr. Polym.* **2009**, *78*, 227–234.
- Yang, L.; Zhao, T.; Wei, H.; Zhang, M.; Zou, Y.; Mao, G.; Wu, X. Carboxymethylation of polysaccharides from *Auricularia auricula* and their antioxidant activities *in vitro*. *Int. J. Biol. Macromol.* **2011**, *49*, 1124–1130.
- Chen, Y.; Zhang, H.; Wang, Y.; Nie, S.; Li, C.; Xie, M. Acetylation and carboxymethylation of the polysaccharide from *Ganoderma atrum* and their antioxidant and immunomodulating activities. *Food Chem.* **2014**, *156*, 279–288.
- Wang, X.; Zhang, Z.; Zhao, M. Carboxymethylation of polysaccharides from *Tremella fuciformis* for antioxidant and moisture-preserving activities. *Int. J. Biol. Macromol.* **2015**, *72*, 526–530.
- Li, J.; Shang, W.; Si, X.; Bu, D.; Strappe, P.; Zhou, Z.; Blanchard, C. Carboxymethylation of corn bran polysaccharide and its bioactive property. *Int. J. Food Sci. Tech.* **2017**, *52*, 1176–1184.
- Wei, D.; Cheng, W.; Wei, Y.; Zhang, L. Phosphorylated modification and *in vitro* antioxidant activity of *Radix Hedysari* polysaccharide. *Glycoconj. J.* **2012**, *29*, 167–172.

20. Chen, T.; Zhu, L.; Liu, X.; Li, Y.; Zhao, C.; Xu, Z.; Yan, W.; Zhang, H. Synthesis and antioxidant activity of phosphorylated polysaccharide from *Portulaca oleracea* L. with H₃PW₁₂O₄₀ immobilized on polyamine functionalized polystyrene bead as catalyst. *J. Mol. Catal. Chem.* **2011**, *342*, 74–82.
21. Liu, X.X.; Wan, Z.J.; Shu, L.; Lu, X.X. Preparation and antiherpetic activities of chemically modified polysaccharides from *Polygonatum cyrtoneuma* Hua. *Carbohydr. Polym.* **2011**, *83*, 737–742.
22. He, Y.; Ye, M.; Jiang, L.; Du, Z.; Surahio, M.; Xu, H.; Li, J. Preparation characterization and bioactivities of derivatives of an exopolysaccharide from *Lachnum*. *Carbohydr. Polym.* **2015**, *117*, 788–796.
23. Wang, Y.; Chen, Y.; Du, H.; Yang, J.; Ming, K.; Song, M.; Liu, J. Comparison of the anti-duck hepatitis A virus activities of phosphorylated and sulfated *Astragalus* polysaccharides. *Exp. Biol. Med.* **2017**, *242*, 344–252.
24. Deng, C.; Fu, H.; Xu, J.; Shang, J.; Cheng, Y. Physicochemical and biological properties of phosphorylated polysaccharides from *Dictyophora indusiata*. *Int. J. Biol. Macromol.* **2015**, *72*, 894–899.
25. Rao, M.R.P.; Warriar, D.U.; Gaikwad, S.R.; Shevate, P.M. Phosphorylation of psyllium seed polysaccharide and its characterization. *Int. J. Biol. Macromol.* **2016**, *85*, 317–326.
26. Wang, X.; Zhang, Z.; Yao, Q.; Zhao, M.; Qi, H. Phosphorylation of low-molecular-weight polysaccharide from *Enteromorpha linza* with antioxidant activity. *Carbohydr. Polym.* **2013**, *96*, 371–375.
27. Song, Y.; Ni, Y.; Hu, X.; Li, Q. Effect of phosphorylation on antioxidant activities of pumpkin (*Cucurbita pepo*, Lady godiva) polysaccharide. *Int. J. Biol. Macromol.* **2015**, *81*, 41–48.
28. Xiong, X.; Huang, G.; Huang, H. The antioxidant activities of phosphorylated polysaccharide from native ginseng. *Int. J. Biol. Macromol.* **2019**, *126*, 842–845.
29. Dace, R.; McBride, E.; Brooks, K.; Gander, J.; Buszko, M.; Doctor, V.M. Comparison of the anticoagulant action of sulfated and phosphorylated polysaccharides. *Thromb. Res.* **1997**, *1*, 113–121.
30. Church, F.C.; Pratt, C.W.; Treavor, R.E.; Whina, H.C. Anticoagulant action of phosvitin and other phosphate containing polyanions is mediated by heparin cofactor II. *FEBS Lett.* **1988**, *237*, 26–30.
31. Lindahl, V.; Thunberg, L.; Backstrom, G.; Risenfeld, T.; Nordling, K.; Bjork, I. Extension and structural variability of the antithrombin-binding sequence in heparin. *J. Biol. Chem.* **1984**, *259*, 12368–12374.
32. Gu, Y.P.; Yang, X.M.; Luo, P.; Li, Y.Q.; Tao, Y.X.; Duan, Z.H.; Xiao, W.; Zhang, D.Y.; Liu, H.Z. Inhibition of acrolein-induced autophagy and apoptosis by a glycosaminoglycan from *Sepia esculenta* ink in mouse Leydig cells. *Carbohydr. Polym.* **2017**, *163*, 270–279.
33. Duan, Z.H.; Duan, W.W.; Li, F.P.; Li, Y.Q.; Luo, P.; Liu, H.Z. Effect of carboxymethylation on properties of fucoidan from *Laminaria japonica*: antioxidant activity and preservative effect on strawberry during cold storage. *Postharvest Biol. Tech.* **2019**, *151*, 127–133.
34. Zhang, M.; Su, N.; Huang, Q.; Zhang, Q.; Wang, Y.; Li, J.; Ye, M. Phosphorylation and antiaging activity of polysaccharide from *Trichosanthes* peel. *J. Food Drug Anal.* **2017**, *25*, 976–983.
35. Sun, X.; Pan, D.D.; Zeng, X.Q.; Cao, J.X. Phosphorylation modification of polysaccharides from *Enteromorpha*. *Food Sci.* **2011**, *24*, 73–77.

