

Promotion of Wound Healing and Prevention of Frostbite Injury in Rat Skin by Exopolysaccharide from the Arctic Marine Bacterium *Polaribacter* sp. SM1127

Mei-Ling Sun ¹, Fang Zhao¹, Xiu-Lan Chen ^{1,2}, Xi-Ying Zhang ^{1,2}, Yu-Zhong Zhang ^{1,2,3}, Xiao-Yan Song ¹, Cai-Yun Sun ¹ and Jie Yang ^{1,*}




Figure S1. ¹H NMR spectrum of SM1127 EPS on a Varian Inova-500 MHz Bruker NMR spectrometer at 70°C ($\delta_{\text{H}}=2.225$ ppm).




Figure S2. COSY spectrum of SM1127 EPS on a Varian Inova-500 MHz Bruker NMR spectrometer at 70°C ($\delta_c=30.89$ ppm).




Figure S3. TOCSY spectrum of SM1127 EPS on a Varian Inova-500 MHz Bruker NMR spectrometer at 70°C ($\delta_C=30.89$ ppm).




Figure S4. HSQC spectrum of SM1127 EPS on a Varian Inova-500 MHz Bruker NMR spectrometer at 70°C ($\delta_H=2.225$ ppm, $\delta_C=30.89$ ppm).