

Article

Oxygenated Theonellastrols: Interpretation of Unusual Chemical Behaviors Using Quantum Mechanical Calculations and Stereochemical Reassignment of 7 α -Hydroxytheonellasterol

A-Young Shin ^{1,2}, Hyi-Seung Lee ^{1,2}, Yeon-Ju Lee ^{1,2}, Jong Seok Lee ^{1,2}, Arang Son ³, Changhoon Choi ³ and Jihoon Lee ^{1,2,*}

¹ Korea Institute of Ocean Science & Technology (KIOST), Busan 49111, Korea; dkdud1624@kiost.ac.kr (A.-Y.S.); hslee@kiost.ac.kr (H.-S.L.); yjlee@kiost.ac.kr (Y.-J.L.); jslee@kiost.ac.kr (J.S.L.)

² Department of Marine Biotechnology, University of Science & Technology, Daejeon 34113, Korea

³ Department of Radiation Oncology, Samsung Medical Center, Seoul 06351, Korea; arang.son@sbsri.co.kr (A.S.); changhoon1.choi@samsung.com (C.C.)

* Correspondence: jihoonlee@kiost.ac.kr; Tel.: +82-51-664-3343

Received: 5 November 2020; Accepted: 26 November 2020; Published: 30 November 2020



Abstract: A total of eight new oxygenated 4-*exo*-methylene sterols, **1–8**, together with one artifact **9** and six known sterols **11–16**, were isolated from the marine sponge *Theonella swinhoei* collected from the Bohol province in Philippines. Structures of sterols **1–8** were determined from 1D and 2D NMR data. Among the sterols, 8 α -hydroxytheonellasterol (**4**) spontaneously underwent an allylic 1,3-hydroxyl shift to produce 15 α -hydroxytheonellasterol (**9**) as an artifact; this was rationalized by quantum mechanical calculations of the transition state. In addition, the 1,2-epoxy alcohol subunit of 8 α -hydroxy-14,15- β -epoxytheonellasterol (**5**) was assigned using the Gauge-Independent Atomic Orbital (GIAO) NMR chemical shift calculations and subsequent DP4+ analysis. Finally, comparison of the ¹³C chemical shifts of isolated 7 α -hydroxytheonellasterol (**6**) with the reported values revealed significant discrepancies at C-6, C-7, C-8, and C-14, leading to reassignment of the C-7 stereochemistry in the known structure.

Keywords: *Theonella swinhoei*; marine natural product; oxygenated theonellasterol; GIAO NMR chemical shift calculation; anti-inflammatory activity

1. Introduction

Marine invertebrates have a broad array of pharmacologically and structurally attractive natural products, some of which exhibit extraordinary potencies and selectivities against human diseases, thus rendering them potential drug candidates [1]. The marine sponge *Theonella swinhoei* has been one of the most diverse sources of natural products. Since the isolation of theonellasterol A from *T. swinhoei* in 1981 [2], various classes of molecules, including polyketide [3–12], peptides [13–18], and sterols [2,19–22], have been identified and evaluated for their biological properties. The characteristic 4-*exo*-methylene-sterols represent the largest family of secondary metabolites isolated from the genus *Theonella* sponges. To date, at least 37 sterols, including swinhoeisterols [21,22], swinhosterols [19,23], conicasterols [2,24–27], and theonellasterols [2,25,26,28,29], have been identified through extensive research on this marine invertebrate. Recently, these marine sterols and their synthetic derivatives have been investigated as potent farnesoid-x-receptor (FXR) antagonists to protect the liver from injuries caused due to bile acid overload [25,30].

Generally, structures of the natural sterols can be analyzed from the key NMR (HMBC, NOESY) correlations arising from the methyl groups at C-18 and C-19. However, precise analyses utilizing conventional NMR techniques are problematic in some cases because of the complex overlapping of non-functionalized sp^3 methylene peaks in the 1H NMR spectrum and the absence of 1H signals from oxygenated tertiary carbon atoms. Current advances in the prediction of NMR shielding constants employing quantum mechanical calculations have provided alternative tools to clarify the ambiguities in the course of structure determination [31,32]. For instance, the structure of conicasterol F, bearing a tetra-substituted epoxide at C-8 and C-14, was deduced from GIAO calculations of ^{13}C NMR chemical shifts and DFT-calculated ROE-distances [26].

As a part of our ongoing research to isolate bioactive and structurally interesting natural products, we investigated the metabolites of *T. swinhoei*, collected from the Bohol province in Philippines and identified eight novel theonellasterol analogs 1–8 (Figure 1), one artifact 9, and six known sterols 11–16 (Figure S2, Supporting Information). Herein, we report the structural assignments highlighted with DFT calculations to provide a rationale for the unusual chemical behaviors of oxygenated 4-*exo*-methylene sterols. The structure of 8 α -hydroxy-14,15- β -epoxytheonellasterol (5) was deduced using GIAO chemical shift calculations. In addition, the structure of 7 α -hydroxytheonellasterol (6), determined by Faulkner and Qureshi in 2000 [28], was reevaluated due to significant discrepancies between the reported ^{13}C NMR chemical shifts and the spectroscopic data obtained in this study.




Figure 1. Structures of 1–8.

2. Results and Discussion

A total of eight new oxygenated theonellasterols 1–8 were obtained from the hexane extract of *T. Swinhoei*. Theonellasterol-5,8-oxide (1) was isolated as a colorless oil. Its molecular formula was determined to be $C_{30}H_{48}O_3$ by HRESIMS (m/z $[M + Na]^+$ 479.3485, calcd 479.3501), indicating seven degrees of unsaturation. Inspection of the 1H and ^{13}C NMR data, including correlations from the HSQC spectrum, provided sufficient information to propose a theonellasterol-type skeleton: two quaternary sp^2 carbon atoms (δ_C 152.6, 150.6), an sp^2 methylene (δ_C 111.2, δ_H 5.13, 5.59), an oxymethine (δ_C 69.8, δ_H 4.64), two singlet methyl (δ_C 20.1/ δ_H 0.68, δ_C 19.0/ δ_H 0.92), three doublet methyl (δ_C 20.0/ δ_H 0.90, δ_C 19.6/ δ_H 0.88, δ_C 19.6/ δ_H 0.98), and one triplet methyl (δ_C 12.9, δ_H 0.93) groups. Additionally, an sp^2 methine (δ_C 116.8, δ_H 5.80) and two oxygenated tertiary carbons (δ_C 91.9, 86.8) were detected as characteristics of compound 1, suggesting identification of a new analog. While the

exocyclic 4,4-di-substituted $\Delta^{4,30}$ -olefin and endocyclic tetra-substituted olefin at $\Delta^{8,14}$ or $\Delta^{8,9}$ were known to be structural features of theonellasterols, the sp^2 methine in **1** indicated the presence of an endocyclic tri-substituted olefin that may be generated by an isomerization or rearrangement of the tetra-substituted olefin. Moreover, the additional oxygenated tertiary carbons and a higher degree of hydrogen deficiency (DBE = 7) compared to theonellasterol A (DBE = 6) indicated the existence of an oxygenated theonellasterol framework bearing an additional ring [33,34].

HMBC correlations from CH_3 -19 (δ_H 0.68)/ H_2 -30 (δ_H 5.59, 5.13)/ H_2 -6 (δ_H 1.77, 1.33) to δ_C 91.9 and from H -9 (δ_H 1.59)/ H_2 -11 (δ_H 1.33, 2H) to δ_C 86.8 corresponded to the oxygenated tertiary carbon at C-5 and C-8, respectively. The carbon-carbon connectivity for the B-ring was determined by HMBC correlations from H_2 -7 (δ_H 1.86, 1.43) to C-9 (δ_C 58.3)/C-6 (δ_C 30.6), from H_2 -6 (δ_H 1.77, 1.33) to C-4 (δ_C 150.6)/C-5 (δ_C 91.9)/C-10 (δ_C 44.2), and from H -9 to C-1 (δ_C 37.8)/C-7 (δ_C 31.5)/C-10 (δ_C 44.2). The sp^2 methine (δ_C 116.8, δ_H 5.80) was located at C-15 to form the $\Delta^{14,15}$ -olefin, as evident from the 1H - 1H COSY cross peak for H -15- H_2 -16- H -17 and the HMBC correlations from CH_3 -18 (δ_H 0.92)/ H_2 -12 (δ_H 1.93, 1.23)/ H_2 -16 (δ_H 2.36, 2.01) to C-14 (δ_C 152.6). Since the carbon framework of **1** turned out to be identical to those of theonellasterols, a covalent bond between the two oxygen atoms at C-5 and C-8 was speculated to fulfill the hydrogen deficiency. The relative configuration of the [2,2,2]-bicyclic B-ring, including those of the consecutive stereocenters at C-13, C-17, and C-20, was determined by NOESY cross peaks between CH_3 -19 and H -6 β (δ_H 1.77)/ H -7 β (δ_H 1.86) and between CH_3 -18 and H -7 β / H -20 (δ_H 1.59) (Figure 2). The absolute configuration of **1**, including the configuration of C-24, was deduced to be (3*S*,5*S*,8*R*,9*R*,10*R*,13*R*,17*R*,20*R*,24*S*), considering its biosynthetic correlation with other theonellasterol analogs. Assignment of the 24*S* configuration using the δ_{H-26} - δ_{H-27} values is discussed later in this article.




Figure 2. Optimized structures of **1** (a) and **2** (b) at the mPW1PW91/6-31G* level of theory and key NOESY correlations (arrows).

8 β -Hydroxytheonellasterol (**2**) was obtained as a colorless oil. Its molecular formula was determined to be $C_{30}H_{50}O_2$ by HRFABMS (m/z [$M - H_2O + H$] $^+$ 425.3780, calcd 425.3783), indicating six degrees of unsaturation. The 1H and ^{13}C NMR data (Tables S1 and S2, Supporting Information) suggested that compounds **1** and **2** had most of the characteristics in common; however, only one oxygenated tertiary carbon (δ_C 83.5) was detected. The oxygenated tertiary carbon exhibited correlations with H -7 (δ_H 2.67, 1.20)/ H -15 (δ_H 5.52) in the HMBC spectrum to be assigned at C-8. The endocyclic olefin was positioned at $\Delta^{14,15}$, based on the interpretation of HMBC correlations from CH_3 -18 (δ_H 1.11) to C-14 (δ_C 151.2) and from H -15 (δ_H 5.52) to C-8/C-13 (δ_C 47.7)/C-14, as well as a spin system for H -15- H -16- H -17 in the 1H - 1H COSY spectrum. Careful inspection of the 1H NMR spectrum and the NOESY spectrum revealed that the axial H -11 β (δ_H 1.82) and H -6 β (δ_H 1.89) were more deshielded

than equatorial H-11 α (δ_{H} 1.30) and H-6 α (δ_{H} 1.44) (Figure 2). Additionally, axially oriented CH₃-18 and CH₃-19 (δ_{H} 0.97) were shifted downfield compared to those in the reported theonellasterols [25]. These features accounted for the β -orientation of the hydroxyl group at C-8, initiating additional 1,3-diaxial interactions [35,36].

15 β -Hydroxytheonellasterol (**3**) was isolated as a colorless oil. Its molecular formula was determined to be C₃₀H₅₀O₂ by HRESIMS (m/z [M – H₂O + H]⁺ 425.3789, calcd 425.3783). The 1D and 2D NMR spectra of **3** were almost identical to those of theonellasterol A. However, an additional oxymethine (δ_{C} 70.3, δ_{H} 4.64) was detected, of which the location was assigned as C-15 by HMBC correlations from δ_{H} 4.64 to C-13 (δ_{C} 43.8)/C-17 (δ_{C} 54.2) and from H₂-16 (δ_{H} 1.95, 1.60) to δ_{C} 70.3. The β -orientation of the hydroxyl group at C-15 was established from the NOESY correlation between H-15 and H-17 (δ_{H} 1.60).

8 α -Hydroxytheonellasterol (**4**) was obtained as a chemically labile compound, and reliable HRMS data could not be obtained. However, the 1D and 2D NMR spectra of **4** suggested the presence of an oxygenated theonellasterol scaffold bearing an oxygenated tertiary carbon (δ_{C} 86.6) and an sp² methine (δ_{C} 123.2, δ_{H} 5.61). HMBC correlations from H₂-6 (δ_{H} 1.54)/H-9 (δ_{H} 2.27) to the carbon at δ_{C} 86.6 and from the proton at δ_{H} 5.61 to C-13 (δ_{C} 48.0)/C-16 (δ_{C} 35.2)/C-17 (δ_{C} 60.0) revealed that the oxygenated tertiary carbon and the sp² methine group were located to C-8 and C-15, respectively. This further suggested that **4** was the C-8 epimer of 8 β -hydroxytheonellasterol (**2**). For this case, the extra 1,3-diaxial interaction, which was observed in the case of **2**, was undetectable from H₂-6 (δ_{H} 1.67, 1.54), H₂-11 (δ_{H} 1.49, 2H), CH₃-18 (δ_{H} 0.83), and CH₃-19 (δ_{H} 0.68), supporting the α -orientation of OH-8. Instead, the ¹H chemical shift of H-5 (δ_{H} 3.03) was further downfield compared to those of the reported sterols [25], and H-7 β (δ_{H} 1.76) exhibited NOESY correlations with Me-18 and Me-19. This phenomenon could be rationalized using quantum mechanical calculations. Geometry optimization of **4** at the mPW1PW91/6-31G* level of theory revealed a boat conformation for the B-ring to initiate 1,4-flagpole interactions between H-5 and OH-8 α ($d = 2.18$ Å), resulting in the downfield shift of H-5 (Figure 3). Additionally, atomic distances from H-7 to H-18 and H-19 were measured as 2.07 Å and 3.91 Å, respectively, which are close enough to exhibit NOESY correlations.




Figure 3. The optimized structure of **4** at the mPW1PW91/6-31G* level of theory and the calculated atomic distance between H-5 and OH-8.

Due to its labile nature, sterol **4** was entirely decomposed into a complex mixture of unidentifiable compounds within several days. However, extended storage in benzene afforded an artifact as a single compound (Figure 4). Its molecular formula was determined to be C₃₀H₅₀O₂ by HRFABMS (m/z [M – H₂O + H]⁺ 425.3781, calcd 425.3783). The 1D and 2D NMR data were almost identical to those of compound **3**, except for the deshielded oxymethine (δ_{C} 84.4, δ_{H} 4.97). The oxymethine exhibited a ¹H-¹H COSY cross peak with H₂-16 (δ_{H} 1.49) and HMBC correlations with C-13 (δ_{C} 43.5)/C-14 (δ_{C} 141.1)/C-17 (δ_{C} 54.1) to be assigned at C-15. Based on the stereochemistry of **3**, the artifact was determined to be 15 α -hydroxytheonellasterol (**9**), which was produced through an allylic 1,3-hydroxyl migration of **4**.




Figure 4. (a) The spontaneous transformation of 8 α -hydroxytheonellasterol (**4**) to an artifact, 15 α -hydroxytheonellasterol (**9**). (b) Time dependent ^1H NMR spectrum in C_6D_6 indicating the transformation.

Calculation of transition state energy using the Linear Synchronous Transit (LST) method revealed that the energy barrier for the transformation of **4** to **9** was only 0.6 kcal/mol, which can explain the instability of **4** (Figure 5). The structure of the transition state (TS) was turned out to be almost identical to that of **9**. In the transition state, the B-ring was flipped to a chair-like conformation, bringing O-8 and sp^2 C-15 (1.43 Å) in close proximity. The atomic distance between C-8 and O-8 in TS was measured to be 3.09 Å, suggesting that the C-8–O-8 bond was actually broken before the TS to form a new C–O bond at C-15. In addition, the C-8–C-14 bond length was estimated to be 1.34 Å, indicating olefin migration from $\Delta^{14,15}$ to $\Delta^{8,14}$ (Figure S4, Supporting Information). Although **9** was slightly more stable than **4** at rt ($\Delta G^\circ = -0.3$ kcal/mol), the low activation energy and the formation of a more rigid tetra-substituted olefin perhaps shifted the chemical equilibrium toward **9**.

8 α -Hydroxy-14,15- β -epoxy-theonellasterol (**5**) was isolated as an amorphous powder. Its molecular formula was determined to be $\text{C}_{30}\text{H}_{50}\text{O}_3$ (m/z [$\text{M} + \text{Na}$] $^+$ 481.3659, calcd 481.3658) by HRESIMS, indicating six degrees of unsaturation. The 1D and 2D NMR data of **5** revealed an oxygenated theonellasterol framework bearing an additional oxymethine (δ_{C} 58.8, δ_{H} 3.33) and two oxygenated tertiary carbons (δ_{C} 75.9, 71.3). In this case, only one olefin (δ_{C} 154.2, 103.3), corresponding to 4-*exo*-methylene, was found, indicating the presence of an additional ring to satisfy the unsaturation index. HMBC correlations from H_2 -7 (δ_{H} 1.82, 1.60) to C-8 (δ_{C} 71.3), from H_2 -16 (δ_{H} 1.96, 0.93) to C-14 (δ_{C} 75.9)/C-15 (δ_{C} 58.8), and from Me-18 (δ_{H} 0.71) to C-14 (δ_{C} 75.9) indicated the presence of a 1,2-epoxy alcohol moiety within the C-8–C-14–C-15. The upfield shift of C-15 oxymethine (δ_{C} 58.8) and the unusual HMBC correlations from the hydroxyl peak at δ_{H} 3.47 to C-7 (δ_{C} 28.3)/C-8 (δ_{C} 71.3)/C-14 (δ_{C} 75.9) suggested the presence of an 8-hydroxy-14,15-epoxide subunit. Additionally, β -orientation of the epoxide was assigned by the NOESY cross peak between H-17 (δ_{H} 1.41) and H-15 (Figure S3, Supporting Information), and the downfield shift of H-5 (δ_{H} 3.35) was perhaps attributable to an α -orientation of the C-8 hydroxyl group, similar to that of 8 α -hydroxytheonellasterol (**4**).




Figure 5. Free energy diagram for 1,3-hydroxyl migration in 8 α -hydroxy-theonellasterol (**4**) to generate 15 α -hydroxytheonellasterol (**9**). Geometry optimizations of compounds **4**, **9**, and the transition state (TS) were performed at the mPW1PW91/6-31G* level of theory.

Due to the limited spectroscopic data for the 1,2-epoxy alcohol subunit of **5**, GIAO NMR chemical shift calculations were employed to support the assignments (Table 1). Although our observations suggested the maximum possibility of the 14,15-epoxide isomers **5-I** and **5-II**, the formation of 8,14-epoxide isomers **5-III** and **5-IV** could not be ruled out. The ^{13}C NMR chemical shift calculations of the four sets of 8,14,15-isomers using the mPW1PW91/6-31G** level of theory and subsequent DP4+ analysis indicated 100% probability of 8 α -hydroxy-14,15- β -epoxy-isomer **5-I** [26]. The correlation coefficient (R^2) in the regression analysis of the experimental versus calculated ^{13}C chemical shifts of **5-I** was calculated to be 0.9908, indicating that the structure assignment was highly reliable. As anticipated from the downfield shift of H-5, the B-ring in the optimized structure adopted a boat conformation to rationalize the downfield shift of H-5 by 1,4-flagpole interactions (Table S4, Supporting Information).

Compound **6** was isolated as a colorless needle-shaped solid. The molecular formula was determined to be $\text{C}_{30}\text{H}_{50}\text{O}_2$ by HRESIMS (m/z [$\text{M} + \text{Na}$] $^+$ 465.3705, calcd 465.3709), indicating six degrees of unsaturation. Comparison of the NMR spectra of **6** with the previously reported data revealed an oxygenated theonellasterol-type framework bearing an additional oxymethine (δ_{C} 66.7, δ_{H} 4.64). The HMBC correlation from H₂-6 (δ_{C} 1.77, 1.59) to δ_{C} 66.7, as well as the ^1H - ^1H COSY cross peak between the protons at δ_{H} 4.64 and H₂-6, suggested that the oxymethine was positioned at C-7.

Because the isolation of 7 α -hydroxytheonellasterol was reported in 2000 by Faulkner and Qureshi [28], compound **6** was initially considered to be 7 β -hydroxytheonellasterol, as deduced from the comparison of ^1H and ^{13}C chemical shifts (Figure 6a). However, a lack of NOESY signals corresponding to H-7 led us to synthesize 3,7-dimethyl ether **10** from **6**. Surprisingly, the NOESY data of **10** indicated a correlation between OMe-7 (δ_{H} 3.14) and H-9 (δ_{H} 2.29), supporting the α -orientation of the C-7 hydroxyl group. In addition, methylation of swinhosterol C (**11**), known as 7 α -OMe-theonellasterol, afforded a compound that was spectroscopically identical to **10** (Figure 6b,c). Single crystal X-ray diffraction of **6** further confirmed a 7 α -hydroxytheonellasterol structure (Figure 7). Considering the large differences in the ^{13}C chemical shifts at C-6, C-7, C-8, and C-14, we speculate that the previously reported compound is the 7 β -epimer of **6**.

Table 1. Comparison of experimental ^{13}C chemical shifts in C_6D_6 with calculated ^{13}C shifts for isomers I–IV.


5-I
 (8- α -hydroxy-14,15- β -epoxide)

5-II
 (8- β -hydroxy-14,15- β -epoxide)

5-III
 (8,14- α -epoxy-15- β -alcohol)

5-IV
 (8,14- β -epoxy-15- β -alcohol)

Position	δ_{exp} (ppm)	δ_{calcd} (ppm)			
		5-I	5-II	5-III	5-IV
C-7	28.3	31.7	35.5	31.8	30.9
C-8	71.3	72.4	74.2	63.4	65.0
C-9	55.3	51.6	55.1	47.2	50.0
C-13	42.5	45.7	47.3	43.3	42.5
C-14	75.9	76.4	75.1	80.0	72.0
C-15	58.8	60.3	62.5	73.9	65.8
C-16	32.3	25.7	27.6	38.7	38.8
DP4+		100%	0%	0%	0%
R^2		0.9908	0.9828	0.9772	0.9845
MAD ^a		1.38	1.91	2.30	1.64

^a Mean absolute deviation.**Figure 6.** Cont.




Figure 6. Structure confirmation of 7 α -hydroxytheonellasterol (**6**): (a) Difference between the ^{13}C chemical shifts of **6** and the reported values in CDCl_3 . (b) Methylation of **9** and swinhosterol C (**11**). (c) ^1H NMR spectrum of **10** synthesized from **6** (red) and ^1H NMR spectrum of crude **10** synthesized from swinhosterol C (**11**) in C_6D_6 .




Figure 7. ORTEP drawing of compound **6** based on X-ray data.


8 β -Hydroxy-7 α -formyl-B-northeonellasterol (**7**) was isolated as a colorless oil. Its molecular formula was determined to be $\text{C}_{30}\text{H}_{50}\text{O}_3$ by HRESIMS (m/z $[\text{M} + \text{Na}]^+$ 481.3660, calcd 481.3658). Analysis of the 1D and 2D NMR data of **7** revealed an oxygenated theonellasterol-like skeleton bearing an aldehyde (δ_{C} 204.2, δ_{H} 9.75) and an additional oxygenated tertiary carbon (δ_{C} 87.2). Since the aldehyde moiety is known as a unique feature of 8 β -hydroxy-B-norconicasta-6 α -aldehyde among the sterols isolated from *T. swinhoei* [37], the 6/5/6/5-fused cyclic backbone of **7** was assigned by comparing the NMR data. Compound **7** could be differentiated from 8 β -hydroxy-B-norconicasta-6 α -aldehyde in the ethyl substituent at C-24, which was assigned based on the HMBC correlations from a triplet methyl group (δ_{H} 0.95) to C-24 (δ_{C} 46.9)/C-28 (δ_{C} 23.8).

28-Homoswinhoeisterol (**8**) was isolated as a yellow oil. The molecular formula was determined to be $\text{C}_{30}\text{H}_{48}\text{O}_2$ by HRESIMS (m/z $[\text{M} + \text{Na}]^+$ 465.3549, calcd 463.3552). The IR spectrum of **8** clearly indicated the presence of a hydroxyl group (3343 cm^{-1}) and a ketone group (1593 cm^{-1}). The features of the IR and NMR spectra of this compound were almost identical to those of swinhoeisterol A. The only difference was found in the triplet methyl group (δ_{H} 0.84), which was involved in a spin system for H-24–H₂-28–CH₃-29, as evident from the ^1H - ^1H COSY spectrum. This suggested that


the ethyl group was located at C-24. In addition, the plausible biogenetic pathway reported by Zhang et al. [21] indicated that compound **8** could be originated from swinhosterol A through an intramolecular aldol-type reaction, which strongly suggested that the absolute configuration of **8** is (3*S*,5*R*,7*R*,10*S*,13*R*,17*R*,20*R*,24*S*). To date, 28-homoswinhoiesterol (**8**) is the only 6/6/5/7-fused cyclic sterol derived from C₃₀ sterols such as theonellasterol A and swinhosterol A.

Generally, the 24*S* configurations of 24-ethyl-sterol analogs are deduced from the ¹³C chemical shift differences between CH₃-26 and CH₃-27 [38]. However, the differences for compounds **2** (0.6 ppm), **5** (0.7 ppm), and **7** (0.8 ppm) were not significant enough for determining the configuration of C-24, and hence, a complementary method was required for the assignment. To establish a universal database using ¹H NMR data, the absolute values of $\delta_{\text{H-26}} - \delta_{\text{H-27}}$ were obtained from the sets of sterol-type compounds bearing a (21*R*,24*R*) or (21*R*,24*S*)-21,26-dimethyl-24-ethylhexane side chain (Table S11, Supporting Information). Values calculated for the (21*R*,24*R*)-set were higher than 0.04 ppm, whereas those for the (21*R*,24*S*)-set were smaller than 0.04 ppm. The validity of our database was evaluated using six known 24*S*-ethyl-sterols isolated in this study: theonellasterol A (**12**), E (**13**), G (**14**), and K (**15**); swinhosterol A (**16**) and C (**11**). In all the cases, the differences were smaller than 0.03 ppm to prove the reliability of the database. Further, this method was extended to the new compounds **1–8**. The differences for all of them were in the desirable range (<0.03 ppm), confirming their 24*S* configuration (Figure S8, Supporting Information).

With the theonellasterol analogs in hand, their anti-inflammatory activities were investigated using the murine macrophage RAW264.7 cells. Treatment with lipopolysaccharide (LPS) in RAW264.7 macrophages stimulates secretion of pro-inflammatory cytokines, including interleukin-6 (IL-6). Levels of IL-6 secreted by the cells were quantified by the enzyme-linked immunosorbent assay (ELISA) method. Unfortunately, useful levels of biological properties for the new oxygenated theonellasterols (**1–9**) were unidentified. However, theonellasterol G (**14**) showed a moderate anti-inflammatory activity with an IC₅₀ of 4.4 $\mu\text{g/mL}$ (9.2 μM), and theonellasterol K (**15**) exhibited a weak anti-inflammatory activity with an IC₅₀ of 16.7 $\mu\text{g/mL}$ (35.2 μM) (Figure 8).



(a)



(b)

Figure 8. Anti-inflammatory effect of (a) theonellasterol G (**14**) and (b) theonellasterol K (**15**). RAW264.7 cells were pre-treated with indicated concentrations of theonellasterols for 3 h, followed by treatment with 5 ng/mL of lipopolysaccharide (LPS). After 24 h of incubation, secreted interleukin-6 (IL-6) levels were determined as described in Experimental section. Data are presented as mean \pm standard deviation ($n = 4$). Statistical analysis was performed using one-way analysis of variance with Dunnett's post-hoc analysis. * $p < 0.05$; *** $p < 0.001$ vs. LPS-stimulated group.

3. Materials and Methods

3.1. General Experimental Procedures

Specific optical rotations were obtained on a Rudolph Research Analytical (Autopol III) polarimeter (Rudolph Research Analytical, Hackettstown, NJ, USA). IR spectra were recorded on a JASCO FT/IR-4100 spectrophotometer (JASCO Corporation, Tokyo, Japan). The 1D (^1H and ^{13}C) and 2D (COSY, HSQC, HMBC, and NOESY) NMR spectra were taken in C_6D_6 or CDCl_3 using Bruker 600 MHz spectrometer (Bruker BioSpin GmbH, Rheinstetten, Germany) at 297.1 K. ^1H NMR spectra were collected after 64 scans, and ^{13}C NMR spectra were collected at a range of 10,000–15,000 scans depending on the sample concentrations. The mixing time for NOESY experiments was set as 0.3 s. Chemical shifts are reported in parts per million relative to C_6D_6 (δ_{H} 7.16, δ_{C} 128.4) and CDCl_3 (δ_{H} 7.26, δ_{C} 77.1). High resolution mass-spectra were obtained on a Waters Q-TOF spectrometer (Waters Corporation, Milford, MA, USA) equipped with an ESI source and a JEOL JMS-700 spectrometer (JEOL Ltd., Tokyo, Japan) with an FAB Source at Korea Basic Science Institute (KBSI) (Seoul, Republic of Korea). MPLC was performed using the TELEDYNE ISCO CombiFlash Companion with the TELEDYNE ISCO RediSep Normal-phase Silica Flash Column (Teledyne ISCO, Lincoln, NE, USA). HPLC was performed on a PrimeLine Binary pump (Analytical Scientific Instruments, Inc., El Sobrante, CA, USA) utilizing Silica columns (YMC-Pack Silica, 250×10 mm I.D. or 250×4.6 mm I.D., $5 \mu\text{m}$; YMC Co. Ltd., Kyoto, Japan), the Shodex RI-101 (Shoko Scientific Co. Ltd., Yokohama, Japan), or the UV-M201.

3.2. Biological Material

The biological material was collected in March 2016 from the Bohol province in Philippines ($9^\circ 43' 31.86''$ N, $124^\circ 32' 35.57''$ E) at a depth of 15 m using scuba diving. The sponge was kept frozen at -20°C until identified as *Theonella swinhoei* and chemically analyzed. A voucher sample (163PIL-102) has been stored at the marine biotechnology center, Korea Institute of Ocean Science & Technology (KIOST).

3.3. Extraction and Isolation

The specimen (wet wt. 1.8 kg) was lyophilized and extracted with MeOH ($2.5 \text{ L} \times 3$) and CH_2Cl_2 ($2.5 \text{ L} \times 3$) repeatedly at room temperature. The extracts were combined and then concentrated under reduced pressure. The residue was partitioned with *n*-butanol (7.0 L) and water (5.0 L) to yield 55.43 g of organic soluble material. The *n*-butanol layer was further partitioned between *n*-hexane (2.0 L) and 15% aqueous methanol (2.0 L). The hexane fraction was concentrated and subjected to flash column chromatography over SiO_2 (0.040–0.063 mm, 230–400 mesh) with a stepwise gradient solvent system (100%, 93.7%, 90%, 83%, 80%, 75%, 50% hexane/EtOAc, 100% EtOAc).

The 90% and 83% hexane fractions were combined (3.22 g) and separated using MPLC on SiO_2 with a gradient solvent system from 100% hexane to 100% EtOAc over 40 min to afford seven subfractions (based on TLC analysis). The third subfraction gave theonellasterol A (**12**) (1.00 g) as a pure compound without further purification. The sixth subfraction was separated using HPLC (hexane/EtOAc = 5/1) to yield compounds **2** (8.9 mg, $t_{\text{R}} = 26$ min) and **1** (3.5 mg, $t_{\text{R}} = 28$ min).

The 80% hexane fraction (273.4 mg) was directly separated using HPLC (hexane/EtOAc = 4/1) to yield compound **8** (31.4 mg, $t_{\text{R}} = 21$ min) and compound **5** (1.2 mg, $t_{\text{R}} = 26$ min). The 75% and 50% hexane fractions were combined (786.0 mg) and separated using MPLC on SiO_2 with a gradient solvent system from 100% hexane to 100% EtOAc over 40 min to afford six subfractions (based on TLC analysis). The third subfraction (200.0 mg) was separated using HPLC (hexane/acetone = 6/1) to yield swinhosterol C (**11**) (1.7 mg, $t_{\text{R}} = 17$ min), **4** (1.2 mg, $t_{\text{R}} = 20$ min), and theonellasterol K (**15**) (30.3 mg, $t_{\text{R}} = 24$ min). The fourth subfraction (57.2 mg) was separated using HPLC ($\text{CH}_2\text{Cl}_2/\text{MeOH} = 100/1$) to yield swinhosterol A (**16**) (12.0 mg, $t_{\text{R}} = 26$ min), **7** (1.6 mg, $t_{\text{R}} = 28$ min), and **3** (3.5 mg, $t_{\text{R}} = 52$ min). The fifth subfraction (72.9 mg) was separated using HPLC (hexane/acetone = 6/1) to yield **6** (3.1 mg, $t_{\text{R}} = 30$ min), and theonellasterol G (**14**) (5.9 mg, $t_{\text{R}} = 52$ min).

The 100% EtOAc fraction (372.3 mg) was separated using MPLC with a gradient solvent system from 70% hexane to 100% EtOAc to afford four subfractions (based on TLC analysis). The fourth subfraction (78.1 mg) was separated using HPLC (hexane/acetone = 4/1) to yield theonellasterol E (**13**) (5.0 mg, $t_R = 40$ min).

Theonellasterol-5,8-oxide (**1**): colorless oil; $[\alpha]_D^{25} + 20.0$ (c 0.1, MeOH); IR (ATR) ν_{\max} 3417, 2957, 2851, 1738, 1455, 1027 cm^{-1} ; ^1H NMR and ^{13}C NMR, see Tables S1 and S2, Supporting Information; HRESIMS m/z 479.3485 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{30}\text{H}_{48}\text{O}_3\text{Na}$, 479.3501).

8β -Hydroxytheonellasterol (**2**): colorless oil; $[\alpha]_D^{25} + 20.0$ (c 0.1, MeOH); IR (ATR) ν_{\max} 3353, 2929, 1751, 1489, 1410, 1101 cm^{-1} ; ^1H NMR and ^{13}C NMR, see Tables S1 and S2, Supporting Information; HRFABMS m/z 425.3780 $[\text{M} - \text{H}_2\text{O} + \text{H}]^+$ (calcd for $\text{C}_{30}\text{H}_{49}\text{O}$, 425.3783).

15β -Hydroxytheonellasterol (**3**): colorless oil; $[\alpha]_D^{25} + 40.0$ (c 0.1, MeOH); IR (ATR) ν_{\max} 3345, 2963, 2938, 2871, 1711, 1379, 1039 cm^{-1} ; ^1H NMR and ^{13}C NMR, see Table S1 and S2, Supporting Information; HRESIMS m/z 425.3789 $[\text{M} - \text{H}_2\text{O} + \text{H}]^+$ (calcd for $\text{C}_{30}\text{H}_{49}\text{O}$, 425.3783).

8α -Hydroxytheonellasterol (**4**): colorless oil; ^1H NMR and ^{13}C NMR, see Tables S1 and S2, Supporting Information.

8α -Hydroxy-14,15- β -epoxy-theonellasterol (**5**): amorphous powder; $[\alpha]_D^{25} + 30.0$ (c 0.1, MeOH); IR (ATR) ν_{\max} 3567, 2954, 2929, 2861, 1727, 1377, 1254, 1035 cm^{-1} ; ^1H NMR and ^{13}C NMR, see Table S1 and S2, Supporting Information; HRESIMS m/z 481.3659 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{30}\text{H}_{50}\text{O}_3\text{Na}$, 481.3658).

7α -Hydroxytheonellasterol (**6**): colorless needle-shaped solid; $[\alpha]_D^{25} + 6.67$ (c 0.1, MeOH); IR (ATR) ν_{\max} 3359, 2929, 1416, 1333, 1100 cm^{-1} ; ^1H NMR and ^{13}C NMR, see Tables S1 and S2, Supporting Information; HRESIMS m/z 465.3705 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{30}\text{H}_{50}\text{O}_2\text{Na}$, 465.3709).

8β -Hydroxy- 7α -formyl-B-northeonellasterol (**7**): colorless oil; $[\alpha]_D^{25} + 40.0$ (c 0.1, MeOH); IR (ATR) ν_{\max} 3434, 2933, 2865, 1710, 1458, 1374, 1031 cm^{-1} ; ^1H NMR and ^{13}C NMR, see Tables S1 and S2, Supporting Information; HRESIMS m/z 481.3660 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{30}\text{H}_{50}\text{O}_3\text{Na}$, 481.3658).

28-Homoswinhoeisterol (**8**): yellow oil; $[\alpha]_D^{25} + 100.0$ (c 0.58, CHCl_3); IR (ATR) ν_{\max} 3343, 2936, 1593, 1458, 1413, 1120, 1042 cm^{-1} ; ^1H NMR and ^{13}C NMR, see Table S1 and S2, Supporting Information; HRESIMS m/z 465.3549 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{30}\text{H}_{48}\text{O}_2\text{Na}$, 463.3552).

15α -Hydroxytheonellasterol (**9**): colorless oil; $[\alpha]_D^{25} + 76.6$ (c 0.1, MeOH); IR (ATR) ν_{\max} 3367, 2961, 2925, 2872, 1702, 1458, 1381 cm^{-1} ; ^1H NMR and ^{13}C NMR, see Tables S1 and S2, Supporting Information; HRFABMS m/z 425.3781 $[\text{M} - \text{H}_2\text{O} + \text{H}]^+$ (calcd for $\text{C}_{30}\text{H}_{49}\text{O}$, 425.3783).

Theonellastrol-3,7-dimethyl ether (**10**): white powder; $[\alpha]_D^{25} + 23.33$ (c, 0.1, MeOH); IR (ATR) ν_{\max} 2957, 2953, 2872, 2353, 1593, 1102 cm^{-1} ; ^1H NMR and ^{13}C NMR, see Supporting Information; HRESIMS m/z 493.4009 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{32}\text{H}_{54}\text{O}_2\text{Na}$, 493.4016).

3.4. ^{13}C Chemical Shift Calculations

The conformational searches were performed using the MacroModel software (Maestro Materials Science 3.7.013 based on Maestro Core 12.3.013, MMshare Version 4.9.013, Release 2020-1, Platform Windows-x64; New York, NY, USA). The conformers within an energy threshold of 5 kJ/mol were optimized employing DFT calculations at the mPW91PW1/6-31G* level of theory to estimate gas phase energies and Gibbs free energies. All of the optimizations were performed at “fine” grid density and “ultrafine” accuracy level. The structure that has the lowest gas phase energy was selected, and NMR shielding constants were calculated with the mPW91PW1/6-31G**/CPCM benzene basis set. The calculated ^{13}C chemical shifts of compounds **5–I–IV** were referenced to the ^{13}C chemical shift of tetramethylsilane (TMS), computed with the same level of theory (for the details, see Supporting Information).

3.5. IL-6 Assay

The murine macrophage RAW264.7 cells were obtained from Dong Hyun Sohn, Pusan National University, Yangsan, South Korea. The RAW264.7 cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% fetal bovine serum (FBS) and antibiotics. For determining interleukin-6 (IL-6) production, RAW264.7 cells were pre-treated with eight new sterols (1–3, 5–9) and six known sterols (11–16) at various concentrations for 3 h, and then were incubated with or without 5 ng/mL of lipopolysaccharide (LPS; eBioscience, San Diego, CA, USA). After 24 h of incubation, the supernatant was collected and subjected to enzyme-linked immunosorbent assay (ELISA). The levels of IL-6 were measured by using mouse IL-6 Quantikine ELISA kit (R&D systems, Minneapolis, MN, USA) according to the manufacturer's instructions.

4. Conclusions

A total of eight new oxygenated 4-*exo*-methylene sterols 1–8 and six known sterols (11–16) were isolated from *T. swinhoei*. The C-7 stereochemistry of the reported 7 α -hydroxytheonellasterol has been revised based on the outcome of a series of chemical modifications and the X-ray crystallography data of 6. The stereo and regiochemistry of the 1,2-epoxyalcohol moiety in 8 α -hydroxy-14,15- β -epoxy-theonellasterol (5) was determined by GIAO chemical shift calculations. The reaction pathway for the 1,3-hydroxyl migration of 4 was calculated using quantum mechanical calculations to explain the observed reaction spontaneity. In addition, the unusual downfield shifts observed for H-5 in compounds 4 and 5 were rationalized through geometry optimizations, which indicated the presence of an 8 α -hydroxyl group in 6/6/6/5-fused cyclic sterols.

Supplementary Materials: The following are available online at <http://www.mdpi.com/1660-3397/18/12/607/s1>, Supporting Information I; I. Experimental procedure, II. Computational methods, III. Determination of the C-24 configuration, IV. X-ray crystallography data; Supporting Information II; Figures SII-1–SII-57: ¹H NMR, ¹³C NMR, COSY, HSQC, HMBC, NOESY spectra of 1–10; CIF file for 6.

Author Contributions: A.-Y.S. worked on isolation, structure elucidation, and DFT calculations. H.-S.L., Y.-J.L., and J.S.L. collected the marine sponge and provided advices on the structure elucidation. A.S. and C.C. performed the biological evaluation. J.L. supervised the whole research work. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by grants from the Ministry of Ocean and Fisheries (PM62060) and the Korea Institute of Ocean Science and Technology (PE99821).

Acknowledgments: We are grateful to Byungho Park at EMNI for the technical support and helpful discussions on the quantum mechanical calculations performed in this study.

Conflicts of Interest: The authors declare no competing financial interest.

References

1. Newman, D.J.; Cragg, G.M. Natural products as sources of new drugs over the nearly four decades from 01/1981 to 09/2019. *J. Nat. Prod.* **2020**, *83*, 770–803. [[CrossRef](#)] [[PubMed](#)]
2. Kho, E.; Imagawa, D.K.; Rohmer, M.; Kashman, Y.; Djerassi, C. Sterols in marine invertebrates. 22. Isolation and structure elucidation of conicasterol and theonellasterol, two new 4-methylene sterols from the Red Sea sponges *Theonella conica* and *Theonella swinhoei*. *J. Org. Chem.* **1981**, *46*, 1836–1839. [[CrossRef](#)]
3. Carmely, S.; Kashman, Y. Structure of swinholide-A, a new macrolide from the marine sponge *Theonella swinhoei*. *Tetrahedron Lett.* **1985**, *26*, 511–514. [[CrossRef](#)]
4. Kobayashi, M.; Tanaka, J.; Katori, T.; Kitagawa, I. Marine natural products. XXIII: Three new cytotoxic dimeric macrolides, Swinholides B and C and Isoswinholide A, congeners of Swinholide A, from the Okinawan marine sponge *Theonella swinhoei*. *Chem. Pharm. Bull.* **1990**, *38*, 2960–2966. [[CrossRef](#)]
5. Tsukamoto, S.; Ishibashi, M.; Sasaki, T.; Kobayashi, J.I. New congeners of swinholides from the okinawan marine sponge *Theonella* sp. *J. Chem. Soc. Perkin Trans. 1* **1991**, 3185–3188. [[CrossRef](#)]

6. Dumdei, E.J.; Blunt, J.W.; Munro, M.H.; Pannell, L.K. Isolation of calyculins, calyculinamides, and swinholide H from the New Zealand deep-water marine sponge *Lamellomorpha strongylata*. *J. Org. Chem.* **1997**, *62*, 2636–2639. [[CrossRef](#)]
7. Youssef, D.T.; Mooberry, S.L. Hurghadolide A and Swinholide I, Potent Actin-Microfilament Disrupters from the Red Sea Sponge *Theonella swinhoei*. *J. Nat. Prod.* **2006**, *69*, 154–157. [[CrossRef](#)]
8. Marino, S.D.; Festa, C.; D’Auria, M.V.; Cresteil, T.; Debitus, C.; Zampella, A. Swinholide J, a potent cytotoxin from the marine sponge *Theonella swinhoei*. *Mar. Drugs* **2011**, *9*, 1133–1141. [[CrossRef](#)]
9. Sinisi, A.; Calcinai, B.; Cerrano, C.; Dien, H.A.; Zampella, A.; D’Amore, C.; Renga, B.; Fiorucci, S.; Tagliatalata-Scafati, O. Isoswinholide B and swinholide K, potentially cytotoxic dimeric macrolides from *Theonella swinhoei*. *Bioorg. Med. Chem.* **2013**, *21*, 5332–5338. [[CrossRef](#)]
10. Sakai, R.; Higa, T.; Kashman, Y. Misakinolide-A, an antitumor macrolide from the marine sponge *Theonella* sp. *Chem. Lett.* **1986**, *15*, 1499–1502. [[CrossRef](#)]
11. Kobayashi, J.; Kondo, K.; Ishibashi, M.; Walchli, M.R.; Nakamura, T. Theonezolid A: A novel polyketide natural product from the Okinawan marine sponge *Theonella* sp. *J. Am. Chem. Soc.* **1993**, *115*, 6661–6665. [[CrossRef](#)]
12. Kondo, K.; Ishibashi, M.; Kobayashi, J. Isolation and structures of Theonezolid B and C from the Okinawan marine sponge *Theonella* sp. *Tetrahedron* **1994**, *50*, 8355–8362. [[CrossRef](#)]
13. Ford, P.W.; Gustafson, K.R.; McKee, T.C.; Shigematsu, N.; Maurizi, L.K.; Pannell, L.K.; Williams, D.E.; Dilip de Silva, E.; Lassota, P.; Allen, T.M. Papuamides A–D, HIV-Inhibitory and Cytotoxic Depsipeptides from the Sponges *Theonella mirabilis* and *Theonella swinhoei* Collected in Papua New Guinea. *J. Am. Chem. Soc.* **1999**, *121*, 5899–5909. [[CrossRef](#)]
14. Fusetani, N.; Matsunaga, S.; Matsumoto, H.; Takebayashi, Y. Bioactive marine metabolites. 33. Cyclotheonamides, potent thrombin inhibitors, from a marine sponge *Theonella* sp. *J. Am. Chem. Soc.* **1990**, *112*, 7053–7054. [[CrossRef](#)]
15. Nakao, Y.; Matsunaga, S.; Fusetani, N. Three more cyclotheonamides, C, D, and E, potent thrombin inhibitors from the marine sponge *Theonella swinhoei*. *Bioorg. Med. Chem.* **1995**, *3*, 1115–1122. [[CrossRef](#)]
16. Hamada, T.; Matsunaga, S.; Yano, G.; Fusetani, N. Polytheonamides A and B, Highly cytotoxic, linear polypeptides with unprecedented structural features, from the marine sponge, *Theonella swinhoei*. *J. Am. Chem. Soc.* **2005**, *127*, 110–118. [[CrossRef](#)] [[PubMed](#)]
17. Matsunaga, S.; Fusetani, N. Theonellamides A–E, cytotoxic bicyclic peptides, from a marine sponge *Theonella* sp. *J. Org. Chem.* **1995**, *60*, 1177–1181. [[CrossRef](#)]
18. Wada, S.-I.; Matsunaga, S.; Fusetani, N.; Watabe, S. Theonellamide F, a bicyclic peptide marine toxin, induces formation of vacuoles in 3Y1 rat embryonic fibroblast. *Mar. Biotechnol.* **1999**, *1*, 337–341. [[CrossRef](#)] [[PubMed](#)]
19. Umeyama, A.; Shoji, N.; Enoki, M.; Arihara, S. Swinhosterols A–C, 4-Methylene Secosteroids from the Marine Sponge *Theonella swinhoei*. *J. Nat. Prod.* **1997**, *60*, 296–298. [[CrossRef](#)]
20. Festa, C.; De Marino, S.; D’Auria, M.V.; Bifulco, G.; Renga, B.; Fiorucci, S.; Petek, S.; Zampella, A. Solomonsterols A and B from *Theonella swinhoei*. The first example of C-24 and C-23 sulfated sterols from a marine source endowed with a PXR agonistic activity. *J. Med. Chem.* **2011**, *54*, 401–405. [[CrossRef](#)]
21. Gong, J.; Sun, P.; Jiang, N.; Riccio, R.; Lauro, G.; Bifulco, G.; Li, T.-J.; Gerwick, W.H.; Zhang, W. New steroids with a rearranged skeleton as (h) P300 inhibitors from the sponge *Theonella swinhoei*. *Org. Lett.* **2014**, *16*, 2224–2227. [[CrossRef](#)]
22. Li, J.; Tang, H.; Kurtán, T.; Mándi, A.; Zhuang, C.-L.; Su, L.; Zheng, G.-L.; Zhang, W. Swinhoeisterols from the South China Sea Sponge *Theonella swinhoei*. *J. Nat. Prod.* **2018**, *81*, 1645–1650. [[CrossRef](#)] [[PubMed](#)]
23. Yang, F.; Li, Y.-Y.; Tang, J.; Sun, F.; Lin, H.-W. New 4-methylidene sterols from the marine sponge *Theonella swinhoei*. *Fitoterapia* **2018**, *127*, 279–285. [[CrossRef](#)]
24. De Marino, S.; Ummarino, R.; D’Auria, M.V.; Chini, M.G.; Bifulco, G.; D’Amore, C.; Renga, B.; Mencarelli, A.; Petek, S.; Fiorucci, S. 4-Methylenesterols from *Theonella swinhoei* sponge are natural pregnane-X-receptor agonists and farnesoid-X-receptor antagonists that modulate innate immunity. *Steroids* **2012**, *77*, 484–495. [[CrossRef](#)]
25. De Marino, S.; Ummarino, R.; D’Auria, M.V.; Chini, M.G.; Bifulco, G.; Renga, B.; D’Amore, C.; Fiorucci, S.; Debitus, C.; Zampella, A. Theonellasterols and conicasterols from *Theonella swinhoei*. Novel marine natural ligands for human nuclear receptors. *J. Med. Chem.* **2011**, *54*, 3065–3075. [[CrossRef](#)] [[PubMed](#)]

26. Chini, M.G.; Jones, C.R.; Zampella, A.; D'Auria, M.V.; Renga, B.; Fiorucci, S.; Butts, C.P.; Bifulco, G. Quantitative NMR-derived interproton distances combined with quantum mechanical calculations of ^{13}C chemical shifts in the stereochemical determination of conicasterol F, a nuclear receptor ligand from *Theonella swinhoei*. *J. Org. Chem.* **2012**, *77*, 1489–1496. [[CrossRef](#)] [[PubMed](#)]
27. Sepe, V.; Ummarino, R.; D'Auria, M.V.; Chini, M.G.; Bifulco, G.; Renga, B.; D'Amore, C.; Debitus, C.; Fiorucci, S.; Zampella, A. Conicasterol E, a small heterodimer partner sparing farnesoid X receptor modulator endowed with a pregnane X receptor agonistic activity, from the marine sponge *Theonella swinhoei*. *J. Med. Chem.* **2012**, *55*, 84–93. [[CrossRef](#)]
28. Qureshi, A.; Faulkner, D.J. 7 α -Hydroxytheonellasterol, a cytotoxic 4-methylene sterol from the Philippines sponge *Theonella swinhoei*. *J. Nat. Prod.* **2000**, *63*, 841–842. [[CrossRef](#)]
29. Guo, J.-K.; Chiang, C.-Y.; Lu, M.-C.; Chang, W.-B.; Su, J.-H. 4-Methylenesterols from a Sponge *Theonella swinhoei*. *Mar. Drugs* **2012**, *10*, 1536–1544. [[CrossRef](#)]
30. Sepe, V.; Ummarino, R.; D'Auria, M.V.; Tagliatalata-Scafati, O.; Marino, S.D.; D'Amore, C.; Renga, B.; Chini, M.G.; Bifulco, G.; Nakao, Y. Preliminary structure-activity relationship on theonellasterol, a new chemotype of FXR antagonist, from the marine sponge *Theonella swinhoei*. *Mar. Drugs* **2012**, *10*, 2448–2466. [[CrossRef](#)]
31. Pu, J.-X.; Huang, S.-X.; Ren, J.; Xiao, W.-L.; Li, L.-M.; Li, R.-T.; Li, L.-B.; Liao, T.-G.; Lou, L.-G.; Zhu, H.-J. Isolation and Structure Elucidation of Kadlongilactones C–F from *Kadsura longipedunculata* by NMR Spectroscopy and DFT Computational Methods. *J. Nat. Prod.* **2007**, *70*, 1706–1711. [[CrossRef](#)] [[PubMed](#)]
32. Brown, S.G.; Jansma, M.J.; Hoye, T.R. Case study of empirical and computational chemical shift analyses: Reassignment of the relative configuration of phomopsichalasin to that of diaporthichalasin. *J. Nat. Prod.* **2012**, *75*, 1326–1331. [[CrossRef](#)] [[PubMed](#)]
33. Mun, B.; Wang, W.; Kim, H.; Hahn, D.; Yang, I.; Won, D.H.; Kim, E.-H.; Lee, J.; Han, C.; Kim, H. Cytotoxic 5 α , 8 α -epidioxo sterols from the marine sponge *Monanchora* sp. *Arch. Pharm. Res.* **2015**, *38*, 18–25. [[CrossRef](#)] [[PubMed](#)]
34. Gunatilaka, A.L.; Gopichand, Y.; Schmitz, F.J.; Djerassi, C. Minor and trace sterols in marine invertebrates. 26. Isolation and structure elucidation of nine new 5 α , 8 α -epidoxo sterols from four marine organisms. *J. Org. Chem.* **1981**, *46*, 3860–3866. [[CrossRef](#)]
35. Vien, L.T.; Hanh, T.T.H.; Hong, P.T.; Thanh, N.V.; Huong, T.T.; Cuong, N.X.; Nam, N.H.; Thung, D.C.; Kiem, P.V.; Minh, C.V. Polar steroid derivatives from the Vietnamese starfish *Astropecten polyacanthus*. *Nat. Prod. Res.* **2018**, *32*, 54–59. [[CrossRef](#)] [[PubMed](#)]
36. Wang, W.; Li, F.; Park, Y.; Hong, J.; Lee, C.-O.; Kong, J.Y.; Shin, S.; Im, K.S.; Jung, J.H. Bioactive Sterols from the Starfish *Certonardoa semiregularis*. *J. Nat. Prod.* **2003**, *66*, 384–391. [[CrossRef](#)]
37. Zhang, H.J.; Yi, Y.H.; Lin, H.W. Oxygenated 4-Methylidene Sterols from the South China Sea Sponge *Theonella swinhoei*. *Helv. Chim. Acta* **2010**, *93*, 1120–1126. [[CrossRef](#)]
38. Sright, J.; McInnes, A.; Shimizu, S.; Smith, D.; Walter, J.; Idler, D.; Khalil, W. Identification of C-24 alkyl epimers of marine sterols by ^{13}C nuclear magnetic resonance spectroscopy. *Can. J. Chem.* **1978**, *56*, 1898–1903. [[CrossRef](#)]

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).