

Supporting Information

Antitumor Effects of a Sesquiterpene Derivative from Marine Sponge in Human

Breast Cancer Cells

Li-Yuan Bai ^{1,2,†}, Jui-Hsin Su ^{3,†}, Chang-Fang Chiu ^{1,4}, Wei-Yu Lin ⁵, Jing-Lan Hu ¹, Chia-Hsien Feng ⁶, Chih-Wen Shu ⁷, and Jing-Ru Weng ^{8,9,10,*}

¹Division of Hematology and Oncology, Department of Internal Medicine, China Medical University Hospital, Taichung 40447, Taiwan

²College of Medicine, China Medical University, Taichung 40402, Taiwan

³National Museum of Marine Biology and Aquarium, Pingtung 94450, Taiwan

⁴Cancer Center, China Medical University Hospital, Taichung 40415, Taiwan

⁵Department of Pharmacy, Kinmen Hospital, the Ministry of Health and Welfare, Kinmen 89142, Taiwan

⁶Department of Fragrance and Cosmetic Science, College of Pharmacy, Kaohsiung Medical University, Kaohsiung 80708, Taiwan

⁷Institute of BioPharmaceutical Sciences, National Sun Yat-sen University, Kaohsiung 80424, Taiwan

⁸Department of Marine Biotechnology and Resources, National Sun Yat-sen University, Kaohsiung 80424, Taiwan

⁹Doctoral Degree Program in Marine Biotechnology, National Sun Yat-sen
University, Kaohsiung 80424, Taiwan

¹⁰Graduate Institute of Natural Products, College of Pharmacy, Kaohsiung Medical
University, Kaohsiung 80708, Taiwan

Figure	Page
<p>S1 Localization analysis of (A) E-cadherin and (B) β-catenin in MCF-7 cells after ilimaquinone (5 μM) treatment. Cells were incubated with DMSO or ilimaquinone (5 μM) for 48 h, then washed three with PBS, fixed with 4% paraformaldehyde (PFA) and immunostained by antibodies against E-cadherin and β-catenin. Bar: 100 μm.</p>	4

Fig. S1

