

Highlighting the biotechnological potential of deep oceanic crust fungi through the prism of their antimicrobial activity

Maxence Quemener¹, Marie Dayras², Nicolas Frotté¹, Stella Debaets¹, Christophe Lemeur¹, Georges Barbier¹, Virginia Edgcomb³, Mohamed Mehiri², Gaëtan Burgaud^{1*}

¹ Univ Brest, Laboratoire Universitaire de Biodiversité et Ecologie Microbienne, France

² Université Côte d'Azur, CNRS, Institut de Chimie de Nice, UMR 7272, Marine Natural Products Team, 06108, Nice, France.

³ Departments of Geology and Geophysics and Biology, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA

*Correspondence: gaetan.burgaud@univ-brest.fr; Tel.: +33290915148

Supplementary Figure S1: Comparative analyses of the HPLC-PDA-ELSD chemical fingerprints of the crude organic extracts F1 of a) (UBOCC-A-118169 *Penicillium olsonii*), b) (UBOCC-A-119011 *Penicillium olsonii*) and c) (UBOCC-A-118156 *Aspergillus conicus*) dichloromethane-ethyl acetate extracts obtained with HPLC-DAD-ELSD (Macherey-Nagel propylphenyl-C18, 250 x 4.6 mm; 5 μ m, gradient: CH₃CN + 0.1% formic acid (FA), H₂O + 0.1% FA, 10:90 to 100:0 in 30 mn).

