

Article

Cosmetic Application of Cyanobacteria Extracts with a Sustainable Vision to Skincare: Role in the Antioxidant and Antiaging Process

Janaína Morone ^{1,2}, Graciliana Lopes ¹, João Morais ^{1,2}, Jorge Neves ¹, Vítor Vasconcelos ^{1,2}
and Rosário Martins ^{1,3,*}

- ¹ CIIMAR/CIMAR—Interdisciplinary Centre of Marine and Environmental Research, University of Porto, Terminal de Cruzeiros do Porto de Leixões, Av. General Norton de Matos s/n, 4450-208 Matosinhos, Portugal
² FCUP—Faculty of Sciences, University of Porto, Rua do Campo Alegre, Edifício FC4, 4169-007 Porto, Portugal
³ Health and Environment Research Centre, School of Health, Polytechnic Institute of Porto, Rua Dr. António Bernardino de Almeida, 400, 4200-072 Porto, Portugal
* Correspondence: mrm@ess.ipp.pt

Abstract: Nature-based and sustainably sourced cosmetics have been dominating the area of skincare products worldwide. Due to their antioxidant and antiaging properties, compounds from cyanobacteria, such as carotenoids and phycobiliproteins, may replace synthetic ingredients in cosmetic formulations and may be used in products such as sunscreens, skincare creams, and makeup. In this study, we evaluated the potential of acetonic and aqueous extracts from cyanobacteria strains of the genera *Cyanobium* and *Leptothoe* and from strains within Synechococcales and Oscillatoriales orders, for use in cosmetics. Extractions were sequentially performed with acetone and water. Extracts were firstly analyzed for their toxicity to keratinocytes, fibroblasts, and endothelial cells (HaCAT, 3T3L1 and hCMEC/D3, respectively). The non-cytotoxic extracts were characterized in terms of total proteins, carotenoids, chlorophyll, phenols, phycobiliproteins, and analyzed for their antioxidant potential against the superoxide anion radical ($O_2^{\bullet-}$), and for their ability to inhibit key enzymes associated with the skin aging process. Aqueous extracts were richer in total proteins and phycobiliproteins. The aqueous extracts of Synechococcales cyanobacterium LEGE 181157 and Synechococcales cyanobacterium LEGE 181150 showed the highest value for total proteins (760.81 and 695.25 $\mu\text{g BSA mL}^{-1}$ dry extract, respectively) and the best values regarding $O_2^{\bullet-}$ scavenging ($IC_{50} = 63.24$ and $112.18 \mu\text{g mL}^{-1}$ dry extract, respectively) with a significant negative correlation observed ($p < 0.01$). Moreover, aqueous extracts of Synechococcales cyanobacterium LEGE 181150 and Synechococcales cyanobacterium LEGE 181157 inhibited hyaluronidase, (IC_{50} of 483.86 and $645.06 \mu\text{g mL}^{-1}$ dry extract, respectively), with a significant negative correlation with total proteins ($p < 0.05$), pointing out the contribution of these compounds to the biological activities observed. Acetonic extracts were richer in carotenoids and phenols. Zeaxanthin and β -carotene were predominant among all strains, being present in higher amount in *Cyanobium* sp. LEGE 07175 ($53.08 \mu\text{g mg}^{-1}$) and *Leptothoe* sp. LEGE 181156 ($47.89 \mu\text{g mg}^{-1}$), respectively. The same strains also showed the highest values for collagenase inhibition at $750 \mu\text{g mL}^{-1}$ dry extract (32.88 and 36.61%, respectively). Furthermore, *Leptothoe* sp. LEGE 181156 exhibited the lowest IC_{50} value for tyrosinase inhibition ($465.92 \mu\text{g mL}^{-1}$ dry extract) and Synechococcales cyanobacterium LEGE 181157 presented the best values for elastase inhibition (IC_{50} of 380.50 and IC_{25} of $51.43 \mu\text{g mL}^{-1}$ dry extract). In general, cyanobacteria extracts demonstrated potential for being used for antiaging purposes, with aqueous extracts being more efficient at free radicals scavenging and acetonic ones at avoiding degradation of dermal matrix components.

Keywords: cyanobacteria; cosmetics; carotenoids; phycobiliproteins; phenols; metalloproteinases; oxidative stress; antiaging



Citation: Morone, J.; Lopes, G.; Morais, J.; Neves, J.; Vasconcelos, V.; Martins, R. Cosmetic Application of Cyanobacteria Extracts with a Sustainable Vision to Skincare: Role in the Antioxidant and Antiaging Process. *Mar. Drugs* **2022**, *20*, 761. <https://doi.org/10.3390/md20120761>

Academic Editor: Alessandro Concas

Received: 7 November 2022

Accepted: 29 November 2022

Published: 2 December 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Thriving cosmetics properties include skin protection against chemical and physical factors such as xenobiotics, ultraviolet radiation (UVR), and dehydration, the main factors involved in skin's loss of integrity and aging. Despite being a result of natural physiological features, the skin aging process may be accelerated by mechanisms such as oxidative stress, which settles down with the accumulation of free radicals. Oxidative stress is mainly responsible for triggering the activation of matrix metalloproteinases (MMPs), namely collagenases, elastases, and hyaluronidases, that contribute to the degradation of collagen, elastin, and hyaluronic acid (HA), as the main components of the extracellular matrix (ECM), which are essential for the maintenance of skin elasticity, firmness, and regeneration [1]. Consequently, the aesthetic appearance of the skin becomes compromised, which affects health, social well-being, and self-esteem. Thus, skincare habits have become a priority among people who seek a healthier and more pleasing lifestyle.

Cosmetic ingredients derived from photosynthetic organisms are a beneficial approach to prioritize the use of natural raw materials into final products that respect the skin's properties and the environment and, consequently, make a commitment to sustainability [2]. Among photosynthetic organisms, the interest in cyanobacteria has been growing, with different species being explored from the natural cosmetics point of view. Cyanobacteria are widespread organisms found in all ecosystems, some with life-threatening conditions such as high temperatures, high salinity, and extreme solar radiation [3]. As extremely adaptable organisms, cyanobacteria have developed a remarkable secondary metabolism, resulting in an arsenal of metabolites with proven pharmaceutical, nutraceutical, and food-conservation potential. Considering skin-related needs, the production of carotenoids, phenolic compounds, phycobiliproteins (PBPs), mycosporin-like amino acids (MAAs), and scytonemin (SCY) places these organisms at the forefront of healthy skincare. All these molecules play a central role as antiaging ingredients, mainly due to their antioxidant potential, sun-protection capacity, and ability to inhibit enzymes responsible for the degradation of the ECM [2,4].

The use of cyanobacteria-derived compounds in skincare products can be expressed in sunscreens, moisturizing creams, formulations with antioxidant and anti-inflammatory properties, as well as in pigmentation disorders, dermatophytosis, and infectious diseases [1,2,5–7]. Recent studies focusing on the potential of cyanobacteria in cosmetics have reported the antioxidant activity attributed to the production of carotenoids from some genera, namely *Leptolyngbya*, *Synechocystis*, and *Wolleea* [2,8], PBP such as phycocyanin (PC) and phycoerythrin (PE) from *Arthrospira* sp. and *Spirulina* sp. [8,9], and phenolic compounds from *Nostoc commune* [10]. Considering the inhibition of MMPs compounds such as mycosporin-2-glycine isolated from a halotolerant cyanobacterium was found to inhibit collagenase activity [11], cyclodepsipeptide from *Schizothrix* sp. and *Coleofasciculus* sp. presented high elastase inhibitory properties [12], and a polyssaccharide from *Nostochopsis lobatus* showed potent hyaluronidase inhibitory activity [13]. Additionally, Favas and co-workers recently showed that some cyanobacteria aqueous extracts had the potential to retard dermal density loss [14]. Additionally, the extract from *Phormidium persicinum* (PHORMISKIN Bioprotech G[®]) has been commercially used as a melanin synthesis reducer [15]. The studies involving cyanobacteria as photoprotectors have also been highlighted [16]. It is known that cyanobacteria synthesize pigments, which are UVR-absorbing compounds, namely carotenoids, PBPs, MAAs, and SCY [16]. The available literature produced so far highlights the biotechnological potential of cyanobacteria as natural and eco-friendly sources of compounds in the field of skincare and triggers further exploitation in this field. Besides being self-renewable, with low nutritional requirements, and presenting high growth and photosynthesis rates, the sustainable cultivation of cyanobacteria is guaranteed [2,17].

In the present work, seven cyanobacteria strains isolated from Portuguese and Cape Verde marine ecosystems (*Cyanobium* sp. LEGE 07175, *Synechococcales* cyanobacterium LEGE 181150, *Leptothoe* sp. LEGE 181155, *Leptothoe* sp. LEGE 181156, *Synechococcales*

cyanobacterium LEGE 181157, Synechococcales cyanobacterium LEGE 181158, and Oscillatoriales cyanobacterium LEGE 181159) were explored for their potential to slow down skin aging. Acetonic and aqueous extracts were evaluated for cytotoxicity in skin cell-lines, phytochemical composition, antioxidant potential, and inhibitory effect in skin key enzymes.

2. Results

2.1. Cytotoxicity

Cytotoxic effects were registered in all cell lines for both acetonic and aqueous extracts of the strain *Leptothoe* sp. LEGE 181155 (g and n in Figures S1–S3) and for the acetonic extract of strain Oscillatoriales cyanobacterium LEGE 181159 (m in Figures S1–S3), with a significant decrease from the lowest concentration ($12.5 \mu\text{g mL}^{-1}$) ($p < 0.05$) after 24 h. For all other strains, none of the extracts showed cytotoxic effects, with no significant differences from the control ($p < 0.05$) for both 24 and 48 h of exposition. Based on these results, the extracts of the strain *Leptothoe* sp. LEGE 181155 and the acetone extract of Oscillatoriales cyanobacterium LEGE 181159 were excluded from the study (Results and Supplementary Material).

2.2. Phytochemical Analysis

For the purpose of comparing the chemical profiles and establishing a relationship between the chemical composition and the biological activities assessed herein, the acetonic and aqueous extracts were characterized in terms of phenolic content, total proteins, and PBP. Moreover, the qualitative and quantitative carotenoids profile was also established.

2.2.1. Carotenoids

The HPLC-PDA analysis of the acetonic extract from the different cyanobacteria strains allowed the detection of 13 carotenoids, 2 chlorophyll-*a* derivatives, and chlorophyll-*a*, and for the aqueous extract 8 carotenoids, 2 chlorophyll-*a* derivatives and chlorophyll-*a* (Table 1). The chromatographic profile of the three different genera is illustrated in Figure 1.

The identified compounds for aqueous extracts consisted of two xanthophylls: zeaxanthin (5) and myxoxanthophyll (7); one carotene: β -carotene (13), chlorophyll-*a* (4) and the chlorophyll-*a* derivative phaeophytin-*a* (14). For acetonic extracts, the identified compounds consisted of four xanthophylls: lutein (2), zeaxanthin (5), myxoxanthophyll (7) and echinenone (10); two carotenes: β -carotene (13) and γ -carotene (16), as well as chlorophyll-*a* (4) and phaeophytin-*a* (14). Seven compounds with the same spectra as the identified carotenoids, however, with different retention times to the standard, were also detected in some strains, being defined as unidentified carotenoids (1, 3, 8, 9, 11, 12, 15), and the same was carried out for the chlorophyll-*a* derivative (6). Regarding aqueous extracts, five compounds were defined as unidentified carotenoids (1, 3, 9, 12, 15).

For aqueous extracts, total carotenoid concentration ranged between 0.05 and $3.00 \mu\text{g mg}^{-1}$ of dry extract. *Cyanobium* sp. LEGE 07175 showed the highest carotenoids content, followed by Synechococcales cyanobacterium LEGE 181150 and Oscillatoriales cyanobacterium LEGE 181159 (3.00 ; 1.55 and $0.84 \mu\text{g mg}^{-1}$, respectively) ($p < 0.05$). All strains contain β -carotene (13), Zeaxanthin (5), and phaeophytin-*a* (14), but in *Cyanobium* sp. LEGE 07175, this chlorophyll-*a* derivative could not be quantified. Myxoxanthophyll (7) was not found in two of six strains. Lutein (2) and γ -carotene (16) were not identified in any aqueous extract.

Table 1. Carotenoid and chlorophyll contents ($\mu\text{g mg}^{-1}$ dry extract) in the aqueous and acetic extracts of the cyanobacteria strains, determined by HPLC-PDA ^{1,2}.

Peak	Compound	RT (min)	Cyanobium sp. LEGE 07175		Synechococcales cyanobacterium LEGE 181150		Leptothoe sp. LEGE 181156		Synechococcales cyanobacterium LEGE 181157		Synechococcales cyanobacterium LEGE 181158		Oscillatoriales cyanobacterium LEGE 191159
			Aqueous Extract	Acetonic Extract	Aqueous Extract	Acetonic Extract	Aqueous Extract	Acetonic Extract	Aqueous Extract	Acetonic Extract	Aqueous Extract	Acetonic Extract	Aqueous Extract
1	Unidentified carotenoid	13.35	nq	7.28 ± 0.03 ^a	nd	2.55 ± 0.04 ^c	nd	3.31 ± 0.07 ^b	nd	nd	nd	nd	nd
2	Lutein	13.44	nd	nd	nd	nd	nd	nd	nd	2.07 ± 0.70	nd	1.96 ± 0.15	nd
3	Unidentified carotenoid	14.81	0.53 ± 0.05	21.03 ± 0.40 ^a	0.04 ± 0.02	7.75 ± 0.39 ^b	nq	8.53 ± 0.03 ^b	nd	nd	nd	0.77 ± 0.12 ^c	nd
4	Chlorophyll- <i>a</i>	17.02	nd	nd	14.77 ± 1.21 ^A	32.19 ± 1.02	0.81 ± 0.06 ^C	nd	0.06 ± 0.002	nd	9.42 ± 1.01 ^B	31.96 ± 2.04	8.69 ± 0.07 ^B
5	Zeaxanthin	17.15	1.58 ± 0.08	53.08 ± 1.48 ^a	nd	nd	nd	39.85 ± 2.69 ^b	nd	24.38 ± 0.33 ^c	nd	nd	nd
6	Chlorophyll- <i>a</i> derivative	17.98	nd	25.27 ± 1.17 ^b	nd	29.23 ± 0.69 ^{a,b}	nd	34.81 ± 1.52 ^a	nd	12.69 ± 2.04 ^c	nq	21.90 ± 3.55 ^b	nd
7	Myxoxanthophyll	21.07	nd	nd	0.29 ± 0.03 ^A	8.22 ± 0.37 ^a	nd	4.07 ± 0.38 ^b	0.01 ± 0.002 ^C	5.41 ± 0.28 ^{a,b}	0.17 ± 0.05 ^{A,B}	8.43 ± 1.41 ^a	0.16 ± 0.01 ^B
8	Unidentified carotenoid	22.18	nd	1.40 ± 0.18 ^a	nd	2.13 ± 0.07 ^a	nd	1.27 ± 0.02 ^b	nd	1.15 ± 0.03 ^b	nd	1.76 ± 0.41 ^a	nd
9	Unidentified carotenoid	25.89	nq	1.21 ± 0.15 ^a	nd	0.70 ± 0.01 ^b	nd	1.37 ± 0.17 ^a	nd	0.26 ± 0.08 ^b	nd	nd	nd
10	Echinenone	27.01	nd	1.17 ± 0.07	nd	nd	nd	nd	nd	nd	nd	nd	nd
11	Unidentified carotenoid	28.93	nd	1.10 ± 0.04 ^b	nd	nd	nd	2.02 ± 0.24 ^a	nd	1.15 ± 0.14 ^b	nd	1.00 ± 0.08 ^b	nd
12	Unidentified carotenoid	29.7	nq	0.41 ± 0.16 ^b	nq	1.15 ± 0.07 ^a	nd	nd	nd	0.26 ± 0.09 ^b	nd	nd	nq
13	β -Carotene	34.32	0.88 ± 0.11 ^B	19.58 ± 0.64 ^c	1.22 ± 0.11 ^A	25.13 ± 0.75 ^{b,c}	0.05 ± 0.002 ^D	47.89 ± 2.81 ^a	0.10 ± 0.002 ^D	29.30 ± 0.56 ^b	0.54 ± 0.05 ^C	30.57 ± 1.63 ^b	0.68 ± 0.01 ^{B,C}
14	Phaeophytin- <i>a</i>	34.69	nq	5.81 ± 1.08 ^a	nd	2.25 ± 0.61 ^b	nd	1.81 ± 0.11 ^b	nd	5.23 ± 0.27 ^a	nd	6.31 ± 0.33 ^a	nd
15	Unidentified carotenoid	35.05	nq	3.92 ± 0.18 ^c	nq	5.23 ± 0.64 ^{a,b}	nq	6.51 ± 0.24 ^a	nq	3.37 ± 0.11 ^c	nq	4.53 ± 0.60 ^{b,c}	nq
16	γ -Carotene	41.78	nd	nq	nd	35.96 ± 3.00 ^a	nd	33.52 ± 3.79 ^a	nd	32.11 ± 1.47 ^a	nd	26.51 ± 1.99 ^b	nd
	Total carotenoids		3.00 ± 0.01 ^A	115.98 ± 0.48 ^c	1.55 ± 0.01 ^B	123.26 ± 0.85 ^b	0.05 ± 0.01 ^F	150.15 ± 1.40 ^a	0.18 ± 0.01 ^E	104.71 ± 0.42 ^d	0.71 ± 0.01 ^D	113.81 ± 0.81 ^c	0.84 ± 0.01 ^C
	Total chlorophylls		nd	25.27 ± 1.17 ^{a,b}	14.77 ± 1.21 ^A	29.23 ± 0.69 ^{a,b}	0.81 ± 0.06 ^C	34.81 ± 1.52 ^a	nd	12.69 ± 2.04 ^c	9.42 ± 1.01 ^B	21.90 ± 3.55 ^{b,c}	8.69 ± 0.07 ^B

¹ Values are expressed as mean ± SD of two determinations. ² Different superscript letters in the same row denote statistical differences at $p < 0.05$ (ANOVA, Tukey's HSD); uppercase, Aqueous extracts; lowercase, Acetonic extracts. nd: Not detected. nq: Not quantified.




Figure 1. Carotenoid and chlorophyll profile of acetonic extracts of cyanobacteria strains. HPLC-PDA recorded at 450 nm. Unidentified carotenoids (1,3,8,9,11,12,15), Zeaxanthin (5), Chlorophyll-*a* derivative (6), Myxoxanthophyll (7), Echinenone (10), β -carotene (13), Phaeophytin-*a* (14), and γ -carotene (16).

According to chlorophylls, the total chlorophylls content ranged from 0.81 up to 14.77 $\mu\text{g mg}^{-1}$. Chlorophyll-*a* (4) was found in all strains, except in *Cyanobium sp. LEGE 07175*, where only a chlorophyll-*a* derivative (6) was found. The highest chlorophylls content was found in *Synechococcales cyanobacterium LEGE 181150* (14.77 $\mu\text{g mg}^{-1}$) ($p < 0.05$), followed by *Synechococcales cyanobacterium LEGE 181158* and *Oscillatoriales cyanobacterium LEGE 181159* (9.42 and 8.69 $\mu\text{g mg}^{-1}$), with no significant differences.

Regarding acetonic extract, the total carotenoid concentration ranged between 104.71 and 150.15 $\mu\text{g mg}^{-1}$ of dry extract, and it differed significantly. The highest carotenoids content was found in *Leptothoe sp. LEGE 181156* followed by *Synechococcales cyanobacterium LEGE 181150* and *Cyanobium sp. LEGE 07175* (150.15; 123.26; and 115.98 $\mu\text{g mg}^{-1}$, respectively) ($p < 0.05$). Lutein (2) was found in two strains: *Synechococcales cyanobacterium LEGE 181157* and *Synechococcales cyanobacterium LEGE 181158*. Zeaxanthin (5) was identified in all strains, but in *Synechococcales cyanobacterium LEGE 181150* and *Synechococcales cyanobacterium LEGE 181158*, this xanthophyll appeared co-eluted with chlorophyll-*a* (4). The highest content of this xanthophyll was found in *Cyanobium sp. LEGE 07175* (53.08 $\mu\text{g mg}^{-1}$) ($p < 0.05$). Even though *Cyanobium sp. LEGE 07175* was the only one that did not present myxoxanthophyll (7), it was the only strain in which echinenone (10) was identified. All the other strains presented β -carotene (13), phaeophytin-*a* (14), and γ -carotene (16) in their profile. The highest content of β -carotene was found in the

strain *Leptothoe* sp. LEGE 181156 ($47.89 \mu\text{g mg}^{-1}$) ($p < 0.05$) followed by Synechococcales cyanobacterium LEGE 181158 and Synechococcales cyanobacterium LEGE 181157 (30.57 and $29.30 \mu\text{g mg}^{-1}$, respectively). The highest phaeophytin-*a* content was detected in Synechococcales cyanobacterium LEGE 181158, followed by *Cyanobium* sp. LEGE 07175 and Synechococcales cyanobacterium LEGE 181157 (6.31 ; 5.81 ; and $5.23 \mu\text{g mg}^{-1}$, respectively) with no significant differences (Table 1 and Figure 1).

Regarding chlorophylls, the total chlorophylls content ranged from 12.69 up to $34.81 \mu\text{g mg}^{-1}$. Chlorophyll-*a* (4) derivative was identified in all strains of acetonetic extracts in considerably high amounts when compared with chlorophyll-*a*. Synechococcales cyanobacterium LEGE 181156 presented the highest content, followed by Synechococcales cyanobacterium LEGE 181150 and Synechococcales cyanobacterium LEGE 181158 (34.81 ; 29.23 and $21.90 \mu\text{g mg}^{-1}$, respectively) with no significant differences. Additionally, in two strains, namely Synechococcales cyanobacterium LEGE 181150 and Synechococcales cyanobacterium LEGE 181158, chlorophyll-*a* (4) was also present; however, it was co-eluted with zeaxanthin (5). Amongst the pigments analyzed, carotenoids were dominant over chlorophylls in all strains for acetonetic extraction. Furthermore, zeaxanthin, phaeophytin-*a*, and γ -carotene were identified in the acetonetic extracts of all the species studied (Figure 1, Table 1).

Acetonetic extracts were richer in pigments than aqueous extracts, considering both carotenoids and chlorophylls. The unique carotenoid identified for all strains, in both extractions, was β -carotene. Moreover, the carotene echinenone was only identified in *Cyanobium* sp. LEGE 07175.

2.2.2. Total Phenolic Content (TPC)

For both extracts, TPC was measured through the Folin–Ciocalteu colorimetric assay; the results were expressed in μg of GAEs per mg of dry extract and are displayed in Table 2.

Table 2. Total Phenolic Content (TPC) and Total Proteins present in cyanobacterial extracts ^{1,2}.

Strains	TPC (μg (GAE) mg^{-1} dry extract)		Total Proteins (μg (BSA) mg^{-1} dry extract)	
	Aqueous Extracts	Acetonetic Extracts	Aqueous Extracts	Acetonetic Extracts
<i>Cyanobium</i> sp. LEGE 07175	$14.79^{a,b} \pm 0.20$	$23.11^{a,b} \pm 0.69$	$309.63^e \pm 4.26$	$167.49^b \pm 4.93$
Synechococcales cyanobacterium LEGE 181150	$14.49^{a,b} \pm 0.48$	$20.14^{b,c} \pm 0.16$	$695.25^b \pm 10.35$	$199.37^a \pm 6.38$
<i>Leptothoe</i> sp. LEGE 181156	$14.93^{a,b} \pm 0.27$	$27.13^a \pm 0.07$	$463.70^d \pm 8.78$	158.67 ± 2.58
Synechococcales cyanobacterium LEGE 181157	$14.92^{a,b} \pm 0.66$	$17.86^c \pm 0.06$	$760.81^a \pm 15.72$	$128.84^c \pm 7.12$
Synechococcales cyanobacterium LEGE 181158	$15.67^a \pm 0.23$	$24.71^{a,b} \pm 0.53$	$579.99^c \pm 12.36$	$209.31^a \pm 4.36$
Oscillatoriales cyanobacterium LEGE 181159	$11.46^b \pm 0.67$	–	$550.59^c \pm 18.14$	–

¹ Mean \pm SD of three independent experiments. ² Different superscript letters in the same column correspond to statistical differences at $p < 0.05$ (ANOVA, Tukey's HSD). – not evaluated.

Acetonetic extracts presented higher TPC values than aqueous extracts. The highest TPC was found in the acetonetic extract of *Leptothoe* sp. LEGE 181156, followed by Synechococcales cyanobacterium LEGE 181158 and *Cyanobium* sp. LEGE 07175 (27.13 ; 24.71 ; and $23.11 \mu\text{g GAE mg}^{-1}$ dry extract, respectively) with not significant differences.

Regarding aqueous extracts, the values ranged from 11.46 up to $15.67 \mu\text{g GAE mg}^{-1}$ dry extract, Synechococcales cyanobacterium LEGE 181158 being the strain with the highest TPC, with no significant differences among the samples, except between Oscillatoriales cyanobacterium LEGE 181159, which showed lower content, and Synechococcales cyanobacterium LEGE 181158 (Table 2).

2.2.3. Total Proteins

The total protein content of both extracts was evaluated by the method of BSA, using the protein assay kit, and the results were expressed in μg of BSA per mg of dry extract

(Table 2). Higher values were found in the aqueous extraction, in which Synechococcales cyanobacterium LEGE 181157 presented the highest content, followed by Synechococcales cyanobacterium LEGE 181150 and Synechococcales cyanobacterium LEGE 181158 (760.81; 695.25 and 579.99 $\mu\text{g BSA mg}^{-1}$ dry extract, respectively) ($p < 0.05$).

Regarding acetonetic extracts, Synechococcales cyanobacterium LEGE 181158 showed the highest content, followed by Synechococcales cyanobacterium LEGE 181150 (209.31 and 199.37 $\mu\text{g BSA mg}^{-1}$ dry extract, respectively) with no significant difference.

2.2.4. Phycobiliproteins (PBPs)

As water-soluble proteins, PBPs were quantified only in aqueous extracts. Regarding PC, the Synechococcales cyanobacterium LEGE 181150 extract presented the highest content, followed by Oscillatoriales cyanobacterium LEGE 181159 and Synechococcales cyanobacterium LEGE 181158 (222.76; 201.15; and 194.43 $\mu\text{g mg}^{-1}$ dry extract, respectively) ($p < 0.05$). In contrast, Synechococcales cyanobacterium LEGE 181157 possessed the highest content of PE, followed by *Leptothoe* sp. LEGE 181156 and Oscillatoriales cyanobacterium LEGE 181159 (275.04; 159.05 and 26.33 $\mu\text{g mg}^{-1}$ dry extract, respectively) ($p < 0.05$) (Table 3).

Table 3. Phycobiliproteins content ($\mu\text{g mg}^{-1}$ dry extract) in cyanobacteria aqueous extracts ^{1,2}.

Strains	Phycobiliproteins ($\mu\text{g mg}^{-1}$ dry aqueous extract)		
	Phycocyanin	Allophycocyanin	Phycoerythrin
<i>Cyanobium</i> sp. LEGE 07175	18.21 ^f \pm 1.36	11.26 ^e \pm 1.20	8.06 ^e \pm 1.17
Synechococcales cyanobacterium LEGE 181150	222.76 ^a \pm 0.79	57.29 ^b \pm 0.17	17.31 ^d \pm 0.11
<i>Leptothoe</i> sp. LEGE 181156	76.10 ^e \pm 0.13	37.05 ^d \pm 0.28	159.05 ^b \pm 0.14
Synechococcales cyanobacterium LEGE 181157	117.94 ^d \pm 0.12	44.57 ^c \pm 0.28	275.04 ^a \pm 0.30
Synechococcales cyanobacterium LEGE 181158	194.43 ^c \pm 0.39	44.52 ^c \pm 0.34	17.42 ^d \pm 0.22
Oscillatoriales cyanobacterium LEGE 181159	201.15 ^b \pm 0.78	65.90 ^a \pm 0.67	26.33 ^c \pm 0.16

¹ Values are expressed as the mean \pm SD of three independent experiments. ² Different superscript letters in the same column correspond to statistical differences at $p < 0.05$.

The strong blue color of PC was notable in the aqueous extracts of Synechococcales cyanobacterium LEGE 181150, Oscillatoriales cyanobacterium LEGE 181159, and Synechococcales cyanobacterium LEGE 181158; and the strong pink/red color of PE was observed in the aqueous extracts of Synechococcales cyanobacterium LEGE 181157 and *Leptothoe* sp. LEGE 181156, which is in accordance with the main phycobiliproteins quantified in each strain.

2.3. Biological Activities

Superoxide Anion Radical ($\text{O}_2^{\bullet-}$) Scavenging

The results of $\text{O}_2^{\bullet-}$ scavenging capacity of both cyanobacteria extracts are summarized in Table 4 and Figure 2. Aqueous extracts were more effective and presented lower IC_{50} values than acetonetic extracts towards radical scavenging. Synechococcales cyanobacterium LEGE 181157 was the most effective strain, showing the lowest IC_{50} value (63.24 $\mu\text{g mL}^{-1}$ dry extract), followed by Synechococcales cyanobacterium LEGE 181150 (112.18 $\mu\text{g mL}^{-1}$ dry extract), with no significant difference, and Oscillatoriales cyanobacterium LEGE 181159 (332.40 $\mu\text{g mL}^{-1}$ dry extract) ($p < 0.05$).

Table 4. Inhibitory concentration (IC) values ($\mu\text{g mL}^{-1}$) of cyanobacteria aqueous and acetic extracts against superoxide anion radical ($\text{O}_2^{\bullet-}$)^{1,2}.

Strains	$\text{O}_2^{\bullet-}$ ($\mu\text{g mL}^{-1}$)			
	Aqueous Extracts		Acetic Extracts	
	IC ₂₅	IC ₅₀	IC ₂₅	IC ₅₀
<i>Cyanobium</i> sp. LEGE 07175	248.31 ^c ± 1.97	816.22 ^c ± 11.48	921.46 ^b ± 77.09	1549.10 ^b ± 65.28
Synechococcales cyanobacterium LEGE 181150	25.48 ^{a,b} ± 9.46	112.18 ^a ± 12.46	538.01 ^a ± 67.00	943.45 ^a ± 45.48
<i>Leptothoe</i> sp. LEGE 181156	41.10 ^{a,b} ± 5.24	nd	647.24 ^a ± 8.95	1097.05 ^a ± 2.41
Synechococcales cyanobacterium LEGE 181157	25.48 ^{a,b} ± 6.81	63.24 ^a ± 24.67	467.26 ^a ± 41.66	847.65 ^a ± 30.70
Synechococcales cyanobacterium LEGE 181158	51.22 ^b ± 24.20	404.59 ^b ± 14.30	1032.15 ^b ± 139.10	1463.69 ^b ± 271.87
Oscillatoriales cyanobacterium LEGE 181159	12.27 ^a ± 0.70	332.40 ^b ± 77.41	—	—

¹ Mean ± SD of at least three independent experiments, performed in duplicate. ² Different superscript letters in the same column correspond to statistical differences at $p < 0.05$ (ANOVA; Tukey's HSD). nd: not determined. —: not evaluated.




Figure 2. Superoxide anion radical ($\text{O}_2^{\bullet-}$) scavenging activity of cyanobacteria acetic (a) and aqueous (b) extracts. Values are expressed as the mean ± SD of at least three independent experiments, performed in duplicate.

Concerning acetic extracts, the lowest IC₅₀ was presented by *Synechococcales* cyanobacterium LEGE 181157, followed by *Synechococcales* cyanobacterium LEGE 181150 and *Leptothoe* sp. LEGE 181156 (847.65; 943.45; and 1097.05 $\mu\text{g mL}^{-1}$ dry extract) with no significant differences among the species (Table 4).

2.4. Enzymes Inhibition

2.4.1. Hyaluronidase (Hase) Inhibition

For both extracts studied herein, the strains of the order *Synechococcales* presented higher potential for Hase inhibition. *Synechococcales* cyanobacterium LEGE 181150 showed the lowest IC₅₀ value for aqueous and acetic extracts (483.86 and 726.29 $\mu\text{g mL}^{-1}$ dry extract, respectively), followed by *Synechococcales* cyanobacterium LEGE 181158 (624.51 and 738.88 $\mu\text{g mL}^{-1}$ dry extract, respectively) and *Synechococcales* cyanobacterium LEGE 181157 (645.06 and 859.83 $\mu\text{g mL}^{-1}$ dry extract, respectively), with no significant differences between them. *Leptothoe* sp. LEGE 181156 was the unique strain for which the acetic extracts did not reach IC₅₀ nor IC₂₅ values (Table 5).

Table 5. Inhibitory concentration (IC) values ($\mu\text{g mL}^{-1}$) of cyanobacteria aqueous and acetic extracts obtained for hyaluronidase, elastase, and tyrosinase ^{1,2}.

Strains	Hyaluronidase ($\mu\text{g mL}^{-1}$)				Elastase ($\mu\text{g mL}^{-1}$)		Tyrosinase ($\mu\text{g mL}^{-1}$)	
	Aqueous Extracts		Acetonic Extracts		Acetonic Extracts		Acetonic Extracts	
	IC ₂₅	IC ₅₀	IC ₂₅	IC ₅₀	IC ₂₅	IC ₅₀	IC ₂₅	IC ₅₀
<i>Cyanobium</i> sp. LEGE 07175	488.39 ^{a,b} ± 97.20	894.59 ^c ± 78.14	796.94 ^a ± 284.55	nd	nd	nd	795.57 ^b ± 198.79	nd
Synechococcales cyanobacterium LEGE 181150	257.05 ^{a,b} ± 1.39	483.86 ^a ± 86.71	595.11 ^a ± 4.77	726.29 ^a ± 16.56	nd	nd	nd	nd
<i>Leptothoe</i> sp. LEGE 181156	800.00 ^c ± 152.73	nd	nd	nd	378.26 ^b ± 24.50	nd	133.47 ^a ± 35.40	465.92 ^a ± 37.59
Synechococcales cyanobacterium LEGE 181157	188.89 ^a ± 86.14	645.06 ^{a,b} ± 118.73	613.47 ^a ± 82.241	859.83 ^a ± 122.08	51.43 ^a ± 20.02	380.50 ± 19.72	381.33 ^a ± 14.378	849.48 ^b ± 75.815
Synechococcales cyanobacterium LEGE 181158	397.72 ^{a,b} ± 70.00	624.51 ^{a,b} ± 19.84	442.63 ^a ± 36.54	738.88 ^a ± 83.79	233.47 ^b ± 89.26	nd	nd	nd
Oscillatoriales cyanobacterium LEGE 181159	518.69 ^{b,c} ± 15.68	801.30 ^{b,c} ± 116.04	–	–	–	–	–	–

¹ Mean ± SD of at least three independent experiments, performed in duplicate. ² Different superscript letters in the same column correspond to statistical differences at $p < 0.05$ (ANOVA; Tukey's HSD). nd: not determined. – not evaluated.

2.4.2. Elastase Inhibition

For elastase, only the acetonic extracts showed inhibitory potential, with *Synechococcales* cyanobacterium LEGE 181157 being the only strain that reached IC₅₀, with a value of 380.50 $\mu\text{g mL}^{-1}$ dry extract. Regarding IC₂₅, this strain reached an interesting value (51.43 $\mu\text{g mL}^{-1}$ dry extract) ($p < 0.05$), followed by *Synechococcales* cyanobacterium LEGE 181158 and *Leptothoe* sp. LEGE 181156 (233.47 and 378.26 $\mu\text{g mL}^{-1}$ dry extract) (Table 5).

2.4.3. Tyrosinase Inhibition

As for elastase, only acetonic extracts were able to promote tyrosinase inhibition (Table 5). *Leptothoe* sp. LEGE 181156 reached IC₅₀, followed by *Synechococcales* cyanobacterium LEGE 181157 (465.92 and 849.48 mg mL^{-1} , respectively) ($p < 0.05$). Additionally, both were also able to reach IC₂₅ with values of 133.47 and 381.33 mg mL^{-1} , respectively) with no significant differences.

2.4.4. Collagenase Inhibition

Contrary to the previous enzymes, the IC values were not able to be determined for collagenase. Due to the scarce amount of enzyme available, we chose to carry out the assay in kinetic mode, using the result of the calculation by slope of the linear equation line. In this sense, the inhibitory activity of the extracts was tested at two different concentrations of 0.75 and 1 mg mL^{-1} .

Acetonic extracts were more effective than aqueous ones towards collagenase inhibition (Table 6). *Leptothoe* sp. LEGE 181156 presented 36.61% of collagenase inhibition for the tested concentration of 0.75 mg mL^{-1} , while *Cyanobium* sp. LEGE 07175 reached 32.88% for the same concentration. Although *Synechococcales* cyanobacterium LEGE 181150 showed 9.81% of collagenase inhibition at 0.75 mg mL^{-1} , this strain presented 44.40% when tested at 1.0 mg mL^{-1} .

Regarding aqueous extracts, *Synechococcales* cyanobacterium LEGE 181158 was the most effective strain against collagenase, presenting 25.97% of inhibition at 1.0 mg mL^{-1} .

Table 6. Percentage (%) of inhibition of collagenase displayed by cyanobacteria aqueous and acetic extracts ^{1,2}.

Strains	% Collagenase Inhibition			
	Aqueous Extracts		Acetic Extracts	
	0.75 mg mL ⁻¹	1 mg mL ⁻¹	0.75 mg mL ⁻¹	1 mg mL ⁻¹
<i>Cyanobium</i> sp. LEGE 07175	nd	8.91 ^b ± 2.1	32.88 ^a ± 1.70	nd
Synechococcales cyanobacterium LEGE 181150	nd	nd	9.81 ^b ± 2.28	44.40 ^a ± 3.87
<i>Leptothoe</i> sp. LEGE 181156	nd	nd	36.61 ^a ± 2.72	36.74 ^{a,b} ± 7.82
Synechococcales cyanobacterium LEGE 181157	nd	nd	nd	nd
Synechococcales cyanobacterium LEGE 181158	nd	25.97 ^a ± 6.22	7.30 ^b ± 1.58	24.82 ^b ± 4.01
Oscillatoriales cyanobacterium LEGE 181159	nd	nd	–	–

¹ Mean ± SD of at least three independent experiments, performed in duplicate. ² Different superscript letters in the same column correspond to statistical differences at $p < 0.05$ (unpaired t-test; ANOVA; Tukey's HSD). nd: not determined. – not evaluated.

3. Discussion

The stratified squamous epithelium that constitutes the epidermis is mainly composed of keratinocytes and corneocytes and is in a constant process of cell renewal. In the dermis, fibroblasts are essential cells involved in the production and maintenance of the collagen and elastic fibers that ensure resistance and the elasticity of the skin. In addition, dermis is richly vascularized, being responsible for the nutrition and oxygenation of the epidermis [1]. The absence of cytotoxicity to main skin cells, both from the epidermis and dermis, is mandatory in a skin formulation. In this work, although seven strains were initially selected, the strain *Leptothoe* sp. LEGE 181155 was excluded due to the cytotoxic effects of both extracts on keratinocytes, fibroblasts, and endothelial cells. Additionally, due to cytotoxic effects in all cell lines tested, the acetone extract of Oscillatoriales cyanobacterium LEGE 181159 was also excluded. Thus, only nontoxic extracts were further analyzed.

3.1. Phytochemical Analysis

3.1.1. Carotenoids and Chlorophylls

Among the studied strains, the acetic extract of the strain *Leptothoe* sp. LEGE 181156 was the richest in carotenoids (150.15 µg mg⁻¹ dry extract) ($p < 0.05$). The highest concentration of carotenoids in aqueous extracts was obtained with strain *Cyanobium* sp. LEGE 07175 with a maximum of 3.00 µg mL⁻¹ dry extract. As expected, the results obtained indicated that acetone extraction is more efficient than the aqueous on both carotenoids and chlorophylls (Table 1), which can be explained by the lower polar nature of these compounds, and therefore a higher affinity for lower polar solvents such as acetone. Additionally, as extractions were carried out in sequential mode, in which the acetic extraction was performed first, most of these pigments were extracted with acetone.

Considering carotenoids identification, the xanthophyll zeaxanthin was found in all acetic extracts of all species studied, ranging from 24.38 to 53.08 µg mg⁻¹ dry extract, with the highest value reported in *Cyanobium* sp. LEGE 07175, followed by *Leptothoe* sp. LEGE 181156 (53.08 and 39.85 µg mg⁻¹ dry extract, respectively) ($p < 0.05$). In a previous study by Morone and co-workers [18], a 70% ethanolic extract by *Cyanobium* sp. LEGE 07175 exhibited 16.31 µg g⁻¹ of dry biomass, with the best value being presented by *Synechocystis salina* LEGE 06099, with 49.82 µg g⁻¹ of dry biomass. In this work, in terms of dry biomass, acetic extracts of *Cyanobium* sp. LEGE 07175 presented a value of 799 µg g⁻¹ and *Leptothoe* sp. LEGE 181156 a value of 498 µg g⁻¹. Zeaxanthin's properties include photoprotection in cyanobacteria and reduction of oxidative damage in humans' eyes [19,20].

Myxoxanthophyll was detected in all cyanobacteria acetic extracts and most aqueous extracts, except for *Cyanobium* sp. LEGE 07175. In contrast, echinenone was only detected in *Cyanobium* sp. LEGE 07175 with a concentration of 1.17 µg mg⁻¹ dry extract. This same carotenoid was detected in a 70% ethanol extract of *Cyanobium gracile* LEGE 12431

(7.31 $\mu\text{g mg}^{-1}$ dry extract) [21]. Echinenone was also found in other genera of cyanobacteria [18,21]. According to previous studies, this carotenoid is important in cyanobacteria photoprotection [19].

Beta-carotene is a terpene of relevant importance due to its important role as an antioxidant. This carotenoid was found in all strains in both acetone and aqueous extracts. For acetonic extracts, the concentrations ranged from 19.58 to 47.89 $\mu\text{g mg}^{-1}$ dry extract. There was a significant difference among the strain *Leptothoe* sp. LEGE 181156 and the Synechococcales, strains LEGE 181158, LEGE 181157, and LEGE 181150, which did not differ significantly from each other. These results may corroborate a study that analyzed the patterns of carotenoid synthesis in different genera and concluded that strains of the same genus present the same patterns of pigments production, namely concerning β -carotene [22]. Previous studies also reported β -carotene in different genera of cyanobacteria, namely, *Synechocystis*, *Nodosilinea*, *Phormidium*, *Leptolyngbya*-like, *Cuspidothrix*, *Anabaena*, *Nostoc*, *Aphanothece*, and *Gloeotheca* [18,21,23–25], which validates the broad spectrum of cyanobacterial genera that produce this compound, and thus its importance in industries such as cosmetics.

Morone et al., 2020 [18], showed that a 70% ethanol extract of *Cyanobium* sp. LEGE 07175 presented 8.06 μg of β -carotene per g of dry biomass. Accounting the extraction yield, in the present work, the acetonic extract of *Cyanobium* sp. LEGE 07175, which exhibited 19.58 $\mu\text{g mg}^{-1}$ dry extract, presented a value of 295 $\mu\text{g g}^{-1}$ in terms of dry biomass. Another example is *Leptothoe* sp. LEGE 181156, which exhibited the highest value for β -carotene (47.89 $\mu\text{g mg}^{-1}$ dry extract) and, in terms of dry biomass, showed a value of 599 $\mu\text{g g}^{-1}$, for acetonic extract.

As β -carotene, γ -carotene was also detected in the acetonic extract of all species studied, although below the limit of quantification in *Cyanobium* sp. LEGE 07175. In addition to the antioxidant potential, β -carotene and γ -carotene are precursors of retinol, which works as a storage form of vitamin A and can be transformed into other modes of activation, such as retinoic acid and retinal [26]. In skin, retinoids are widely used in the treatment of diseases such as cancer, psoriasis, acne, ichthyosis, and even wrinkles due to their effect on cell differentiation, proliferation, and apoptosis [27]. Therefore, β -carotene and γ -carotene are two interesting carotenoids to be explored in cosmetics [28].

In this study, we found that zeaxanthin, β -carotene, and γ -carotene were the main carotenoids detected in most of the samples analyzed. A relevant aspect in the production of carotenoids in cyanobacteria is that by changing the culture conditions, it is possible to increase the production of these compounds. A previous study involving the optimization of culture conditions for cyanobacteria revealed that high light radiation may increase the production of carotenoids, such as β -carotene, echinenone, and myxoxanthophyll, and may exhibit strong protection under photo-oxidative conditions [29]. The results from the present study indicated that, despite the low yield achieved with the acetonic extraction (around 1.5 to 2% per gram of dry biomass (Table 7), the amount of carotenoids extracted with acetone may be profitable when compared to water, which presented percentages at least 20 times higher.

Table 7. Yield (% *w/w*) obtained from sequential acetonic and aqueous extractions ¹.

Strains	Solvents	
	Acetone	Water
<i>Cyanobium</i> sp. LEGE 07175	1.2 ± 0.3	17.2 ± 1.4
Synechococcales cyanobacterium LEGE 181150	1.4 ± 0.4	39.0 ± 5.2
<i>Leptothoe</i> sp. LEGE 181156	1.9 ± 0.6	26.6 ± 5.3
Synechococcales cyanobacterium LEGE 181157	1.6 ± 0.6	31.7 ± 8.4
Synechococcales cyanobacterium LEGE 181158	1.5 ± 0.3	23.4 ± 5.6
Oscillatoriales cyanobacterium LEGE 181159	–	38.9 ± 1.0

¹ Values are expressed as Mean ± SD of at least four extractions. – not evaluated.

Amongst the pigments analyzed herein, the total chlorophylls content, including chlorophyll-*a* and its derivatives, was lower than total carotenoids, except for aqueous extracts. Unlike the results of our study, Lopes et al., 2020 [21], showed that chlorophylls were dominant over carotenoids in both, ethanol and acetic extractions; however, the strains evaluated by the author were of terrestrial origin, which may explain the differences. Chlorophylls, as well as carotenoids, act as antioxidants and are widely used in the food, cosmetic, and pharmaceutical industries [30].

3.1.2. Total Phenolic Content (TPC)

Despite the limitations inherent to colorimetric assays, the determination of TPC by the Folin–Ciocalteu method is widely used to determine total phenols in plant extracts, as well as to infer about their antioxidant potential. Overall, acetic extracts had higher phenolic content than aqueous ones. Although phenolic compounds are soluble in solvents with higher polarity, the fact that the extraction was carried out sequentially with acetone and water led to the extraction of most of the compounds with acetone. The highest TPC was found in the acetone extract of *Leptothoe* sp. LEGE 181156, with $27.13 \mu\text{g GAE mg}^{-1} \text{ dry extract}$ with no significant differences when compared to Synechococcales cyanobacterium LEGE 181158 and *Cyanobium* sp. LEGE 07175 (24.71 and $23.11 \mu\text{g GAE mg}^{-1} \text{ dry extract}$, respectively). In a previous study by Morone et al. [18], using a 70% ethanol extract, *Cyanobium* sp. LEGE 07175 had a value of $1.09 \text{ mg GAE g}^{-1} \text{ dry biomass}$, which was higher than the result obtained herein for the same strain ($0.29 \text{ mg GAE g}^{-1}$, converted according to extraction yield). Another study with the same genus [31], revealed values around $82 \mu\text{g GAE mg}^{-1} \text{ dry extract}$ for acetic extracts of another *Cyanobium* sp. Favas et al. [14], presented a value of $17.59 \mu\text{g GAE mg}^{-1} \text{ dry extract}$ for acetic extract of *Leptolyngbya* cf. *ectocarpus* LEGE 11479.

Regarding aqueous extracts, although the values are lower, between 14.49 to $15.67 \mu\text{g GAE mg}^{-1} \text{ dry extract}$, when converted to dry biomass, the results turn out to be promising, considering the excellent yield that aqueous extraction may provide; for example, this was true for Synechococcales cyanobacterium LEGE 181150 and Synechococcales cyanobacterium LEGE 181157, which were the strains that presented the best results (6.48 and $4.33 \text{ mg GAE g}^{-1} \text{ dry biomass}$, respectively). However, regarding acetic extracts, the results were much lower for the same strains in terms of dry biomass (0.39 and $0.30 \text{ mg GAE g}^{-1} \text{ dry biomass}$, respectively). Phenolic compounds are known for their beneficial effect against humans, including skin diseases, namely due to their anti-inflammatory and antioxidant potential [32]. In the skin, phenolic compounds were found to be active as antioxidants and anti-inflammatories [33] and are effective at inhibiting the enzyme tyrosinase, and thus at treating aesthetic problems such as hyperpigmentation [34]. Regarding marine cyanobacteria, as far as we are aware, there are no other previous reports on the TPC, corroborating the importance of further exploitation of cyanobacteria extracts concerning antioxidant activity.

3.1.3. Proteins

Proteins play a key role in the structure and integrity of the skin. The main benefit of using proteins in cosmeceuticals is related to improving skin hydration, as they act as water-binding molecules. In addition to being responsible for forming collagen, which is a protein that gives firmness to skin and prevents the formation and exacerbation of wrinkles and expression lines, proteins are the main basis for the formation of new tissues, also being implicated in improving wound healing. Furthermore, proteins present UV protection capacity, antimicrobial, and antioxidant properties. Therefore, there is a growing search for products with antiaging action based on bioactive peptides [35]. In this study, the characterization of the total protein content of the extracts studied was carried out. It was confirmed that the aqueous extracts presented much higher values than the acetic ones, with emphasis on strains used in this work within the order Synechococcales, which showed promising results. Synechococcales cyanobacterium LEGE 181157 ranked first with $760.81 \mu\text{g (BSA) mg}^{-1} \text{ dry extract}$ ($p < 0.05$), followed by Synechococcales

cyanobacterium LEGE 181150 ($p < 0.05$) and Synechococcales cyanobacterium LEGE 181158 (695.25 and 579.99 $\mu\text{g (BSA) mg}^{-1}$ dry extract, respectively). Converting to extraction yield, the values are 220, 331, and 100 mg (BSA) g^{-1} dry biomass, respectively. Therefore, in terms of biomass, Synechococcales cyanobacterium LEGE 181150 exhibited the highest value. Favas and coworkers, 2022 [14] also reported higher values for aqueous extracts than acetonetic ones, in which the freshwater strain *Cephlothrix lacustris* LEGE 15493 presented 521.18 $\mu\text{g (BSA) mg}^{-1}$ dry extract.

3.1.4. Phycobiliproteins (PBP)

Phycobiliproteins are water-soluble proteins, which are deep-colored, formed by a complex between proteins and covalently bound phycobilins. Phycocyanin (PC), allophycocyanin (APC), and phycoerythrin (PE) are the three main PBP groups, absorbing light within specific regions of the spectrum, according to their structure and types of bilins [9]. In recent years, the demand for microorganisms to synthesize natural colorants for cosmetics has increased in order to replace synthetic ones used in lipsticks, eyeliners, and blushes [2]. The PBP PC, of blue color; APC, blue-green; and PE and PEC, both of red/pink color, have been already used as a natural coloring in foods, nutritional supplements, cosmetics, and as fluorescent markers in immunoassays [36].

As water-soluble compounds, PBPs were only analyzed in aqueous extracts. As visually confirmed through its intense pink color, the aqueous extract of the strain Synechococcales cyanobacterium LEGE 181157 was the richest in PE (275.04 $\mu\text{g mg}^{-1}$ dry extract) followed by the strain *Leptothoe* sp. LEGE 181156 (159.05 $\mu\text{g mg}^{-1}$ dry extract) ($p < 0.05$). Validated by the strong blue color, Synechococcales cyanobacterium LEGE 181150 aqueous extract was the richest in PC (222.76 $\mu\text{g mg}^{-1}$ dry extract) ($p < 0.05$). Regarding APC, Oscillatoriales cyanobacterium LEGE 181159 presented the highest value, followed by Synechococcales cyanobacterium LEGE 181150 (65.90 and 57.29 $\mu\text{g mg}^{-1}$ dry extract, respectively). Only *Cyanobium* sp. LEGE 07175, with a light green extract, showed significantly lower values. In terms of dry biomass, Synechococcales cyanobacterium LEGE 181157 exhibited 113.18 mg g^{-1} dry biomass for PE and Synechococcales cyanobacterium LEGE 181150 83.53 mg g^{-1} dry biomass for PC. These values are in accordance with literature data, namely for PE concentration in the thermotolerant cyanobacterium *Leptolyngbya* sp. KC45 (100 mg g^{-1}) [37]. For PC, the value described is within the concentrations obtained with different strains of *Spirulina*, as exposed in a recent review by Jaeschke and co-workers [38] in which different methods of PC extraction from *Spirulina* strains are compared. Concerning total PBPs by dry biomass, the results obtained are also in accordance with reported data for the strain Synechococcales cyanobacterium LEGE 181157 (179 mg g^{-1} dry biomass) compared with in the strain *Anabaena* sp. NCCU-9 (91 mg g^{-1} dry biomass) [39] and *Cyanobium* sp. (200 mg g^{-1} dry biomass) [31]. The results obtained confirm the use of cyanobacteria to obtain natural dyes and expand the range of genera with this potential. Along with the coloring role, PBPs are described as potent antioxidant and anti-inflammatory agents, properties that are beneficial in cosmetic formulations [40].

3.1.5. Antioxidant Potential

In this work, the relevant physiological free radical $\text{O}_2^{\bullet-}$, one of the major ROS that provoke oxidative damage in the human body, was used to determine the antioxidant potential of cyanobacteria extracts. In terms of radical scavenging, the aqueous extract of the strain Synechococcales cyanobacterium LEGE 181157 was the most effective, presenting the best IC_{50} value of 63.24 $\mu\text{g mL}^{-1}$ dry extract, followed by Synechococcales cyanobacterium LEGE 181150 (112.18 $\mu\text{g mL}^{-1}$ dry extract) ($p > 0.05$). Synechococcales cyanobacterium LEGE 181157 was also the strain that presented the highest value for total proteins (760.81 $\mu\text{g BSA mg}^{-1}$ dry extract) ($p < 0.05$), followed by Synechococcales cyanobacterium LEGE 181150, with 695.25 $\mu\text{g BSA mg}^{-1}$ dry extract. Moreover, Synechococcales cyanobacterium LEGE 181157 showed the highest value for PE (275.04 $\mu\text{g mg}^{-1}$ dry extract) ($p < 0.05$),

and Synechococcales cyanobacterium LEGE 181150 presented the highest content for PC ($222.76 \mu\text{g mg}^{-1}_{\text{dry extract}}$) ($p < 0.05$). According to the statistical analyses, a significant negative correlation was found between the total proteins and the IC values (-0.982 , $p < 0.01$ for IC_{50} and -0.739 , $p < 0.01$ for IC_{25}). In addition, a negative correlation was also observed between PE and IC values, and the same occurred for the TPC. Furthermore, a significant negative correlation was found between ICs and PC (-0.680 , $p < 0.05$ and -0.749 , $p < 0.01$, for IC_{50} and IC_{25} , respectively). Although *Leptothoe* sp. LEGE 181156 was the only strain that did not reach the IC_{50} for aqueous extract, this strain managed to obtain more than 40% of enzymatic inhibition at the concentration $104.00 \mu\text{g mL}^{-1}_{\text{dry extract}}$, and its IC_{25} reached $41.10 \mu\text{g mL}^{-1}_{\text{extract dry}}$ (Figure 2). This antioxidant activity observed in the aqueous extracts may be attributed to their richness in proteins and PBP.

Regarding the acetonetic extracts, strains belonging to Synechococcales cyanobacterium order, followed by *Leptothoe* sp. LEGE 181156, showed the best results for free-radical sequestration ($p > 0.05$). However, the values were much higher than those displayed by the aqueous extract. These acetonetic extracts are richer in carotenoids than aqueous extracts (Table 1). According to the statistical analyses, although they lacked statistical significance, negative correlations were noted with total carotenoids content, lutein, β -carotene, and γ -carotene. Favas et al. [14], using both extractions, also reported a higher effectiveness for the aqueous extracts when compared to the acetonetic ones. In their work, aqueous extracts of *Cephalothrix lacustris* LEGE 15493 showed an IC_{50} of $65.50 \mu\text{g mL}^{-1}_{\text{dry extract}}$, while the IC_{50} of its acetonetic extract was not detected. Morone et al. [18], who studied ethanol extracts of other species, reported the lowest IC_{50} value for *Phormidium* sp. LEGE 05292 ($822.70 \mu\text{g mL}^{-1}$). Amaro and co-workers [23] revealed that *Scenedesmus obliquus* (M2-1) showed an IC_{50} of $826 \mu\text{g mL}^{-1}$ for acetonetic extract. According to the studies published so far, aqueous extracts have shown great potential in terms of their ability to scavenge $\text{O}_2^{\bullet-}$. In this way, we may consider the interest of the aqueous extracts of the strains explored in our work for cosmeceutical applications.

3.1.6. Inhibition of Metalloproteinases (MMPs)

The dermis ECM is extremely important for the maintenance of skin structure. When collagen, elastin, and HA are degraded or deficient, the effects are expressed in the visual appearance of the skin. Under oxidative stress, an inflammatory response with activation of MMPs such as collagenase, elastase, and HAase is triggered, contributing to the degradation of these ECM vital components, and promoting wrinkles formation [1,41]. Tyrosinase is another enzyme that is also activated mainly by external factors, such as exposure to UV rays. This enzyme acts in photoprotection as a catalyst in the synthesis of melanin, but when overactivated, it causes pigmentation lesions, such as vitiligo, or hyperpigmentation marks, such as melasma and age spots [42]. Triggered by the importance of MMPs inhibitors in skin, the ability of the aqueous and acetonetic extracts to inhibit these key enzymes was explored.

Regarding HAase, the lowest results for IC_{50} were presented by the aqueous extracts of the species belonging to the order Synechococcales, in which the strain Synechococcales cyanobacterium LEGE 181150 presented the best value ($483.86 \mu\text{g mL}^{-1}_{\text{dry extract}}$). For acetonetic extracts, the results were around $750 \mu\text{g mL}^{-1}_{\text{dry extract}}$. Regarding statistical analysis, there is a strong negative correlation between total proteins and the IC_{50} for both aqueous and acetonetic extracts (-0.593 , $p < 0.05$ and -1.000 , $p < 0.01$, respectively) and a negative correlation between TPC and IC_{50} , also for both extracts, suggesting that both proteins and phenols may be implicated in the inhibition of this enzyme. Synechococcales cyanobacterium LEGE 181150 was the strain exhibiting one of the highest total proteins for both extractions, which corroborates the deductions above. In this study, the value of total proteins for aqueous extracts of *Cyanobium* sp. was not as high as that presented by the strains belonging to Synechococcales order, which is also in line with the statistical correlations found. Our results highlight the importance of aqueous extracts as potential antiaging ingredients, possibly due to their protein content. Although the aqueous extract

of *Cyanobium* sp. LEGE 07175 displayed an IC_{50} of $894.59 \mu\text{g mL}^{-1}$, in a previous study realized by us [18], a 70% ethanol extract of this strain, *Cyanobium* sp. LEGE 07175, showed stronger inhibitory activity ($IC_{50} = 208.36 \mu\text{g mL}^{-1}$). This may be due to a synergistic effect between the compounds extracted by ethanol 70%, since this solvent has more affinity to a mixture of polar and less polar compounds, contrary to water, from which compounds of more polar nature are mainly extracted. Furthermore, Pagels et al. [43], showed considerable IC_{50} for aqueous and acetonic extracts, using another strain of the genus *Cyanobium* (67.25 and $108.74 \mu\text{g mL}^{-1}$, respectively). Favas et al. [14], presented the lowest value for aqueous extract of *Leptolyngbya* cf. *ectocarpus* LEGE 11479 ($863 \mu\text{g mL}^{-1}$ dry extract) and only IC_{25} for acetonic extracts. A previous work focusing polysaccharides from cyanobacteria [13] reported that *Nostochopsis lobatus* MAC0804NAN had a notable IC_{50} of $7.18 \mu\text{g mL}^{-1}$ for HAase inhibition. These results caught our interest since, considering the polar nature of polysaccharides, these compounds may also be present in the aqueous extracts and, together with proteins, contribute to the inhibition of the enzyme.

Inhibition of elastase was only achieved with acetonic extracts, Synechococcales cyanobacterium LEGE 181157 being the unique one that reached IC_{50} value ($380.50 \mu\text{g mL}^{-1}$ dry extract). In addition, its IC_{25} presented an interesting value of $51.43 \mu\text{g mL}^{-1}$ dry extract. On the other hand, this strain was not able to inhibit collagenase. For acetonic extracts, collagenase inhibition was achieved at $750 \mu\text{g mL}^{-1}$ for *Leptothoe* sp. LEGE 181156 and *Cyanobium* sp. LEGE 07175 (36.61 and 32.88%, respectively), and at $1000 \mu\text{g mL}^{-1}$ for Synechococcales cyanobacterium LEGE 181150, with 44.40%. Favas et al. [14], reported that acetonic extracts of *Leptolyngbya* cf. *ectocarpus* LEGE 11479 reached IC_{50} ($391 \mu\text{g mL}^{-1}$ dry extract) for elastase inhibition, and Pagels et al. [43] showed collagenase inhibition for aqueous extract of the strain *Cyanobium* sp. LEGE 06113 ($IC_{50} = 582.82 \mu\text{g mL}^{-1}$) but with no results below $1000 \mu\text{g mL}^{-1}$ for elastase and tyrosinase inhibition. Regarding aqueous extracts, Synechococcales cyanobacterium LEGE 181158 presented 25.97% of inhibition at $1000 \mu\text{g mL}^{-1}$. As presented above, this strain also exhibited inhibition of hyaluronidase.

Finally, IC_{50} for tyrosinase was displayed for the acetonic extracts for two strains. *Leptothoe* sp. LEGE 181156 was the only strain that exhibited IC results for both tyrosinase ($IC_{50} = 465.92 \mu\text{g mL}^{-1}$ and $IC_{25} = 133.47 \mu\text{g mL}^{-1}$) and elastase ($IC_{25} = 378.26 \mu\text{g mL}^{-1}$). In addition, this strain also showed collagenase inhibition of 36.61% at a concentration of $750 \mu\text{g mL}^{-1}$ dry extract. With respect to Synechococcales cyanobacterium LEGE 181157, ICs values for hyaluronidase, elastase and tyrosinase inhibition were found, the IC_{25} being more attractive (613.47 , 51.43 , and $381.33 \mu\text{g mL}^{-1}$ dry extract, respectively). A more detailed analysis showed a significant negative correlation between TPC (-0.972 , $p < 0.05$), Total Proteins (-0.984 , $p < 0.05$) and Total Carotenoids (-0.974 , $p < 0.05$), and IC_{50} . Additionally, zeaxanthin, β -carotene (-0.956 , $p < 0.05$), and γ -carotene showed a negative correlation in relation to tyrosinase inhibition. For the same enzyme, Favas et al. [14], reported that only the acetonic extract of *Nodosilinea nodulosa* LEGE 06104 was able to reach IC_{50} ($989.26 \mu\text{g mL}^{-1}$ dry extract). Ethanol extracts studied by Morone et al. [18] did not show tyrosinase inhibition.

In a general way, extracts richer in proteins seem to be more effective for free radicals scavenging, being more attractive to counteract oxidative stress, while acetonic extracts, which are richer in carotenoids and phenols, seem more effective as enzyme inhibitors. Regarding phenolic compounds, it seems that they contribute to both biological activities evaluated.

4. Materials and Methods

4.1. Cyanobacteria Strains and Biomass Production

Seven cyanobacterial strains isolated from the Portuguese and Cape Verde marine ecosystems and maintained in the Blue Biotechnology and Ecotoxicology Culture Collection (LEGE-CC) at the Interdisciplinary Centre of Marine and Environmental Research (CIIMAR) [44] were randomly chosen for this study, with the aim of conducting a more

comprehensive screening of the most promising cyanobacteria strains. The strains panel included the picoplanktonic *Cyanobium* sp. LEGE 07175 from Portuguese marine ecosystem and the filamentous Synechococcales cyanobacterium LEGE 181150 [45], *Leptothoe* sp. LEGE 181155, *Leptothoe* sp. LEGE 181156, Synechococcales cyanobacterium LEGE 181157, Synechococcales cyanobacterium LEGE 181158, and Oscillatoriales cyanobacterium LEGE 181159, from Cape Verde marine ecosystems, which are in the process of publication. For biomass production, a scale-up culture scheme up to 4 L was set. The strains were grown in Z8 medium [46], supplemented with 10 µg/L vitamin B12 and 25 g/L of synthetic sea salts (Tropic marin, Berlin, Germany). Cultures were maintained at 25 °C, with a light intensity of 10–30 µmol photons m⁻² s⁻¹ and with a photoperiod of 14h light:10h dark. The fresh biomass was collected by filtration, frozen, freeze-dried, and stored at –20 °C until extracts preparation.

4.2. Extract Preparation

Acetone and aqueous extracts were sequentially prepared from each strain by following the procedure described by Favas et al. [14]. First, the acetonic extract was prepared using 2 g of dry biomass. The biomass was suspended in 80 mL of acetone and extracted for 10 min in an ultrasonic bath (Fisherbrand®-FB15053, Loughborough, UK). The supernatant was collected, and the biomass was reextracted three more times. After the acetonic extraction, the resulting pellet was left to dry in the fume hood until the remaining solvent was completely evaporated. The pellet was further extracted with 70 mL of distilled water, following the same procedure. Cell debris was removed by centrifugation (5000 Gs, 5 min, 4 °C) (Thermo Scientific™ HERAUS Megafuge™ 16R, Waltham, MA, USA). Supernatants from each extraction were evaporated under reduced pressure (acetone) (BUCHI R-210 Rotary Evaporator, Cambridge, MA, USA) or frozen and lyophilized (water). The dry extracts were kept at –20 °C until further chemical and biological analysis. The extractions yield is displayed below (Table 7).

4.3. Phytochemical Analysis

4.3.1. Determination of Pigments Profile by HPLC-PDA

Dried cyanobacteria extracts were dissolved in HPLC-grade methanol (acetonic extract) or Milli-Q water (aqueous extracts) to a final concentration of 5 and 10 mg mL⁻¹, respectively, and filtered through a 0.2 µm pore membrane. Carotenoids analysis was performed following the method previously described [47], with slight modifications. A Waters Alliance 2695 high-performance liquid chromatography (HPLC) with photodiode array (PDA) detector (USA) was employed to resolve, detect, and identify the compounds of interest. The stationary phase was a YMC Carotenoid C30 (250 × 4.6 mm; 5 µm) column, kept at constant temperature (25 °C) with a column heater (Waters Corporation, Milford, CT, USA). The mobile phase consisted of 2 solvents: methanol (A) and tert butyl methyl ether (B) (VWR Prolabo), starting with 95% A and installing a gradient to obtain 10% B at 5 min, 18% B at 20 min, 30% B at 28 min, 50% B from 31 to 37 min, 80% B from 38 to 47 min, and 5% B from 48 to 50 min. The flow rate was 0.90 mLmin⁻¹ and the injection volume was 5 µL. Data were processed using Empower chromatography software (Waters, USA). Spectra data from all peaks were collected in the range 250 to 750 nm.

Compounds were identified by comparing their retention times and UV-Vis spectra with those of authentic standards. Carotenoids quantification was achieved by measuring the absorbance recorded in the chromatograms relative to external standards at 450 nm.

Zeaxanthin, lutein, echinenone, myxoxanthophyll, phaeophytin-*a*, β-carotene, γ-carotene, and chlorophyll-*a* (Extrasynthese, Genay, France; Sigma-Aldrich, St. Louise, MO, USA; DHI, Horsholm, Denmark) were quantified with the authentic standards; unidentified carotenoids were quantified as zeaxanthin, the most representative xanthophyll, and chlorophyll derivatives and phaeophytin-*a* as chlorophyll-*a*, the major cyanobacteria chlorophyll. Calibration curves were performed with five different concentrations of standards, selected

as representative of the range of compounds concentrations in the samples. The calibration plots and r^2 values for the analyzed carotenoids and chlorophyll-*a* are shown in Table 8.

Table 8. Calibration curves of authentic standards used for quantification of different carotenoids and chlorophylls.

Standards	Calibration Curve	r^2
Lutein	$y = 31,188,975x + 81,368$	0.9987
Chlorophyll- <i>a</i>	$y = 5,647,422x + 14,838$	0.9989
Zeaxanthin	$y = 40,108,171x + 97,810$	0.9994
Cantaxanthin	$y = 39,997,059x + 197,655$	0.9992
Myxoxantophyll	$y = 30,518,380x + 5976$	0.9993
Echinenone	$y = 74,770,292x + 126,878$	0.9997
β -Carotene	$y = 31,852,521x + 16,127$	0.9999
γ -Carotene	$y = 452,252x + 17,574$	0.9913

4.3.2. Total Phenolic Content (TPC)

The TPC of the cyanobacterial extracts was determined using the colorimetric assay of Folin–Ciocalteu, according to Barroso et al. [18,48]. The acetonic extracts were solubilized in DMSO and the aqueous extracts in water. Briefly, a volume of 25 μL of each extract (10 mg mL^{-1}) was thoroughly mixed with 25 μL of Folin–Ciocalteu reagent (Sigma-Aldrich, St. Louis, MO, USA), 200 μL of Na_2CO_3 solution (75 g L^{-1}) and 500 μL of deionized water. After the incubation period (60 min at room temperature), the absorbance of the colored product was measured at 725 nm, using a Synergy HT Multi-detection microplate reader (Biotek, Bad Friedrichshall, Germany) operated by GEN5TM software. Standard calibration curves ($y = 2.097x + 0.01560$, $R^2 = 0.9989$, for aqueous extracts and $y = 2.204x + 0.01401$, $R^2 = 0.9982$, for acetonic extracts) were obtained with seven concentrations of gallic acid (GA) (0.025 to 0.5 mg mL^{-1}). TPC in each extract was expressed in μg of gallic acid equivalents (GAE) per mg of dry biomass. Three independent determinations were carried out in triplicate.

4.3.3. Total Proteins

Total proteins concentration was determined using the BSA Protein Assay kit (n° 23227, Thermo-Scientific, Waltham, MA, USA) by following the manufacturer's instructions and according to Favas et al. [14]. Aqueous extracts were prepared in water, while acetone extracts were prepared in DMSO. Briefly, in a 96-well plate, 25 μL of each extract (1 mg mL^{-1}) was mixed with 200 μL of working reagent. The absorbance was measured at 562 nm using a Synergy HT Multi-detection microplate reader (Biotek, Bad Friedrichshall, Germany) operated by GEN5TM software. Standard curves ($y = -126.87x^3 + 547.73x^2 + 483.85x - 10.017$; $R^2 = 0.999$ and $y = 162.87x^3 - 248.51x^2 + 932.13x - 11.715$; $R^2 = 0.999$) were obtained for each extract (aqueous and acetonic, respectively), using nine concentrations of albumin (BSA) (25 to 2000 $\mu\text{g mL}^{-1}$) to quantify the proteins. Three independent experiments were carried out in triplicate. The total proteins were expressed as μg of bovine serum albumin (BSA) equivalents per mg of dry extracts.

4.3.4. Phycobiliproteins

The pigments present in aqueous extracts were determined spectrophotometrically. Aqueous extracts were resuspended in water. PBP's were determined by measuring the ab-

sorbances at different wavelengths (562, 615 and 645 nm), in a cell with 1 cm of optical path. The corresponding formulas were applied, as previously described by Pagels et al. [31]:

$$\text{Phycocyanin}(PC) = \frac{A_{615nm} - 0.474 \times A_{652nm}}{5.34}$$

$$\text{Allophycocyanin}(APC) = \frac{A_{652nm} - 0.208 \times A_{615nm}}{5.09}$$

$$\text{Phycocerythrin}(PE) = \frac{A_{562nm} - 2.41 \times PC - 0.849 \times APC}{9.62}$$

Aqueous extracts were resuspended to a final concentration of 0.5 mg mL⁻¹. The experiment was carried out in triplicate, and the results were expressed in µg of the respective phycobiliprotein per mg of dry extract.

4.4. Cell Assays

4.4.1. Cell Culture

To assess the safety of the cyanobacteria acetonic and aqueous extracts, a preliminary in vitro cytotoxicity assay was performed using three different cell lines, the human keratinocytes HaCAT (ATCC), the mice fibroblasts 3T3L1 (ATCC) and the human endothelial cells hCMEC/D3 (provided by Dr. PO Couraud (INSERM, Paris, France)). Cell culture was performed as already described by us in Morone et al. and Favas et al. Briefly, cells were cultured in DMEM Glutamax medium (Dulbecco's Modified Eagle Medium DMEM GlutaMAX™—Gibco, Glasgow, UK), supplemented with 10% (*v/v*) of fetal bovine serum (Biochrom, Berlin, Germany), 0.1% of amphotericin B (GE Healthcare, Little Chalfont, UK) and 1% of Pen-Strep (penicillin-streptomycin, 100 IU mL⁻¹ and 10 mg mL⁻¹, respectively) (Gibco, Berlin, Germany). Cells maintenance and assays were performed in a humidified atmosphere of 5% CO₂ at 37 °C. Between 80–90% cell confluence, adherent cells were washed with phosphate-buffered saline (PBS, Gibco), detached with a small amount of TrypLE Express enzyme (Gibco), passed for maintenance, and seeded for the planned assays.

4.4.2. Cytotoxicity—MTT Assay

Cytotoxicity assay was performed by measuring the reduction of 3-(4,5-dimethylthiazole-2-yl)-2,5-diphenyltetrazolium bromide (MTT), as previously described [18]. Keratinocytes, fibroblasts and endothelial cells were seeded in 96-well plates at a density of 2.5 × 10⁴ cells mL⁻¹; 3.3 × 10⁴ cells mL⁻¹ and 1.0 × 10⁵ cells mL⁻¹, respectively. After 24 h of adhesion, the culture medium was removed and cells were exposed for 24 and 48 h to fresh medium supplemented with 1% of extracts in five serial concentrations, from 12.5 to 200 µg mL⁻¹. Acetone extracts were prepared in dimethyl sulfoxide (DMSO, Gibco) and diluted with DMEM prior to cells exposure; the maximum DMSO concentration did not exceed 1%. DMSO at 1% and 20% were used as solvent and positive controls, respectively. Aqueous extracts were prepared in PBS and diluted with DMEM prior to cells exposition. PBS at 1% and DMSO at 20% were used as solvent and positive controls, respectively. After each incubation time, 20 µL of 1 mg mL⁻¹ MTT (Sigma-Aldrich), was added to each well and incubated at 37 °C for 3h. Following incubation, the media were removed, and the purple-colored formazan salts were dissolved in DMSO. The absorbance was read at 550 nm in a Synergy HT Multi-detection microplate reader (Biotek, Bad Friedrichshall, Germany) operated by GEN5™ software. The assay was run in quadruplicate and averaged. Cytotoxicity was expressed as a percentage of cell viability, considering 100% viability in the solvent control. For reproducibility, each assay was independently repeated three times.

4.5. Biological Activities

4.5.1. Superoxide Anion Radical ($O_2^{\bullet-}$) Scavenging

The superoxide anion radical ($O_2^{\bullet-}$) scavenging activity of the extracts was determined as described previously [49] with some modifications. The acetone extracts were resuspended in DMSO, while the aqueous extracts were resuspended in water. Six serial dilutions, from 10 mg mL^{-1} , were prepared for each extract and tested in order to evaluate the extracts' behavior and IC values. All reagents were dissolved in phosphate buffer ($19\text{ }\mu\text{M}$, pH 7.4). A volume of $50\text{ }\mu\text{L}$ of serial dilutions of the cyanobacteria extracts (0.104 to 1.667 mg mL^{-1} for acetonic extracts; and 0.013 to 1.667 mg mL^{-1} , for aqueous extracts) was mixed with $50\text{ }\mu\text{L}$ of $166\text{ }\mu\text{M}$ β -nicotinamide adenine dinucleotide reduced-form (NADH) solution and $150\text{ }\mu\text{L}$ of $43\text{ }\mu\text{M}$ nitro tetrazolium blue chloride (NBT), in a 96-well plate. A volume of $50\text{ }\mu\text{L}$ of $2.7\text{ }\mu\text{M}$ phenazine methosulphate (PMS) was added to each well. The radical scavenging activity of the samples was monitored with a Synergy HT Multi-detection Microplate Reader operated by GEN5TM (Biotek, Bad Friedrichshall, Germany), in kinetic function, at room temperature, for 2 min, at 562 nm. All reagents were dissolved in phosphate buffer ($19\text{ }\mu\text{M}$, pH 7.4). Three independent assays were performed in triplicate. GA was used as positive control. The results were expressed as a percentage of radical scavenging in comparison to the untreated control. The results for the calculated IC values were expressed as mean \pm SD ($\mu\text{g/mL}$) of at least three independent assays performed in duplicate. The IC values and the corresponding dose–response curves were calculated with Graphpad Prism[®] software (version 9, for MacOS).

4.5.2. Hyaluronidase Inhibition

HAase Inhibition assay was determined as proposed by Ferreres et al., [50] with some modifications. The acetone extracts were resuspended in DMSO, while the aqueous extracts were resuspended in water. Six serial dilutions were prepared for each extract (0.125 to 1.0 mg mL^{-1} and 0.063 to 1.0 mg mL^{-1} , respectively) and tested to evaluate the extracts' behavior and IC values. Twenty-five microliters of each sample were added to each reaction tube. A volume of $175\text{ }\mu\text{L}$ of HA solution (0.7 mg mL^{-1} in water:buffer, 5:2 *v/v*, kept at $37\text{ }^\circ\text{C}$) was added to each reaction tube and gently mixed. The reaction started by adding $25\text{ }\mu\text{L}$ of HAase (900 U mL^{-1} in NaCl 0.9%). After 30 min incubation at $37\text{ }^\circ\text{C}$, the enzymatic reaction was stopped with $25\text{ }\mu\text{L}$ of disodium tetraborate 0.8 M, followed by subsequent heating for 3 min in a boiling water bath. After cooling to room temperature, $375\text{ }\mu\text{L}$ of DMBA solution was added and gently mixed (2 g of DMAB dissolved in a mixture of 2.5 mL of 10 N HCl and 17.5 mL of glacial acetic acid and further diluted 1:2 with glacial acetic acid immediately before use). The tubes were incubated at $37\text{ }^\circ\text{C}$ for 20 min and the absorbance of the colored product was measured at 560 nm in a Synergy HT Multi-detection microplate reader (Biotek, Bad Friedrichshall, Germany) operated by GEN5TM software. Negative control was performed in the absence of extract. Disodium cromoglycate (DSCG) was used as positive control. Three independent assays were performed in triplicate.

4.5.3. Elastase Inhibition

Porcine pancreatic elastase inhibition assay was slightly modified from those proposed by Mota et al. [51]. Acetonic and aqueous extracts were resuspended in DMSO and water, respectively. Briefly, in a 96-well plate, $50\text{ }\mu\text{L}$ of extract (0.05 to 0.4 mg mL^{-1} for each extract) was mixed with $90\text{ }\mu\text{L}$ of HEPES buffer (0.1 M), $10\text{ }\mu\text{L}$ of N-succinyl-Ala-Ala-Ala *p*-nitroanilide substrate ($100\text{ }\mu\text{M}$), $70\text{ }\mu\text{L}$ of acetate buffer (200 mM), and $30\text{ }\mu\text{L}$ of elastase (1 U mL^{-1}). The plate was incubated at $37\text{ }^\circ\text{C}$ for 10 min, and the absorbance of the reaction product was measured at 405 nm, in a Synergy HT Multi-detection microplate reader (Biotek, Bad Friedrichshall) operated by GEN5TM. Negative control was performed in the absence of extract, and ascorbic acid was used as positive control. Three independent assays were performed in triplicate. The results were expressed as a percentage of enzyme inhibition in comparison to the untreated control.

4.5.4. Collagenase Inhibition

Collagenase inhibition assay was determined as reported by Andrade et al. [52] with some modifications. The substrate *N*-(3-furyl-acryloyl)-Leu-Gly-Pro-Ala (FALGPA) 1 mM was dissolved in tricine buffer (50 mM, pH 7.5). Collagenase enzyme was prepared in the buffer at 1 U mL⁻¹. First, 10 µL of extract dilutions (0.75 and 1.0 mg mL⁻¹), 45 µL of tricine buffer, and 50 µL of collagenase (1 U mL⁻¹) were added to a 96-well plate and kept at 37 °C for 15 min. The reaction was started by adding 120 µL of FALGPA. The substrate hydrolysis was monitored with a Synergy HT Multi-detection Microplate Reader operated by GEN5™ (Biotek, Bad Friedrichshall, Germany), operating in kinetic function, at room temperature, for 8 min, at 345 nm. Negative control was performed in the absence of extract, and gallic acid was used as positive control. Results were expressed as percentage of enzyme inhibition in comparison to the untreated control. The assay was performed in triplicate.

4.5.5. Tyrosinase Inhibition

The tyrosinase inhibition assay was performed according to Adhikari and co-workers [53], with some modifications. Five serial dilutions, from 0.063 to 1.0 mg mL⁻¹, were prepared for each extract. Briefly, in a 96-well plate, 20 µL of each extract was mixed with 100 µL of tyrosinase (30 U mL⁻¹ in phosphate buffer). Acetonic extracts and aqueous extracts were resuspended in DMSO and water, respectively. The mixture was incubated at 30 °C during 8 min. After that, 80 µL of L-DOPA (L-3,4-dihydroxyphenylalanine) solution (2.5 mM in phosphate buffer) was added, and the absorption was immediately read (T0) with a Synergy HT Multi-detection microplate reader (Biotek, Bad Friedrichshall, Germany) operated by GEN5™ software, at 475 nm. After 8 min, the absorbance was measured again (T8). Negative control was performed in the absence of extract, and kojic acid was used as positive control. Three independent assays were performed in triplicate. The results were expressed as a percentage of enzyme inhibition in comparison to the untreated control.

4.6. Statistical Analysis

Statistical analysis was performed using IBM SPSS STATISTICS software, version 28.0.1.0, IBM Corporation, New York, NY, USA (2021). Data were analyzed for normality and homogeneity of variances using Kolmogorov–Smirnov and Leven's tests, and then submitted to one-way ANOVA, using a Tukey's HSD (honest significant difference) as a post hoc test. A Pearson correlation test was used to compare normalized expression data between the chemical profile and the biological activities of cyanobacteria extracts.

5. Conclusions

Many studies have shown that, to protect themselves from the damage associated with extreme environmental conditions, such as high desiccation and UV radiation, cyanobacteria produce compounds such as carotenoids, phenolic compounds and phycobiliproteins.

This study revealed that the cyanobacteria strains evaluated stood out regarding cosmetic purposes and sustainability. The extractions were performed sequentially, allowing the obtention of different bioactive extracts from the same biomass. This increases biomass rentability, has a lower environmental impact and becomes economically attractive. The aqueous extracts of *Synechococcales* cyanobacterium LEGE 181157 and *Synechococcales* cyanobacterium LEGE 181150 strains, rich in phycobiliproteins, showed antioxidant activity, reaching an attractive IC₅₀ for superoxide radical scavenging. Furthermore, these extracts had the ability to inhibit the enzyme HAase. As such, they seem interesting for antioxidant purposes. Moreover, PBPs, the major components of aqueous extracts, are photoprotective compounds, and their natural and attractive colors may add value in cosmetic products. On the other hand, phenols and carotenoids present in the acetonic extracts, especially in *Leptothoe* sp. LEGE 181156 and *Synechococcales* cyanobacterium LEGE 181150, demonstrated potential to preserve dermal matrix components through the inhibition of MMPs. Overall, the results obtained drive the studies on cyanobacteria

towards the cosmetic industry, as a promising source of natural extracts with potential in cosmetic formulations.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/xxx/s1>, Figure S1: Keratinocyte (HaCat) viability after 24 and 48 h of incubation with cyanobacteria aqueous (a–g) and acetone (h–n) extracts. (a,h) *Cyanobium* sp. LEGE 07175, (b,i) Synechococcales cyanobacterium LEGE 181150, (c,j) *Leptothoe* sp. LEGE 181156 and (d,k) Synechococcales cyanobacterium LEGE 181157, (e,l) Synechococcales cyanobacterium LEGE 181158, (f,m) Oscillatoriales cyanobacterium LEGE 181159, (g,n) *Leptothoe* sp. LEGE 181155. Results are expressed as % of MTT reduction vs. the untreated control. DMSO (20%) represents the positive control. Results are expressed as mean \pm SD of at least three independent assays, performed in quadruplicate. Statistical differences at * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ (One-way ANOVA; Tukey's HSD multiple comparisons test). Figure S2: Fibroblast (3T3L1) viability after 24 and 48 h of incubation with cyanobacteria aqueous (a–g) and acetone (h–n) extracts. (a,h) *Cyanobium* sp. LEGE 07175, (b,i) Synechococcales cyanobacterium LEGE 181150, (c,j) *Leptothoe* sp. LEGE 181156 and (d,k) Synechococcales cyanobacterium LEGE 181157, (e,l) Synechococcales cyanobacterium LEGE 181158, (f,m) Oscillatoriales cyanobacterium LEGE 181159, (g,n) *Leptothoe* sp. LEGE 181155. Results are expressed as % of MTT reduction vs. the untreated control. DMSO (20%) represents the positive control. Results are expressed as mean \pm SD of at least three independent assays, performed in quadruplicate. Statistical differences at * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ (One-way ANOVA; Tukey's HSD multiple comparisons test). Figure S3: Endothelial cell (hCMEC/d3) viability after 24 and 48 h of incubation with cyanobacteria aqueous (a–g) and acetone (h–n) extracts. (a,h) *Cyanobium* sp. LEGE 07175, (b,i) Synechococcales cyanobacterium LEGE 181150, (c,j) *Leptothoe* sp. LEGE 181156 and (d,k) Synechococcales cyanobacterium LEGE 181157, (e,l) Synechococcales cyanobacterium LEGE 181158, (f,m) Oscillatoriales cyanobacterium LEGE 181159, (g,n) *Leptothoe* sp. LEGE 181155. Results are expressed as % of MTT reduction vs. the untreated control. DMSO (20%) represents the positive control. Results are expressed as mean \pm SD of at least three independent assays, performed in quadruplicate. Statistical differences at * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ (One-way ANOVA, Tukey's HSD multiple comparisons test).

Author Contributions: Conceptualization, R.M. and G.L.; methodology, J.M. (Janaina Morone), G.L., J.M. (João Morais) and J.N.; data analysis, J.M. (Janaina Morone) and G.L.; investigation, J.M. (Janaina Morone); writing—original draft preparation, J.M. (Janaina Morone); writing—review and editing, J.M. (Janaina Morone), G.L., J.M. (João Morais), J.N., R.M. and V.V.; supervision, G.L., V.V. and R.M.; funding acquisition, V.V. All authors have read and agreed to the published version of the manuscript.

Funding: This work was partially performed in the framework of the project ALGAVALOR-MicroALGAs: integrated production and valorization of biomass and its various applications-SI I&DT n.o 352234-supported by the PORTUGAL 2020 through the European Regional Development Fund; and supported by the FCT Projects UIDB/04423/2020 and UIDP/04423/2020.

Institutional Review Board Statement: Not applicable.

Data Availability Statement: Data are available upon request to the corresponding author.

Acknowledgments: Janaína Morone is grateful to Fundação para a Ciência e a Tecnologia (FCT) for the Grant n.º. UI/BD/150903/2021. Graciliana Lopes thanks FCT for the financial support for her work contract through the Scientific Employment Stimulus-Individual Call (CEECIND/01768/2021). The authors acknowledge the support and the use of resources of MIRRI-ERIC, specifically of the Portuguese infrastructure node of the Microbial Resource Research Infrastructure (MIRRI-PT) NORTE-01-0145-FEDER-084445, financed by the European Regional Development Fund (ERDF) and national funds UIDB/04423/2021 and UIDP/04423/2021 through FCT/MCTES. Sampling was supported through project EMERTOX (grant 734748), funded by H2020-MSCA-RISE 2016. We also thank the Ministério da Agricultura e Ambiente de Cabo Verde under the sampling authorization of the National Environment Directorate N°094/2019.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. Morone, J.; Lopes, G.; Oliveira, B.; Vasconcelos, V.; Martins, R. *Cyanobacteria in Cosmetics: A Natural Alternative for Anti-Aging Ingredients*; Academic Press: Cambridge, MA, USA, 2022; pp. 257–286.
2. Morone, J.; Alfeus, A.; Vasconcelos, V.; Martins, R. Revealing the potential of cyanobacteria in cosmetics and cosmeceuticals—New bioactive approach. *Algal Res.* **2019**, *41*, 101541. [[CrossRef](#)]
3. Chen, L.; Li, D.; Song, L.; Hu, C.; Wang, G.; Lui, Y. Effects of Salt Stress on Carbohydrate Metabolism in Desert Soil Alga *Microcoleus vaginatus* Gom. *J. Integr. Plant Biol.* **2006**, *48*, 914–919. [[CrossRef](#)]
4. Favas, R.; Morone, J.; Martins, R.; Vasconcelos, V.; Lopes, G. Cyanobacteria and microalgae bioactive compounds in skin-ageing: Potential to restore extracellular matrix filling and overcome hyperpigmentation. *J. Enzym. Inhib. Med. Chem.* **2021**, *36*, 1829–1838. [[CrossRef](#)] [[PubMed](#)]
5. Pagels, F.; Guedes, A.; Vasconcelos, V.; Lopes, G. Anti-inflammatory compounds from cyanobacteria. In *The Pharmacological Potential of Cyanobacteria*; Academic Press: Cambridge, MA, USA, 2022.
6. Assunção, J.; Amaro, H.; Malcata, F.; Guedes, A. Cyanobacterial pigments: Photosynthetic function and biotechnological purposes. In *The Pharmacological Potential of Cyanobacteria*; Academic Press: Cambridge, MA, USA, 2022; pp. 201–256.
7. Singh, S.K.; Kaur, R.; Bansal, A.; Kapur, S.; Sundaram, S. Biotechnological exploitation of cyanobacteria and microalgae for bioactive compounds. In *Biotechnological Production of Bioactive Compounds*; Elsevier: Amsterdam, The Netherlands, 2020.
8. Nowruzi, B.; Sarvari, G.; Blanco, S. The cosmetic application of cyanobacterial secondary metabolites. *Algal Res.* **2020**, *49*, 101959. [[CrossRef](#)]
9. Pagels, F.; Guedes, A.C.; Amaro, H.M.; Kijjoo, A.; Vasconcelos, V. Phycobiliproteins from cyanobacteria: Chemistry and biotechnological applications. *Biotechnol. Adv.* **2019**, *37*, 422–443. [[CrossRef](#)] [[PubMed](#)]
10. Jerez-Martel, I.; García-Poza, S.; Rodríguez-Martel, G.; Rico, M.; Afonso-Olivares, C.; Gómez-Pinchetti, J.L. Phenolic Profile and Antioxidant Activity of Crude Extracts from Microalgae and Cyanobacteria Strains. *J. Food Qual.* **2017**, *2017*, 2924508. [[CrossRef](#)]
11. Tarasuntisuk, S.; Patipong, T.; Hibino, T.; Waditee-Sirisattha, R.; Kageyama, H. Inhibitory effects of mycosporine-2-glycine isolated from a halotolerant cyanobacterium on protein glycation and collagenase activity. *Lett Appl. Microbiol.* **2018**, *67*, 314–320. [[CrossRef](#)]
12. Keller, L.; Canuto, K.M.; Liu, C.; Suzuki, B.M.; Almaliti, J.; Sikandar, A.; Naman, C.B.; Glukhov, E.; Luo, D.; Duggan, B.M.; et al. Tutuilamides A-C: Vinyl-Chloride-Containing Cyclodepsipeptides from Marine Cyanobacteria with Potent Elastase Inhibitory Properties. *ACS Chem. Biol.* **2020**, *15*, 751–757. [[CrossRef](#)]
13. Yamaguchi, Y.; Koketsu, M. Isolation and analysis of polysaccharide showing high hyaluronidase inhibitory activity in *Nostocopsis lobatus* MAC0804NAN. *J. Biosci. Bioeng.* **2016**, *121*, 345–348. [[CrossRef](#)]
14. Favas, R.; Morone, J.; Martins, R.; Vasconcelos, V.; Lopes, G. Cyanobacteria Secondary Metabolites as Biotechnological Ingredients in Natural Anti-Aging Cosmetics: Potential to Overcome Hyperpigmentation, Loss of Skin Density and UV Radiation-Deleterious Effects. *Mar. Drugs* **2022**, *20*, 183. [[CrossRef](#)]
15. CODIF. Phormiskin Bioprotech G. Available online: <https://www.codif-tn.com/en/principesactifs/phormiskin-bioprotech-g/> (accessed on 1 May 2022).
16. Gao, X.; Jing, X.; Liu, X.; Lindblad, P. Biotechnological Production of the Sunscreen Pigment Scytonemin in Cyanobacteria: Progress and Strategy. *Mar. Drugs* **2021**, *19*, 129. [[CrossRef](#)] [[PubMed](#)]
17. Lopes, G.; Silva, M.; Vasconcelos, V. *The Pharmacological Potential of Cyanobacteria*; Elsevier: Amsterdam, The Netherlands, 2022.
18. Morone, J.; Lopes, G.; Preto, M.; Vasconcelos, V.; Martins, R. Exploitation of Filamentous and Picoplanktonic Cyanobacteria for Cosmetic Applications: Potential to Improve Skin Structure and Preserve Dermal Matrix Components. *Mar. Drugs* **2020**, *18*, 486. [[CrossRef](#)] [[PubMed](#)]
19. Kusama, Y.; Inoue, S.; Jimbo, H.; Takaichi, S.; Sonoike, K.; Hihara, Y.; Nishiyama, Y. Zeaxanthin and Echinone Protect the Repair of Photosystem II from Inhibition by Singlet Oxygen in *Synechocystis* sp. PCC 6803. *Plant Cell. Physiol.* **2015**, *56*, 906–916. [[CrossRef](#)] [[PubMed](#)]
20. Krinsky, N.I.; Landrum, J.T.; Bone, R.A. Biologic mechanisms of the protective role of lutein and zeaxanthin in the eye. *Annu. Rev. Nutr.* **2003**, *23*, 171–201. [[CrossRef](#)] [[PubMed](#)]
21. Lopes, G.; Clarinha, D.; Vasconcelos, V. Carotenoids from Cyanobacteria: A Biotechnological Approach for the Topical Treatment of Psoriasis. *Microorganisms* **2020**, *8*, 302. [[CrossRef](#)]
22. Palinska, K.A.; Deventer, B.; Hariri, K.; Łotocka, M. A taxonomic study on Phormidium-group (cyanobacteria) based on morphology, pigments, RAPD molecular markers and RFLP analysis of the 16S rRNA gene fragment. *Fottea* **2011**, *11*, 41–55. [[CrossRef](#)]
23. Amaro, H.M.; Fernandes, F.; Valentao, P.; Andrade, P.B.; Sousa-Pinto, I.; Malcata, F.X.; Guedes, A.C. Effect of Solvent System on Extractability of Lipidic Components of *Scenedesmus obliquus* (M2-1) and *Gloeotheca* sp. on Antioxidant Scavenging Capacity Thereof. *Mar. Drugs* **2015**, *13*, 6453–6471. [[CrossRef](#)]
24. Patias, L.D.; Fernandes, A.S.; Petry, F.C.; Mercadante, A.Z.; Jacob-Lopes, E.; Zepka, L.Q. Carotenoid profile of three microalgae/cyanobacteria species with peroxy radical scavenger capacity. *Food Res. Int.* **2017**, *100*, 260–266. [[CrossRef](#)]
25. Hashtroudi, M.S.; Shariatmadari, Z.; Riahi, H.; Ghassempour, A. Analysis of *Anabaena vaginicola* and *Nostoc calcicola* from Northern Iran, as rich sources of major carotenoids. *Food Chem.* **2013**, *136*, 1148–1153. [[CrossRef](#)]

26. Arnum, V.; Susan, D. Vitamin A. In *Kirk-Othmer Encyclopedia of Chemical Technology*; John Wiley & Sons: New York, NY, USA, 2000; pp. 99–107.
27. Szymanski, L.; Skopek, R.; Palusinska, M.; Schenk, T.; Stengel, S.; Lewicki, S.; Kraj, L.; Kaminski, P.; Zelent, A. Retinoic Acid and Its Derivatives in Skin. *Cells* **2020**, *9*, 2660. [[CrossRef](#)]
28. Abiri, B.; Vafa, M. *Antioxidant Vitamins in Acute Lymphoblastic Leukemia*, 2nd ed.; Academic Press: Cambridge, MA, USA, 2021; Chapter 48.
29. Steiger, S.; Schafer, L.; Sandmann, G. High-light-dependent upregulation of carotenoids and their antioxidative properties in the cyanobacterium *Synechocystis* PCC 6803. *J. Photochem. Photobiol.* **1999**, *52*, 14–18. [[CrossRef](#)]
30. Koller, M.; Muhr, A.; Braunegg, G. Microalgae as versatile cellular factories for valued products. *Algal Res.* **2014**, *6*, 52–63. [[CrossRef](#)]
31. Pagels, F.; Salvaterra, D.; Amaro, H.M.; Lopes, G.; Sousa-Pinto, I.; Vasconcelos, V.; Guedes, A.C. Bioactive potential of *Cyanobium* sp. pigment-rich extracts. *J. Appl. Phycol.* **2020**, *32*, 3031–3040. [[CrossRef](#)]
32. Rahman, M.M.; Rahaman, M.S.; Islam, M.R.; Rahman, F.; Mithi, F.M.; Alqahtani, T.; Almikhlaifi, M.A.; Alghamdi, S.Q.; Alruwaili, A.S.; Hossain, M.S.; et al. Role of Phenolic Compounds in Human Disease: Current Knowledge and Future Prospects. *Molecules* **2021**, *27*, 233. [[CrossRef](#)]
33. Boo, Y.C. Can Plant Phenolic Compounds Protect the Skin from Airborne Particulate Matter? *Antioxidants* **2019**, *8*, 379. [[CrossRef](#)]
34. Orhan, I.E.; Khan, M.T.H. Flavonoid Derivatives as Potent Tyrosinase Inhibitors—A Survey of Recent Findings between 2008–2013. *Top. Med. Chem.* **2014**, *14*, 1486–1493. [[CrossRef](#)]
35. Ahsan, H. Immunopharmacology and immunopathology of peptides and proteins in personal products. *J. Immunoass. Immunochem.* **2019**, *40*, 439–447. [[CrossRef](#)]
36. Hossain, M.F.; Ratnayake, R.R.; Meerajini, K.; Wasantha Kumara, K.L. Antioxidant properties in some selected cyanobacteria isolated from fresh water bodies of Sri Lanka. *Food Sci. Nutr.* **2016**, *4*, 753–758. [[CrossRef](#)]
37. Pumas, C.; Vacharapiyasophon, P.; Peerapornpisal, Y.; Leelapornpisid, P.; Boonchum, W.; Ishii, M.; Khanongnuch, C. Thermostability of phycobiliproteins and antioxidant activity from four thermotolerant cyanobacteria. *Phycol. Res.* **2011**, *59*, 166–174. [[CrossRef](#)]
38. Pez Jaeschke, D.; Rocha Teixeira, I.; Damasceno Ferreira Marczak, L.; Domeneghini Mercali, G. Phycocyanin from Spirulina: A review of extraction methods and stability. *Food Res. Int.* **2021**, *143*, 110314. [[CrossRef](#)]
39. Hemlata; Fatma, T. Screening of cyanobacteria for phycobiliproteins and effect of different environmental stress on its yield. *Bull. Environ. Contam. Toxicol.* **2009**, *83*, 509–515. [[CrossRef](#)] [[PubMed](#)]
40. Eriksen, N.T. Production of phycocyanin—A pigment with applications in biology, biotechnology, foods and medicine. *Appl. Microbiol. Biotechnol.* **2008**, *80*, 1–14. [[CrossRef](#)] [[PubMed](#)]
41. Freitas-Rodriguez, S.; Folgueras, A.R.; Lopez-Otin, C. The role of matrix metalloproteinases in aging: Tissue remodeling and beyond. *Biochim. Biophys. Acta Mol. Cell Res.* **2017**, *1864*, 2015–2025. [[CrossRef](#)] [[PubMed](#)]
42. Del Bino, S.; Bernerd, F. Variations in skin colour and the biological consequences of ultraviolet radiation exposure. *Br. J. Derm.* **2013**, *169*, 33–40. [[CrossRef](#)]
43. Pagels, F.; Almeida, C.; Vasconcelos, V.; Guedes, A.C. Cosmetic Potential of Pigments Extracts from the Marine Cyanobacterium *Cyanobium* sp. *Mar. Drugs* **2022**, *20*, 481. [[CrossRef](#)]
44. Ramos, V.; Morais, J.; Castelo-Branco, R.; Pinheiro, A.; Martins, J.; Regueiras, A.; Pereira, A.L.; Lopes, V.R.; Frazao, B.; Gomes, D.; et al. Cyanobacterial diversity held in microbial biological resource centers as a biotechnological asset: The case study of the newly established LEGE culture collection. *J. Appl. Phycol.* **2018**, *30*, 1437–1451. [[CrossRef](#)]
45. Ferreira, L.; Morais, J.; Preto, M.; Silva, R.; Urbatzka, R.; Vasconcelos, V.; Reis, M. Uncovering the Bioactive Potential of a Cyanobacterial Natural Products Library Aided by Untargeted Metabolomics. *Mar. Drugs* **2021**, *19*, 633. [[CrossRef](#)]
46. Kotai, J. *Instructions for Preparation of Modified Nutrient Solution Z8 for Algae*; Norwegian Institute for Water Research: Oslo, Norway, 1972; Volume 15, pp. 155–164.
47. Fernandes, F.; Barbosa, M.; Oliveira, A.P.; Azevedo, I.C.; Sousa-Pinto, I.; Valentão, P.; Andrade, P.B. The pigments of kelps (*Ochrophyta*) as part of the flexible response to highly variable marine environments. *J. Appl. Phycol.* **2016**, *28*, 3689–3696. [[CrossRef](#)]
48. Barroso, M.F.; Ramalhosa, M.J.; Alves, R.C.; Dias, A.; Soares, C.M.D.; Oliva-Teles, M.T.; Delerue-Matos, C. Total antioxidant capacity of plant infusions: Assessment using electrochemical DNA-based biosensor and spectrophotometric methods. *Food Control* **2016**, *68*, 153–161. [[CrossRef](#)]
49. Barbosa, M.; Fernandes, F.; Carlos, M.J.; Valentão, P.; Andrade, P.B. Adding value to marine invaders by exploring the potential of *Sargassum muticum* (Yendo) Fensholt phlorotannin extract on targets underlying metabolic changes in diabetes. *Algal Res.* **2021**, *59*, 102455. [[CrossRef](#)]
50. Ferreres, F.; Lopes, G.; Gil-Izquierdo, A.; Andrade, P.B.; Sousa, C.; Mouga, T.; Valentao, P. Phlorotannin extracts from fucales characterized by HPLC-DAD-ESI-MSn: Approaches to hyaluronidase inhibitory capacity and antioxidant properties. *Mar. Drugs* **2012**, *10*, 2766–2781. [[PubMed](#)]

51. Mota, F.A.R.; Pereira, S.A.P.; Araujo, A.; Saraiva, M. Evaluation of Ionic Liquids and Ionic Liquids Active Pharmaceutical Ingredients Inhibition in Elastase Enzyme Activity. *Molecules* **2021**, *26*, 200. [[CrossRef](#)] [[PubMed](#)]
52. Andrade, J.M.; Dominguez-Martin, E.M.; Nicolai, M.; Faustino, C.; Rodrigues, L.M.; Rijo, P. Screening the dermatological potential of plectranthus species components: Antioxidant and inhibitory capacities over elastase, collagenase and tyrosinase. *J. Enzym. Inhib. Med. Chem.* **2021**, *36*, 257–269. [[CrossRef](#)] [[PubMed](#)]
53. Adhikari, A.; Devkota, H.P.; Takano, A.; Masuda, K.; Nakane, T.; Basnet, P.; Skalko-Basnet, N. Screening of Nepalese crude drugs traditionally used to treat hyperpigmentation: In vitro tyrosinase inhibition. *Int. J. Cosmet. Sci.* **2008**, *30*, 353–360. [[CrossRef](#)] [[PubMed](#)]