

Supplementary information for

Heteronemin and Tetrac Induce Anti-proliferation by Blocking EGFR-mediated Signaling in Colorectal Cancer Cells

Sukanya Unson¹, Tung-Cheng Chang², Yung-Ning Yang³, Shwu-Huey Wang⁴, Chi-Hung Huang⁵, Dana R. Crawford⁶, Haw-Ming Huang⁷, Zi-Lin Li^{8,9}, Hung-Yun Lin^{9,10,11,12,13*}, Jacqueline Whang-Peng^{12*}, Kuan Wang³, Paul J. Davis^{13, 14}, Wen-Shan Li^{15, 16}

Correspondence to: jqpeng@nhri.edu.tw and linhy@tmu.edu.tw

This PDF file includes

Supplementary figures S1 and S2.




Figure S1. Heatmap of cell motility genes that showed significant ($p < 0.05$) upregulation or downregulation. HCT-116 cells were treated with heteronemin, tetrac, and their combination for 24 h. mRNA expression levels were identified by a NanoString® analysis using the nCounter PanCancer Progression Panel.




Figure S2. Heatmap of gene expressions involved in the cell cycle showed significant ($p < 0.05$) upregulation or downregulation. HCT-116 cells were treated with heteronemin, tetrac, and their combination for 24 h. mRNA expressions were identified by NanoString® analysis using the nCounter PanCancer Progression Panel.