

Protection activity of 1,4-naphthoquinones in rotenone-induced models of neurotoxicity.

Irina Agafonova^{1§}, Ekaterina Chingizova^{1§}, Elena Chaikina¹, Ekaterina Menchinskaya¹, Sergey Kozlovskiy¹, Galina Likhatskaya¹, Yuri Sabutski¹, Sergey Polonik¹, Dmitry Aminin^{1,2} and Evgeny Pislyagin^{1*}

¹ G.B. Elyakov Pacific Institute of Bioorganic Chemistry, Far-Eastern Branch of the Russian Academy of Science, 690022 Vladivostok, Russia

² Department of Biomedical Science and Environmental Biology, Kaohsiung Medical University, No. 100, Shichuan 1st Road, Sanmin District, Kaohsiung City 80708, Taiwan

[§] Equal contribution

* Correspondence: pislyagin@hotmail.com (EP); Tel.: +7(4232)319932

Table of Contents

Figure S1. ¹H and ¹³C NMR spectra of compound **4** (U-573)

Figure S2. ¹H and ¹³C NMR spectra of compound **8** (U-443)

Figure S3. HPLC chromatogram of compound **4** (U-573)

Figure S4. HPLC chromatogram of compound **8** (U-443)





Figure S1. ¹H and ¹³C NMR spectra of compound 4 (U-573) (solvent - DMSO-d₆)




Figure S2. ¹H and ¹³C NMR spectra of compound 8 (U-443) (solvent – CDCl₃)




Figure S3. HPLC chromatogram of compound 4 (U-573)




Figure S4. HPLC chromatogram of compound 8 (U-443)