

Supplemental Material

Optimization of the preparation process of glucuronomannan oligosaccharides and their effects on the gut microbiota in MPTP-induced PD model mice

Baoxiang Wang^{1,2}, Lihua Geng^{2,3,4,*}, Jing Wang^{2,3,4}, Yuxi Wei¹, Changhui Yan^{2,5}, Ning Wu^{2,4}, Yang Yue^{2,3,4}, Quanbin Zhang^{2,3,4,*}

¹ College of Life Sciences, Qingdao University, 308 Ningxia Road, Qingdao 266003, China; wangbaoxiangcn@163.com; yuxiw729@163.com

² CAS and Shandong Province Key Laboratory of Experimental Marine Biology, Institute of Oceanology, Chinese Academy of Sciences, Qingdao 266071, China; jingwang@qdio.ac.cn (J.W.); yich2747141850@163.com (C.Y.); wuning@qdio.ac.cn (N.W.); yueyang@qdio.ac.cn (Y.Y.)

³ Laboratory for Marine Biology and Biotechnology, Qingdao Marine Science and Technology Center, Qingdao 266237, China

⁴ Center for Ocean Mega-Science, Chinese Academy of Sciences, Qingdao 266071, China

⁵ Key Laboratory of Optic-electric Sensing and Analytical Chemistry for Life Science, MOE, College of Chemistry and Molecular Engineering, Qingdao University of Science and Technology, Qingdao 266042, PR China; yich2747141850@163.com

* Correspondence: lhgeng@qdio.ac.cn, Tel: +86-532-8289-8703; qbzhang@qdio.ac.cn, Tel: +86-532-8289-8708.




Figure S1. ^1H -NMR spectra of GMn. (A) ^1H -NMR spectra of G2 (GM1) (B) ^1H -NMR spectra of G3 (GM2).