



Article

The Effects of PM_{2.5} from Asian Dust Storms on Emergency Room Visits for Cardiovascular and Respiratory Diseases

Ssu-Ting Liu ¹, Chu-Yung Liao ², Cheng-Yu Kuo ³ and Hsien-Wen Kuo ^{1,4,*}

¹ Institute of Environmental and Occupational Health Sciences, National Yang-Ming University, 155 Li-Nong 2nd Street, Taipei 114, Taiwan; rs86229@yahoo.com.tw

² Department of Early Childhood Educare, College of Health, Chung Chou University of Science and Technology, Changhua 510, Taiwan; cyliao@dragon.ccut.edu.tw

³ College of Arts and Sciences, University of San Francisco, San Francisco, CA 94117, USA; taiwan22361397@gmail.com

⁴ School of Public Health, National Defense Medical Center, Taipei 114, Taiwan

* Correspondence: hwkuo@ym.edu.tw; Tel.: +886-2-28272294; Fax: +886-2-28278254

Academic Editor: Paul B. Tchounwou

Received: 23 January 2017; Accepted: 13 April 2017; Published: 16 April 2017

Abstract: A case-crossover study examined how PM_{2.5} from Asian Dust Storms (ADS) affects the number of emergency room (ER) admissions for cardiovascular diseases (CVDs) and respiratory diseases (RDs). Our data indicated that PM_{2.5} concentration from ADS was highly correlated with ER visits for CVDs and RDs. The odds ratios (OR) increased by 2.92 (95% CI: 1.22–5.08) and 1.86 (95% CI: 1.30–2.91) per 10 µg/m³ increase in PM_{2.5} levels, for CVDs and RDs, respectively. A 10 µg/m³ increase in PM_{2.5} from ADSs was significantly associated with an increase in ER visits for CVDs among those 65 years of age and older (an increase of 2.77 in OR) and for females (an increase of 3.09 in OR). In contrast, PM_{2.5} levels had a significant impact on RD ER visits among those under 65 years of age (OR = 1.77). The risk of ER visits for CVDs increased on the day when the ADS occurred in Taiwan and the day after (lag 0 and lag 1); the corresponding risk increase for RDs only increased on the fifth day after the ADS (lag 5). In Taiwan's late winter and spring, the severity of ER visits for CVDs and RDs increases. Environmental protection agencies should employ an early warning system for ADS to reduce high-risk groups' exposure to PM_{2.5}.

Keywords: air pollution; PM_{2.5}; emergency room; Asian dust storms; case-crossover

1. Introduction

Asian dust storms (ADS) frequently occur in northern China in late winter and spring (from November to May). Asian dust follows westerly winds that move toward the east, impacting Japan, Korea, and other neighboring areas. Only under specific meteorological conditions will an ADS impact Taiwan. On 25 April 2009, ADS arrived in Taiwan, bringing with it PM₁₀ (particles are represented by the mass concentration of particles smaller than 10 µm) exceeding 1000 µg/m³. This was the most serious case in Taiwan. Since then, the situation has worsened, along with the desertification of China's Inner Mongolia, a global increase in droughts, and the uneven distribution of rain. These increases in both the frequency and severity of ADS have impacted Taiwan [1], worsening air quality and visibility. The increase in the number of particulates floating in the air has led to an increase a number of problems, including allergic nasal inflammation, asthma, eye problems, skin allergies, and skin irritability. Asian Dust Storms in Taiwan have led to an increase of 7.66% in the risk for respiratory diseases (RDs) one day after the ADS (lag 1), 4.92% in total deaths two days after the ADS (lag 2), and 2.59% in circulatory diseases two days after the ADS (lag 2). None of these effects

were statistically significant [2]. However, there is a statistically significant association between ADS events and daily primary intracerebral hemorrhagic stroke visits three days after the event (relative risk of 1.15; 95% CI: 1.01–10.10) [3].

Previous environmentally epidemiological studies have shown positive correlations of particulate matter (PM) on the risk of death [4–9], hospitalizations [3,10–17], and emergency room (ER) visits [8,18–21]. In Barcelona, Spain, a case-crossover study from 2003 to 2007 indicated that PM_{2.5} from car exhaust, gas combustion, hyponitrites, hydrosulfites, and road dust are correlated to CVD mortality ($p < 0.05$); for every 10 $\mu\text{g}/\text{m}^3$ increase in PM_{2.5} (the mass concentration of particles smaller than 2.5 μm), cardiovascular-related mortality increased by 2.9% (95% CI: 1.4–4.4) [9]. Previous research on PM has focused on its effects on deaths or hospital visits due to diseases. Few studies have investigated the effects of ADS on ER visits. In addition, the majority of previous ADSs studies used PM₁₀ as the exposure index. However, recent epidemiological studies [22,23] have found PM_{2.5} to pose a larger health threat due to its smaller size and larger surface area, which allows it to carry more toxic substances. Hence, the World Health Organization (WHO) has already recommended the replacement of PM₁₀ with PM_{2.5} for studies on air quality index [24]. Following that recommendation, Taiwan's EPA began measuring PM_{2.5} levels at its environmental monitoring stations and performing research on the health impacts of PM_{2.5}'s. Due to Taiwan's small geographic area and its intermixing of traffic pollutants with industry pollutants, the PM_{2.5} composition might differ from that in the United States or Europe [25,26]. Thus, the PM_{2.5} of other countries might not be the same as Taiwanese PM_{2.5}. Whereas the characteristics of PM_{2.5} in others countries might not represent exposure to PM_{2.5} in Taiwan, we performed a case-crossover design assessing the PM_{2.5} of ADS. This investigation particularly focused on the effects of PM_{2.5} has on the risk of ER visits for CVDs and RDs in central Taiwan.

2. Materials and Methods

2.1. Study Design

The case-crossover design is based on one proposed by Maclure in 1991 [27] to investigate the health effects of short-term exposure to hazards; it is a form of observational study design. This method has already been applied in the literature on the short-term effects of air pollution on health. In this design, each participant is compared to himself before or after the exposure. Because this design compares each participant to himself, it relieves researchers of the need to find a control. In addition, it allows for the control of each individual confounder. For these reasons, we chose the unidirectional case-crossover design for our study. In our study, the day of ER visits acted as the exposure period, while 14 days prior acted as the control period. To estimate relative risk, the exposure frequency during ADS periods just before outcome onset is compared with exposure frequencies during control times.

2.2. ER Visits Data

We used a National Health Insurance (NHI) database in central Taiwan as our dataset. NHI began in 1995, and as of 2013 has a coverage rate of 99.9%. It is thus representative of the entire Taiwanese population and its medical institutions. Demographic information in the registry of ED visit included gender, age and delivering time and diagnosis of diseases, etc. No other detailed personal information was recorded. According to the International Classification of Diseases, 9th Revision (ICD-9), we filtered the dataset to extract those with CVDs (ICD-9 codes 90-459) and those with RDs (ICD-9 codes 460-519). The study was conducted in accordance with the Declaration of Helsinki, and the protocol was approved by the Ethics Committee of the Institutional Review Board (IRB) of National Yang-Ming University (Project identification code: YM102007E).

2.3. Exposure Data and Meteorological Data

The so-called "Asian Dust Storm" (ADS) is a strong wind that collects large amounts of dust from the ground, creating a weather condition that negatively affects visibility. Generally, it is a result

of drought and desertification. Meteorologists define ADS which the visibility is below 1 km and levels of PM_{10} exceeded $150 \mu\text{g}/\text{m}^3$ in consecutive three-days period [1]. Our study relied on the data from 2006 to 2008 from nine Taiwan EPA air-quality monitoring stations, which fully cover cases' residential address. As per the Taiwanese EPA's definition of an ADS, we determined the presence of ADS if two conditions were met: a visibility of less than 1 kilometer and a PM_{10} level of over $150 \mu\text{g}/\text{m}^3$. With this definition, we selected 16 ADSs that occurred between 2006 and 2008, six events in 2006, eight in 2007, and two in 2008. Monitoring stations were fully automated and provided daily readings of sulfur dioxide (SO_2) (by ultraviolet fluorescence), PM_{10} (by β -ray absorption), nitrogen dioxide (NO_2) (by ultraviolet fluorescence), carbon monoxide (CO) (by non-dispersive infrared photometry), and ozone (O_3) (by ultraviolet photometry) levels. Daily meteorological information, including the mean temperature and mean humidity, was provided by the Central Weather Bureau. The monitoring of $PM_{2.5}$ was done by tapered element oscillating microbalance (TEOM), which began in 2006. TEOM allowed for two types: manual monitoring and automatic monitoring. As per the standard measurement guidelines for measuring air quality in Taiwan, we used manual monitoring to collect data on $PM_{2.5}$. To ensure consistency between the manual and automatic monitoring styles, we referenced the American EPA methodology, which suggests the application of simple linear regression on the manual and automatic data, allowing for the correction of the automatic data and immediate reporting of changes as a warning system. We calculated the average of the data in each 24-h period for each of the nine monitoring stations in central Taiwan, and then this data combined with ER visits databases.

2.4. Statistics Analysis

We began by organizing the data in Microsoft Excel 2013. Using SAS 9.3, we ran conditional logistic regression comparisons on the exposure and control periods. To understand the effects of short-term $PM_{2.5}$ exposure on CVDs and RDs in ER visits, we estimated the odds ratios (OR) and took their 95% confidence intervals (CI). Because the temperature changes between the seasons in Taiwan are not significant, we ignored the interaction effects of $PM_{2.5}$ and season on CVDs and RDs. In consideration of age and sex effects, we added an age variable (≥ 65 or < 65 years of age) and a sex variable. In addition, due to individual differences in the physical reaction to $PM_{2.5}$ exposure, we added a lag effect variable (lag 0 to lag 5). A lag effect of zero (lag 0) corresponds to the same day of exposure to $PM_{2.5}$ —that is, the day of arrival at the hospital for ER visits, which means the day of the individual CVDs or RDs event (depending on the time scale being studied) was the initial exposure period (lag 0) considered for that case. For the daily time scale analysis, we examined the association at the day of onset (lag 0 day) and the 1 to 5 before onset (lag 1, 2, 3, 4, 5). Single-pollution model, the two-pollution model, and the multi-pollution model were used in our analysis of $PM_{2.5}$ and other pollutants (SO_2 , CO, O_3 , NO_2 , NO and NO_x) effects on the ER visits for CVDs and RDs.

3. Results

Table 1 indicates significant differences for PM_{10} , $PM_{2.5}$, SO_2 , O_3 , average daily temperature (ADT), DTR (diurnal temperature range), and relative humidity (RH) between ADS days (index days) and comparison days. During ADS periods, PM_{10} , $PM_{2.5}$, SO_2 and O_3 were higher in concentration on index days. However, the average temperature, relative humidity, NO, and NO_x were lower. We found no significant differences in the levels of CO and NO_2 across index days and comparison days.

Table 2 uses Pearson's correlation coefficients to assess the correlations among different air pollutants. We found positive correlations between PM_{10} and $PM_{2.5}$ ($r = 0.76$), SO_2 ($r = 0.51$), as well as CO ($r = 0.34$); we found negative correlations between PM_{10} and RH ($r = -0.34$). $PM_{2.5}$ had positive correlations with SO_2 and CO ($r = 0.45$ and $r = 0.54$, respectively). SO_2 was positively correlated with CO, NO and NO_x , with correlation coefficients of 0.55, 0.36 and 0.28; it had negative correlations with O_3 , ADT and RH ($r = -0.32$, $r = -0.31$ and $r = -0.48$). O_3 had negative correlations with CO

($r = -0.63$), NO_2 ($r = -0.75$), NO ($r = -0.69$), and NO_x ($r = -0.74$). ADT was positively correlated with RH ($r = 0.66$), though DTR was negatively correlated with RH ($r = -0.48$).

Table 1. Comparison of various pollutants and metrological data between index and comparisons days.

Variable	Index Days (Mean \pm SD)	Comparison Days (Mean \pm SD)	<i>p</i> -Value
PM ₁₀ ($\mu\text{g}/\text{m}^3$)	133.02 \pm 31.62	77.77 \pm 31.48	<0.001
PM _{2.5} ($\mu\text{g}/\text{m}^3$)	61.93 \pm 17.11	48.87 \pm 22.34	<0.001
SO ₂ (ppb)	7.16 \pm 2.45	5.66 \pm 2.78	<0.001
CO (ppm)	0.63 \pm 0.12	0.65 \pm 0.21	0.425
O ₃ (ppb)	32.44 \pm 9.90	23.31 \pm 8.65	<0.001
NO (ppb)	6.45 \pm 4.59	9.69 \pm 7.28	<0.001
NO ₂ (ppb)	25.14 \pm 8.21	25.28 \pm 8.53	0.881
NO _x (ppb)	31.50 \pm 12.52	34.93 \pm 14.65	0.021
ADT ($^{\circ}\text{C}$)	18.91 \pm 3.84	21.25 \pm 3.84	<0.001
DTR ($^{\circ}\text{C}$)	6.67 \pm 2.02	6.15 \pm 1.67	0.030
RH (%)	56.37 \pm 12.14	70.63 \pm 4.56	<0.001

Table 2. Correlations between concentrations of pollutants and meteorology on Asian Dust Storms (ADS) periods.

Variable	PM _{2.5}	SO ₂	O ₃	CO	NO ₂	NO	NO _x	ADT	DTR	RH
PM ₁₀	0.76 **	0.51 **	0.01	0.34 *	-0.03	-0.11	-0.06	0.10	0.11	-0.34 *
PM _{2.5}		0.45 *	-0.04	0.54 **	-0.07	-0.15	-0.10	0.22	-0.15	-0.02
SO ₂			-0.32 *	0.55 **	0.22	0.36 *	0.28 *	-0.31 *	0.05	-0.48 **
O ₃				-0.63 **	-0.75 **	-0.69 **	-0.74 **	-0.09	0.16	-0.20
CO					0.57 **	0.46 *	0.55 **	0.38 *	-0.09	0.28 *
NO ₂						0.92 **	0.99 **	0.32 *	0.28 *	0.16
NO							0.97 **	0.05	0.31 *	-0.02
NO _x								0.23	0.30 *	0.10
ADT									-0.12	0.66 **
DTR										-0.48 **

* $p < 0.05$, ** $p < 0.0001$.

Table 3 shows the effects of PM_{2.5} on ER visits for the susceptible groups with CVDs and RDs. For those 65 years of age, a 10 $\mu\text{g}/\text{m}^3$ increase in PM_{2.5} led to a 2.77-fold ($p < 0.05$) increase in the risk of ER visits for those with CVDs. Likewise, we found a 3.94-fold increase in the risk of ER visits for those with RDs. For those <65 years of age, a 10 $\mu\text{g}/\text{m}^3$ increase in PM_{2.5} led to OR of 1.77 ($p < 0.05$) for the risk of ER visits for RDs. A 10 $\mu\text{g}/\text{m}^3$ increase in PM_{2.5} for men, OR was 1.80 ($p < 0.05$) for the risk of ER visits for RDs, and OR was 1.99 for the risk of ER visits for CVDs. We found significant effects on women whose ORs were 3.09 and 1.86 for ER visits for CVDs and RDs, respectively.

Table 3. A 10 $\mu\text{g}/\text{m}^3$ increase in PM_{2.5} during ADS correlated with cardiovascular and respiratory diseases in emergency room (ER) based on age and sex.

Variable	Cardiovascular Diseases			Respiratory Diseases		
	OR	95% CI		OR	95% CI	
≥ 65	2.77 *	1.01	26.69	3.94	0.75	NA
<65	1.92	0.95	5.31	1.77 *	1.24	2.79
Male	1.99	0.80	8.77	1.80 *	1.13	3.58
Female	3.09 *	1.15	26.80	1.86 *	1.15	3.45

* $p < 0.05$.

Table 4 shows the lag effect of PM_{2.5} on CVD and RD emergency visits. During the first and second day of the sand-dust storm arriving in Taiwan (lag 0 and lag 1, respectively), the increase in PM_{2.5} seems to have led to an increase in the ER visits for CVDs. As for the exposure period's effect on ER visits for RDs, every day (lag 0 to lag 5) besides lag 3 had a significant effect.

Table 4. Lag effects for PM_{2.5} during ADS correlated with cardiovascular diseases (CVDs) and respiratory diseases (RDs) in the single-pollutant model.

Lag	Cardiovascular Diseases			Respiratory Diseases		
	OR	95% CI		OR	95% CI	
Lag 0	2.20 *	1.22	5.08	1.86 *	1.30	2.91
Lag 1	1.77 *	1.05	3.97	1.48 *	1.03	2.31
Lag 2	1.86	0.78	5.94	3.78 *	1.84	10.55
Lag 3	0.91	0.652	1.23	1.24	0.95	1.62
Lag 4	0.82	0.60	1.11	1.20 *	1.01	1.41
Lag 5	1.79	0.94	3.43	1.54 *	1.14	2.07

* $p < 0.05$.

Table 5 shows the three models' (single-, two- and multi-pollutant) descriptions of the effects of the heightened PM_{2.5} during the exposure period on ER visits for CVDs and RDs. In the single-pollutant model, which does not account for other pollutants, a 10 µg/m³ increase in PM_{2.5} levels led to an OR of 2.20 (95% CI: 1.22–5.08) for ER visits for CVDs and an OR of 1.86 (95% CI: 1.30–2.91) for ER visits for RDs. For the two-pollutant model, the models that added NO₂, CO, NO or NO_x showed a significant correlation between PM_{2.5} and ER visits for CVDs; models that added SO₂, NO₂, O₃, CO, NO or NO_x showed a significant correlation between PM_{2.5} and ER visits for RDs. In the multi-pollutant model, PM_{2.5} showed no statistical correlation with ER visits for CVDs, but we found statistical significant correlations with RDs in the models including SO₂ + NO₂, SO₂ + CO₂, SO₂ + NO_x, NO₂ + CO, O₃ + NO_x and CO + NO_x.

Table 5. Odds ratio (OR) of disease associated with 10 µg/m³ increase of PM_{2.5} with single- and multiple-pollutant models.

Pollutant	Cardiovascular Diseases			Respiratory Diseases		
	OR	95% CI		OR	95% CI	
Single pollutants models						
PM _{2.5}	2.20 *	1.22	5.08	1.86 *	1.30	2.91
Two pollutant model						
+SO ₂	7.29	0.95	55.96	2.66 *	1.18	6.00
+NO ₂	2.24 *	1.06	4.74	2.20 *	1.35	3.59
+O ₃	1.70	0.86	3.36	1.60 *	1.03	2.48
+CO	4.39 *	1.10	17.55	5.43 *	1.58	18.64
+NO	3.15 *	1.16	8.57	2.97 *	1.12	7.90
+NO _x	2.60 *	1.18	5.77	2.51 *	1.33	4.76
Three pollutant model						
+SO ₂ + NO ₂	9.24	0.91	94.27	2.52 *	1.26	5.03
+SO ₂ + O ₃	4.08	0.47	35.03	1.74	0.91	3.32
+SO ₂ + CO	7.32	0.88	61.18	5.24 *	1.45	18.94
+SO ₂ + NO	5.59	0.41	76.68	2.89	0.74	11.32
+SO ₂ + NO _x	5.94	0.76	46.29	2.51 *	1.12	5.65
+NO ₂ + O ₃	NA	NA	NA	1.63	0.96	2.78
+NO ₂ + CO	11.16	0.56	222.27	4.73 *	1.54	14.52
+NO ₂ + NO	NA	NA	NA	3.61	0.98	13.35
+NO ₂ + NO _x	NA	NA	NA	3.59	0.96	13.34
+O ₃ + CO	2.51	0.34	18.57	4.46	0.95	20.87

Table 5. Cont.

Pollutant	Cardiovascular Diseases			Respiratory Diseases		
	OR	95% CI		OR	95% CI	
Three pollutant model						
+O ₃ + NO	3.07	0.53	17.65	NA	NA	NA
+O ₃ + NO _x	1.32	0.59	2.97	2.96 *	1.13	7.78
+CO + NO	2.42	0.48	12.20	2.27	0.84	6.17
+CO + NO _x	6.53	0.88	48.34	3.50 *	1.26	9.73
+NO + NO _x	NA	NA	NA	3.62	0.98	13.36

* $p < 0.05$, NA: non-available.

4. Discussion

The desertification of northern China has worsened in recent years, which, when combined with the global increase of drought and uneven distribution of rainfall, has led to a rise in both the frequency and severity of Asian Dust Storms. Moreover, ADSs differ from other dust storms: First, the particulates of ADS reaching Taiwan are very small. This is because the long distance the ADS must travel prevents the transportation of larger particles; gravity eventually brings those larger particles back to the ground or sea. As a result, the particulate size ranges from 1.35~10 μm . Second, the average concentration of air pollutants during ADS is significantly high mainly because ADS form as clouds of dust, not as high-speed winds that carry dust with them. For this reason, ADS tend to carry with them a larger amount of toxins [23,24]. Because the voyage of ADS is affected by dissipation, subsidence, and filtering due to rainfall, the Taiwanese EPA uses meteorological observatory data and other countries' dust storm models to track ADS, confirming whether they will impact Taiwan. After a detailed analysis that predicts a decrease in Taiwan's air quality as a result of the ADSs, the EPA will issue a warning to the citizens of Taiwan, encouraging them to take preventative measures against the negative effects of ADS. The observation of the ADS continues until it leaves Taiwan. An ADS increases the concentrations of PM_{2.5} in central Taiwan. During an ADS, the PM_{2.5} level is 61.93 $\mu\text{g}/\text{m}^3$, while that on comparison days is 48.87 $\mu\text{g}/\text{m}^3$. Our findings are the same as those of Kamouchi et al. [28] and Kwon et al. [29]. Yet the levels of combustion particulates (CO and NO₂) do not differ during an ADS, a result that matches that of Yang et al. [30].

When attempting to understand the effects of air pollutants on daily emergency visits, deaths, and hospital visits, we must also control for seasonal and other time variables. Our study design employed a unidirectional recall case-crossover methodology instead of a bidirectional case-crossover design. Using the time point of 14 days prior to the sand-dust storm as our control, we were able to control confounders for the long-term and for different season. In addition, by having looked to the past, we could control for the risks of physical pain that occur after emergency visits and exposure to PM_{2.5}. Through this case-crossover design, we were able to control for many confounders without having to rely on statistical models. However, some research has shown that unidirectional case-crossover study designs can overestimate odds ratios. Because ER visits and air pollution are both susceptible to time trends, when looking at exposure intensity, the control period could be systematically higher or lower than the exposure period [15].

Our study makes a contribution to research on the correlation between PM_{2.5} from ADS and CVDs from RDs ER visits. Compared to other studies, our study found a larger effect when PM_{2.5} levels increased every 10 $\mu\text{g}/\text{m}^3$. One possibility for this is that the PM_{2.5} of our research could have a different chemical composition [23]. Principal component analysis (PCA) was used to analyze the components of Taiwanese ADS and found three principle components were able to explain 88% of the variation. The first component (PC1) explained 52% of the variation and represented a larger-than-average concentration of NO and NO_x (factor loading >0.70). These components mainly came from industry and traffic exhaust. The second component (PC2) explained 24% of the variation

and consisted of $PM_{2.5}$, SO_4^{2-} , NO_3^- , OC, as well as highly dense mineral components and pollutions such as sulfate and nitrate. These components mainly came from Chinese factories [31]. The main component in the ADSs of 2002 in Taiwan was sulfate, as ADSs do not only carry dust but also combustion by-products [32].

Our study shows that, according to the single-pollutant model (i.e., a model that does not adjust for the effects of other pollutants), a $10 \mu\text{g}/\text{m}^3$ increase in $PM_{2.5}$ corresponds with statistical significance to an increase in CVD and RD emergency room visits. For the two-pollutant models that adjusted for NO_2 , CO, NO and NO_x , a $10 \mu\text{g}/\text{m}^3$ increase in $PM_{2.5}$ led to an increase in ER visits for CVDs in these models that controlled for CO showed the largest effect. In the two-pollutant models adjusted for SO_2 , NO_2 , O_3 , CO, NO and NO_x , a $10 \mu\text{g}/\text{m}^3$ increase in $PM_{2.5}$ led to more RD emergency visits. Again, that controlling for CO showed the largest effect. Other research [33] in Taiwan has pointed out that despite other air pollutants having an effect on ER visits (e.g., NO_2 , CO and O_3 increasing visits for RDs as well as CO, NO_2 and SO_2 increasing ER visits for heart diseases), the relative risks of these pollutants is less than that of $PM_{2.5}$. In multi-pollutant models, adjusting for CO and NO_2 still results in the relative risk (RR) for $PM_{2.5}$ being higher. We created cut-off points for the groups most easily affected by $PM_{2.5}$ and discovered that people over the age of 65 or women were more likely to ER visits during the extra $PM_{2.5}$ exposure that ADSs brought. Seniors typically have higher ER visits for CVD, which leads to a higher susceptibility to $PM_{2.5}$. Other research [17] has supported this fact by showing that those under the age of 65 are less likely than their older counterparts to be hospitalized for CVDs during $PM_{2.5}$ exposure. Women might be more susceptible because of hormones, but research on the interaction between sex and air pollution has not yet proved such a thing [34]. ER visits for respiratory problems occurs more often in the <65 years of age, and sex differences in this case do not differ significantly in the OR. Nevertheless, regardless of sex, increased $PM_{2.5}$ exposure increases the risk of respiratory problems. Other research [14] on why those under the age of 65 are more likely to treatment for respiratory problems during an ADS has investigated outside activity time and immune system mechanisms, citing them as probable factors.

Evidence has shown that exposure to air pollution can have an adverse effect on the protective abilities of the lungs, such as the dynamic filtering of air, mucociliary cleaning, particulate dispersion, and alveolar macrophage expulsion. Macrophage can inhibit the propagation of viruses, destroying fragmented cells and restricting the transmission of the virus through the production of antigens for T lymph cells. Exposure to $PM_{2.5}$ can lead to an inflammatory reaction in the lungs, exacerbating lung diseases. It can also release harmful enzymes in the cell as well as alter other reactions related to blood coagulation and its triggering. Consequences of this include cardiovascular problems that require urgent care and the development of atherosclerosis [35]. Furthermore, air particulate pollution significantly correlated with autonomic function measured by changes in heart rate variability (HRV) and blood markers of inflammation. The result is an increase in heart rate and the variance of that speed [13]. Because $PM_{2.5}$ acts on different disease mechanisms for CVDs and RDs, we used a lag effect to investigate the differences in timing when ADS's $PM_{2.5}$ levels increases affects these two diseases. As $PM_{2.5}$ levels increases led to more emergency visits for CVDs on the day of the ADS (lag 0) and on the day after (lag 1), we conclude that exposure to $PM_{2.5}$ during a ADS has a more urgent effect on CVDs, a result that matches that of Kamouchi et al. [25]. Exposure to $PM_{2.5}$ brought by ADS affects RDs over the course of all five days, with only lag 3 not having statistical significance. This result shows that the exposure to $PM_{2.5}$ from ADS does not have such an immediate effect on RDs as it does on CVDs. The reason for this is that the $PM_{2.5}$ in ADS differs from microorganisms. A microorganism must attach to the sand and dust, then enter the body, where they can finally do harm to the respiratory system [25].

Our study has some limitations. First, we used outdoor monitoring stations to collect the average daily air pollution levels and other weather variables as our representatives for human exposure to air pollutants. However, other factors—such as the use of air conditioning and the ventilation between indoors and outdoors—might influence how representative our sample truly is, thereby affecting

our results and their implications [34]. Second, our study from NHI data did not include individual behavioral data. Other studies have shown how heavy smokers or drinkers are more susceptible to the negative health effects of ADSs [25]. Third, we assumed that PM_{2.5} to be located in specific areas and that our designated areas were homogenous in airborne quality. We determined that these areas can represent the average air quality situation in central Taiwan as pertains to PM_{2.5}. However, we were without data on the components of PM_{2.5} and therefore could not establish exactly what part of PM_{2.5} is responsible for the negative health effects we found. Hence, during an ADS, the concentrations and types of suspended fine particles could have different levels of negative effects on health. From a study in Beijing [36], the PM_{2.5} concentration was approximately 230 µg/m³ and the crustal elements constituted about 66.4% of the chemical composition of PM_{2.5} while sulfate and nitrate contributed much less during the ADS period. We therefore can only suggest that susceptible groups stay indoors during an ADS. However, the doors and windows of a household can only offer limited protection, and the quality as well as the maintenance of the air conditioning systems might be lacking. PM_{2.5} can easily enter households, carrying along with them pollutants [37]. Thus, environmental protection organizations should not only implement measures to offer earlier warnings of ADSs but should also advise at-risk groups (asthmatics, those with CVD or RD, people with allergies—and especially the elderly as well as young children) to remain indoors. These organizations should also emphasize that people check the sealing quality on door and screens.

5. Conclusions

This study has shown that exposure to high levels of PM_{2.5} as brought by ADSs increases the risk of ER visits for CVDs and RDs, especially in people above the age of 65 or women. However, those under the age of 65 are also affected by the increase in PM_{2.5} brought by ADSs in that they are more likely to require ER visits for RDs. For this reason, Asian nations should track the ADSs arriving from China as part of a precautionary system that can ensure their citizens' health.

Acknowledgments: We thank Taiwan's Environmental Protection Administration provide air pollution online Database. Grant of project was supported from the Ministry of Science and Technology (Code: 105-2314-B-010-018-).

Author Contributions: Ssu-Ting Liu participated in the design of the study, data check, and performed the statistical analysis. Chu-Yung Liao took study designing and data collection. Cheng-Yu Kuo took related literature review, English editing and revision, and write first draft. Hsien-Wen Kuo participated in reviewing article and idea generation the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

Abbreviations

ADS	Asian dust storm
EPA	Taiwan Environmental Protection Administration
TEOM	Tapered Element Oscillating Microbalance
CI	Confidence interval
ER	Emergency room
ICD-9	International Classification of Diseases, 9th Revision
NO ₂	Nitrogen dioxide
NO _x	Nitrogen oxides
PM _{2.5}	Fine particulate matters with an aerodynamic diameter of less than 2.5 µm
SO ₂	Sulfur dioxide
CO	Carbon monoxide
O ₃	Ozone
ADT	Average daily temperature
DTR	Diurnal temperature range
RH	Relative humidity
CVD	Cardiovascular diseases
RD	Respiratory diseases
WHO	World Health organization
NHI	National Health Insurance
OR	Odds ratios

References

1. Taiwan EPA. *Environmental Protection Annual*; Environmental Protection Agency: Taipei, Taiwan, 2014.
2. Chen, Y.S.; Sheen, P.C.; Chen, E.R.; Liu, Y.K.; Wu, T.N.; Yang, C.Y. Effects of Asian Dust Storm Events on Daily Mortality in Taipei, Taiwan. *Environ. Res.* **2004**, *95*, 151–155. [[CrossRef](#)] [[PubMed](#)]
3. Yang, C.Y.; Chang, C.C.; Chuang, H.Y.; Tsai, S.S.; Wu, T.N.; Ho, C.K. Relationship between air pollution and daily mortality in a subtropical city: Taipei, Taiwan. *Environ. Int.* **2004**, *30*, 519–523. [[CrossRef](#)] [[PubMed](#)]
4. Milojevic, A.; Wilkinson, P.; Armstrong, B.; Bhaskaran, K.; Smeeth, L.; Hajat, S. Short-term effects of air pollution on a range of cardiovascular events in England and Wales: Case-crossover analysis of the MINAP database, hospital admissions and mortality. *Heart* **2014**, *100*, 1093–1098. [[CrossRef](#)] [[PubMed](#)]
5. Son, J.Y.; Cho, Y.S.; Lee, J.T. Effects of air pollution on postneonatal infant mortality among firstborn infants in Seoul, Korea: Case-crossover and time-series analyses. *Arch. Environ. Occup. Health* **2008**, *63*, 108–113. [[CrossRef](#)] [[PubMed](#)]
6. Tsai, S.S.; Chang, C.C.; Liou, S.H.; Yang, C.Y. The effects of fine particulate air pollution on daily mortality: A case-crossover study in a subtropical city, Taipei, Taiwan. *Int. J. Environ. Res. Public Health* **2014**, *11*, 5081–5093. [[CrossRef](#)] [[PubMed](#)]
7. Ueda, K.; Nitta, H.; Ono, M.; Takeuchi, A. Estimating mortality effects of fine particulate matter in Japan: A comparison of time-series and case-crossover analyses. *J. Air Waste Manag. Assoc.* **2009**, *59*, 1212–1218. [[CrossRef](#)] [[PubMed](#)]
8. Samoli, E.; Stafoggia, M.; Rodopoulou, S.; Ostro, B.; Declercq, C.; Alessandrini, E.; Diaz, J.; Karanasiou, A.; Kelessis, A.G.; Le Tertre, A.; et al. Associations between fine and coarse particles and mortality in Mediterranean cities: Results from the MED-PARTICLES project. *Environ. Health Perspect.* **2013**, *121*, 932–938. [[CrossRef](#)] [[PubMed](#)]
9. Neuberger, M.; Moshhammer, H.; Rabczenko, D. Acute and subacute effects of urban air pollution on cardiopulmonary emergencies and mortality: Time series studies in Austrian cities. *Int. J. Environ. Res. Public Health* **2013**, *10*, 4728–4751. [[CrossRef](#)] [[PubMed](#)]
10. Ostro, B.; Tobias, A.; Querol, X.; Alastuey, A.; Amato, F.; Pey, J.; Perez, N.; Sunyer, J. The effects of particulate matter sources on daily mortality: A case-crossover study of Barcelona, Spain. *Environ. Health Perspect.* **2011**, *119*, 1781–1787. [[CrossRef](#)] [[PubMed](#)]
11. Dominici, F.; Peng, R.D.; Bell, M.L.; Pham, L.; McDermott, A.; Zeger, S.L.; Samet, J.M. Fine particulate air pollution and hospital admission for cardiovascular and respiratory diseases. *JAMA* **2006**, *295*, 1127–1134. [[CrossRef](#)] [[PubMed](#)]
12. Zanobetti, A.; Schwartz, J. The effect of particulate air pollution on emergency admissions for myocardial infarction: A multicity case-crossover analysis. *Environ. Health Perspect.* **2005**, *113*, 978–982. [[CrossRef](#)] [[PubMed](#)]
13. Pope, C.A., III; Hansen, M.L.; Long, R.W.; Nielsen, K.R.; Eatough, N.L.; Wilson, W.E.; Eatough, D.J. Ambient particulate air pollution, heart rate variability, and blood markers of inflammation in a panel of elderly subjects. *Environ. Health Perspect.* **2004**, *112*, 339–345. [[CrossRef](#)] [[PubMed](#)]
14. Ueda, K.; Nitta, H.; Odajima, H. The effects of weather, air pollutants, and Asian dust on hospitalization for asthma in Fukuoka. *Environ. Health Prev. Med.* **2010**, *15*, 350–357. [[CrossRef](#)] [[PubMed](#)]
15. Kanatani, K.T.; Ito, I.; Al-Delaimy, W.K.; Adachi, Y.; Mathews, W.C.; Ramsdell, J.W. Desert dust exposure is associated with increased risk of asthma hospitalization in children. *Am. J. Respir. Crit. Care Med.* **2010**, *182*, 1475–1481. [[CrossRef](#)] [[PubMed](#)]
16. Lin, M.; Chen, Y.; Burnett, R.T.; Villeneuve, P.J.; Krewski, D. The influence of ambient coarse particulate matter on asthma hospitalization in children: Case-crossover and time-series analyses. *Environ. Health Perspect.* **2002**, *110*, 575–581. [[CrossRef](#)] [[PubMed](#)]
17. Lin, M.; Stieb, D.M.; Chen, Y. Coarse particulate matter and hospitalization for respiratory infections in children younger than 15 years in Toronto: A case-crossover analysis. *Pediatrics* **2005**, *116*, e235–e240. [[CrossRef](#)] [[PubMed](#)]
18. Haley, V.B.; Talbot, T.O.; Felton, H.D. Surveillance of the short-term impact of fine particle air pollution on cardiovascular disease hospitalizations in New York State. *Environ. Health* **2009**, *8*, 42. [[CrossRef](#)] [[PubMed](#)]

19. Cadelis, G.; Tourres, R.; Molinie, J. Short-term effects of the particulate pollutants contained in saharan dust on the visits of children to the emergency department due to asthmatic conditions in Guadeloupe (French Archipelago of the Caribbean). *PLoS ONE* **2014**, *9*, e91136. [[CrossRef](#)] [[PubMed](#)]
20. Rosenthal, F.S.; Carney, J.P.; Olinger, M.L. Out-of-hospital cardiac arrest and airborne fine particulate matter: A case-crossover analysis of emergency medical services data in Indianapolis, Indiana. *Environ. Health Perspect.* **2008**, *11*, 631–636. [[CrossRef](#)] [[PubMed](#)]
21. Santus, P.; Russo, A.; Madonini, E.; Allegra, L.; Blasi, F.; Centanni, S.; Miadonna, A.; Schiraldi, G.; Amaducci, S. How air pollution influences clinical management of respiratory diseases. A case-crossover study in Milan. *Respir. Res.* **2012**, *13*, 95. [[PubMed](#)]
22. Lee, J.T.; Son, J.Y.; Cho, Y.S. A comparison of mortality related to urban air particles between periods with Asian dust days and without Asian dust days in Seoul, Korea, 2000–2004. *Environ. Res.* **2007**, *105*, 409–413. [[CrossRef](#)] [[PubMed](#)]
23. Chen, S.Y.; Lin, Y.L.; Chang, W.T.; Lee, C.T.; Chan, C.C. Increasing emergency room visits for stroke by elevated levels of fine particulate constituents. *Sci. Total Environ.* **2014**, *473–474*, 446–450.
24. Krzyzanowski, M.; Cohen, A. Update of WHO air quality guidelines. *Air Qual. Atmos. Health* **2008**, *1*, 7–13. [[CrossRef](#)]
25. Newby, D.E.; Mannucci, P.M.; Tell, G.S.; Baccarelli, A.A.; Brook, R.D.; Donaldson, K.; Forastiere, F.; Franchini, M.; Franco, O.H.; Graham, I.; et al. Expert position paper on air pollution and cardiovascular disease. *Eur. Heart J.* **2015**, *36*, 83–93. [[CrossRef](#)] [[PubMed](#)]
26. Franklin, M.; Koutrakis, P.; Schwartz, J. The role of particle composition on the association between PM_{2.5} and mortality. *Epidemiology* **2008**, *19*, 680–689. [[PubMed](#)]
27. Maclure, M. The Case-Crossover Design: A Method for Studying Transient Effects on the Risk of Acute Events. *Am. J. Epidemiol.* **1991**, *133*, 144–153. [[CrossRef](#)] [[PubMed](#)]
28. Kamouchi, M.; Ueda, K.; Ago, T.; Nitta, H.; Kitazono, T. Relationship between asian dust and ischemic stroke: A time-stratified case-crossover study. *Stroke* **2012**, *43*, 3085–3087. [[CrossRef](#)] [[PubMed](#)]
29. Kwon, H.J.; Cho, S.H.; Chun, Y.; Lagarde, F.; Pershagen, G. Effects of the Asian Dust Events on Daily Mortality in Seoul, Korea. *Environ. Res.* **2002**, *90*, 1–5. [[CrossRef](#)] [[PubMed](#)]
30. Yang, C.Y.; Chen, Y.S.; Chiu, H.F.; Goggins, W.B. Effects of Asian dust storm events on daily stroke admissions in Taipei, Taiwan. *Environ. Res.* **2005**, *99*, 79–84. [[CrossRef](#)] [[PubMed](#)]
31. Liang, C.S.; Yu, T.Y.; Chang, Y.Y.; Syu, J.Y.; Lin, W.Y. Source Apportionment of PM_{2.5} Particle Composition and Submicrometer Size Distribution during an Asian Dust Storm and Non-Dust Storm in Taipei. *Aerosol Air Qual. Res.* **2013**, *13*, 545–554.
32. Lee, C.T.; Chuang, M.T.; Chan, C.C.; Cheng, T.J.; Huang, S.L. Aerosol characteristics from the Taiwan aerosol supersite in the Asian yellow-dust periods of 2002. *Atmos. Environ.* **2006**, *40*, 3409–3418. [[CrossRef](#)]
33. Hwang, J.S.; Hu, T.H.; Chan, C.C. Air Pollution Mix and Emergency Room Visits for Respiratory and Cardiac Diseases in Taipei. *J. Data Sci.* **2004**, *2*, 311–327.
34. Yang, C.; Peng, X.; Huang, W.; Chen, R.; Xu, Z.; Chen, B.; Kan, H. A time-stratified case-crossover study of fine particulate matter air pollution and mortality in Guangzhou, China. *Int. Arch. Occup. Environ. Health* **2012**, *85*, 579–585. [[CrossRef](#)] [[PubMed](#)]
35. Terzano, C.; Stefano, F.D.; Conti, E.; Graziani, C.E.; Petroianni, A. Air pollution ultrafine particles: Toxicity beyond the lung. *Eur. Rev. Med. Pharmacol. Sci.* **2010**, *14*, 809–821. [[PubMed](#)]
36. Liang, Y.; Fang, L.; Pan, H.; Zhang, K.; Kan, H.; Brook, J.R.; Sun, Q. PM_{2.5} in Beijing-temporal pattern and its association with influenza. *Environ. Health* **2014**, *13*, 102. [[PubMed](#)]
37. Kuo, H.W.; Shen, H.Y. Indoor and outdoor PM_{2.5} and PM₁₀ concentrations in the air during a dust storm. *Build. Environ.* **2010**, *45*, 610–614.

