

Table S1. Distribution of characteristics of all pharmacists finally analyzed

| Characteristics | Community pharmacist (<i>n</i> = 463) |
|---|---|
| Age group | |
| ≤34 years | 57 (12.3%) |
| 35–44 years | 147 (31.7%) |
| 45–54 years | 154 (33.3%) |
| ≥55 years | 105 (22.7%) |
| Sex | |
| Female | 360 (77.8%) |
| Male | 103 (22.2%) |
| Professional category | |
| Titular pharmacist | 280 (60.5%) |
| Assistant pharmacist | 58 (12.5%) |
| Substitute pharmacist | 125 (27.0%) |
| Years of work | |
| ≤ 9 years | 57 (12.3%) |
| 10–29 years | 310 (67.0%) |
| ≥ 30 years | 96 (20.7%) |
| Type of population | |
| Rural | 120 (25.9%) |
| Urban | 324 (70.0%) |
| Seasonal vaccination 2013/14 | |
| Yes | 116 (25.1%) |
| Seasonal vaccination in any of the three preceding seasons | |
| Yes | 167 (36.1%) |
| Seasonal vaccination in all three preceding seasons | |
| Yes | 111 (24.0%) |
| Vaccination with pandemic vaccine 2009-2010 | |
| Yes | 167 (36.1%) |
| I recommend the vaccine to pregnant women in their first trimester | |
| Yes | 138 (29.8%) |
| I recommend the vaccine to pregnant women in their second or third trimester | |
| Yes | 204 (44.1%) |
| I recommend the vaccine to post-partum women | |
| Yes | 143 (30.9%) |
| I recommend the vaccine to people with chronic disorders | |
| Yes | 439 (94.8%) |
| I recommend the vaccine to immunosuppressed people | |
| Yes | 295 (63.7%) |
| I recommend the vaccine to persons aged > 65 years | |
| Yes | 440 (95.0%) |