

SUPPLEMENTAL MATERIAL

Determination and prediction of respirable dust and crystalline-free silica in the Taiwanese foundry industry

Ching-Tang Kuo¹, Fen-Fen Chiu², Bo-Ying Bao^{3,4}, Ta-Yuan Chang^{2,*}

¹ Department of Public Health, College of Public Health, China Medical University, No.

91, Hsueh-Shih Road, Taichung 40402, Taiwan; ctkuo228@gmail.com

² Department of Occupational Safety and Health, College of Public Health, China

Medical University, No. 91, Hsueh-Shih Road, Taichung 40402, Taiwan;

jessica.cy@msa.hinet.net(F.-F.C.); tychang@mail.cmu.edu.tw(T.-Y.C.)

³ Department of Pharmacy, College of Pharmacy, China Medical University,

Taichung, Taiwan. Department of Pharmacy, College of Pharmacy, China Medical

University, No. 91, Hsueh-Shih Road, Taichung 40402, Taiwan; bao@mail.cmu.edu.tw

⁴ Department of Nursing, Asia University, No. 500, Lioufeng Road, Wufeng, Taichung

41354, Taiwan

* Correspondence: tychang@mail.cmu.edu.tw; Tel.: +886-4-22053366 ext. 6203. Fax:

+886-4-22079225.

Figure of contents:	Page
Figure S 1. Residual diagnostics of the predictive regression model for base-10 logarithmically translated levels of respirable dust.....	3
Figure S 2. Residual diagnostics of the predictive regression model for base-10 logarithmically translated levels of respirable crystalline silica	4

Figure S 1. Residual diagnostics of the predictive regression model for base-10 logarithmically translated levels of respirable dust




Figure S 2. Residual diagnostics of the predictive regression model for base-10 logarithmically translated levels of respirable crystalline silica

