



Article

Association of Anthropometric Indices of Obesity with Hypertension in Chinese Elderly: An Analysis of Age and Gender Differences

Qian Wang ¹, Lingzhong Xu ^{1,2,*}, Jiajia Li ^{1,2}, Long Sun ^{1,2}, Wenzhe Qin ¹, Gan Ding ¹, Jing Zhu ¹, Jiao Zhang ¹, Zihang Yu ¹ and Su Xie ¹

¹ School of Public Health, Shandong University, Jinan 250012, China; wangqian0519@126.com (Q.W.); lijiajia@sdu.edu.cn (J.L.); sunlong@sdu.edu.cn (L.S.); qinwenzhe09@163.com (W.Q.); dinggan90@163.com (G.D.); zj1176559133@163.com (Jin.Z.); 201614324@mail.sdu.edu.cn (Jia.Z.); 13249220350@163.com (Z.Y.); xs340823@163.com (S.X.)

² Collaborative Innovation Center of Social Risks Governance in Health, School of Public Health, Fudan University, Shanghai 200032, China

* Correspondence: lzxu@sdu.edu.cn; Tel.: +86-0531-8838-2648

Received: 14 March 2018; Accepted: 17 April 2018; Published: 19 April 2018



Abstract: This study aims to explore the association of anthropometric indices of obesity with hypertension in Chinese elderly and its possible gender and age differences. A total of 7070 adults age 60 or older were interviewed in a cross-sectional study conducted in 2017. Anthropometric indices for each participant were measured by using standard methods of trained doctoral/master students. We performed two binary logistic regression models to examine the association of the nine different anthropometric indices and hypertension by gender. Lastly, analyses were performed in two steps stratified for age. Comparing individuals with and without hypertension, there were statistically significant differences in anthropometric indices except height, a body shape index (ABSI), and hip index (HI) in males; and except height in females. There were gender differences in the relationship between anthropometric indices and the prevalence of hypertension in Chinese older adults. After stratification by age, the associations of all anthropometric indices became weaker, disappeared, or even went in the opposite direction. Furthermore, body mass index (BMI) in men (except individuals older than 80) and hip circumference (HC) in women showed a significant impact on the risk of hypertension. The association of anthropometric indices of obesity with hypertension in Chinese elderly differ by gender and age. These findings indicate a need to develop gender-specific strategies for the male and female elderly in the primary and secondary prevention of hypertension.

Keywords: hypertension; anthropometric indices; obesity; elderly; China

1. Introduction

Hypertension is the leading global risk factor of cardiovascular disease (CVD). It contributes to nearly 9.4 million deaths worldwide annually [1,2]. In China, hypertension is common and its prevalence is rising. Every year, almost 2.1 million cardiovascular deaths and 1.2 million premature cardiovascular deaths are attributed to hypertension [3]. A recent study showed that nearly half of Chinese adults ages 35–75 have hypertension. Among these hypertensive patients, the elderly—aged 60 or older—accounted for 50.1% higher than the sum of other age groups [4].

Obesity is increasing globally and has received widespread attention [5]. Data from the World Health Organization estimated that more than 1.9 billion adults were overweight in 2016, and that more than 650 million were obese [6]. Many studies have suggested that the high incidence of hypertension is

associated with the dramatic increase in the prevalence of obesity [7–10]. Other studies have identified several mechanisms as potential causes of obesity-hypertension to verify this association [11–13].

Obesity can be defined by different anthropometric measurements and indices. In recent years, there has been increasing speculation over which measure of obesity predicts CVD better [14]. However, previous studies on this remain controversial. Body mass index (BMI) is commonly used in many epidemiologic studies on obesity. However, it is reported to be incapable of differentiating body fat and lean body mass in many studies [15–18]. Some studies found statistical evidence that supports the superiority of measures of centralized obesity (such as waist circumference (WC), waist-to-hip ratio (WHR), or waist-to-height ratio (WHtR)) over BMI, for detecting cardiovascular risk factors in both men and women [14,19–23]. Others found that BMI and central obesity indicators were equally well regardless of age and sex [24]. Moreover, many new anthropometric indices were developed recently, such as a new hip index (HI) and a body shape index (ABSI), and these measures proved to be complementary to BMI and other known risk factors [25,26]. Therefore, in our current study, we included more indices (such as height, weight, WC, hip circumference (HC), WHR, WHtR, BMI, HI, and ABSI) than previous studies to make our results more comprehensive.

Aging population has become a worldwide phenomenon. According to China National Bureau of Statistics, the population age 60 and above accounted for 16.2% of the total population at the end of 2015 [27]. Furthermore, hypertension mostly affects elderly people, these individuals are more likely to have organ damages or clinical CVD [4]. However, the majority of studies on the association between anthropometric indices and hypertension to date have targeted general adults rather than the elderly [15–18,20–24]. Thus, studies focused on these vulnerable population are more necessary.

Although there have been some studies aimed to explore which anthropometric indicators are more strongly associated with hypertension, controversies exist within the results. For example, some studies reported that general obesity (BMI) rather than abdominal obesity linked with hypertension directly [15]; some studies found the superiority of abdominal obesity (such as WC, WHtR) in detecting hypertension [14,19]; and others indicated that these indicators were equally strongly associated with hypertension [24]. In China, many studies explored the association between anthropometric indicators and hypertension from a different gender perspective and tried to find out the better predictor for hypertension in males and females [28,29]. However, few studies stratified for age at the same time, which may significantly influence the association and its gender differences. Therefore, in this study we aim to: (1) explore the association of anthropometric indices of obesity with hypertension among Chinese elderly; (2) find out its possible gender and age differences.

2. Materials and Methods

2.1. Study Participants and Data Collection

This study was collected from the 2017 Survey of the Shandong Elderly Family Health Service, which was conducted by Shandong University. Stratified multi-stage random sampling was applied: in the first stage, six counties were selected from 137 counties as the primary sampling units (PSUs) throughout the eastern, central and western regions of Shandong province (which were divided into three districts and three counties that represented urban and rural areas separately). From each PSU, 18 villages in rural area and 18 communities in suburban and urban area were selected as the secondary sampling units (SSUs). In the third stage, based on the roster of the residents by age and total elderly population of each selected site provided by the local residential committee, an average of 66 individuals were stratified and randomly selected from each SSU making up the total sample. The eligible participants for this survey were those age 60 or older with local household registrations at the time of the interview. Initially, 7088 elderly individuals were selected and interviewed. Of these, 18 did not complete the survey. Finally, a total of 7070 individuals were included in the sample (Figure 1).

All data were collected in the participant's home using a study questionnaire by trained doctoral/master students. Informed consent for the collection and use of information was obtained from

all participants. Medical ethics approval was not required since particular treatments or interventions were not offered or withheld from respondents as a consequence of participating in the study.




Figure 1. Flow chart of participant enrolment.

2.2. Dependent Variable

The elderly with hypertension were self-reported, which derived from a question “Have you been diagnosed with hypertension by a doctor”. If they said “yes,” our investigators (master/doctor students with medical background) checked further. The inclusion criteria were: (i) have been diagnosed with hypertension by doctor; (ii) need to be on long-term medication. Temporary increases of blood pressure were excluded carefully.

2.3. Anthropometric Measurements

To obtain a more reliable result, our study included most of the traditional anthropometric indices (height, weight, WC, HC, WHR, WHtR, BMI) and some newly developed anthropometric indices (HI and ABSI).

Height without shoes was measured in centimeters (accuracy 1.0 cm), and weight in light clothing was measured in kilograms (accuracy 0.01 kg). Height was classified into quartiles: Quartile 1 (<154), Quartile 2 (154–160), Quartile 3 (160–165), and Quartile 4 (≥ 165). Weight was also classified into quartiles: Quartile 1 (<55), Quartile 2 (55–62), Quartile 3 (62–70), and Quartile 4 (≥ 70).

WC was measured at the level midway between the lower rib margin and the iliac crest, with participants in standing position without heavy outer garments and with emptied out pockets, breathing out gently (accuracy 1 cm). The cut-off points for abdominal obesity were chosen as a WC ≥ 90 cm for men and ≥ 80 cm for women [30].

HC was recorded as the maximum circumference over the buttocks (accuracy 1 cm). HC was classified into quartiles: Quartile 1 (<92), Quartile 2 (92–98), Quartile 3 (98–103), and Quartile 4 (≥ 103).

BMI was calculated as weight divided by height squared (kg/m^2). BMI was classified based on WHO guidelines as follows: underweight was a BMI < 18.50; normal weight was a BMI of 18.5–24.9; overweight was a BMI of 25.0–29.9; obese was a BMI ≥ 30.0 [31].

WHR was calculated as WC divided by HC. The cut-off value for central obesity was considered ≥ 0.8 for females and ≥ 0.9 for males [29].

WHtR was calculated as WC divided by height. The cut-off value for central obesity was considered ≥ 0.5 for both males and females [32].

ABSI based on WC adjusted for height and weight was developed by Krakauer N.Y. et al. [26].

$$\text{ABSI} = \frac{\text{WC}}{\text{BMI}^{\frac{2}{3}} \text{height}^{\frac{1}{2}}}$$

HI based on HC, height and weight was developed by Krakauer N.Y. et al. [25]

$$HI = HC \left(\frac{H}{\langle H \rangle} \right)^{0.310} \left(\frac{W}{\langle W \rangle} \right)^{-0.428}$$

where $\langle H \rangle = 166$ cm and $\langle W \rangle = 73$ kg.

ABSI and HI were both classified into quartiles: Quartile 1 was the lowest value and Quartile 4 (Q4) was the highest value.

2.4. Covariates

Socio-demographic factors included gender (male, female), age (60–64, 65–69, 70–74, 75–79, ≥ 80), marital status (single, married), empty nest (empty nest, non-empty nest), education level (no formal education, primary education, secondary or above), employment (employed, unemployed), and personal income (0–2500, 2500–5000, 5000–15,000, $\geq 15,000$ RMB Yuan). Health behavior-related factors included smoking status (nonsmoker, ex-smoker, current smoker) and drinking status (never drink, past drink, drink seldom, drink occasionally, drink more often). Health-related factors included self-rated health (very bad, bad, moderate, very good, excellent) and activities of daily living (ADL). Activities of daily living were assessed by using the Activities of Daily Living Scale (ADLs), which includes physical self-maintenance scale (PSMS) and instrumental activities of daily living scale (IADL). The ADL scale is used to evaluate people's simple and basic ability to practice one's normal life independently. The reliability and validity of ADL instrument in Chinese-language version was verified to be good [33]. Scores of ADL can be divided into three levels: no obvious dysfunction (≤ 14 scores), functional decline (15–21 scores), and obvious dysfunction (≥ 22 scores).

2.5. Statistical Analysis

We used SPSS 24.0 (IBM Corp, Armonk, NY, USA) to analyze the data. The socio-demographic characteristics of individuals with and without hypertension were compared by chi-square for categorical variables. Distributions of anthropometric indices between hypertensive and normotensive participants in both males and females were compared by chi-square for categorical variables. Binary logistic regression models with an enter method were used to examine the independent association of the nine different anthropometric indices and hypertension by gender. Model 1: binary logistic regression model by gender, a basic model adjusted for socio-demographic factors characteristics (gender, age, marital status, education level, employment, personal income, smoking status, drinking status, empty nest) and health-related factors (self-rated health, ADL scores), entered each time one of the anthropometric indices that show significant difference in chi-square. Model 2: binary logistic regression model by gender, the basic model further adjusted for all the other anthropometric indices that show significant difference in model 1. Associations were determined by odds ratios (ORs) and 95% confidence intervals. In addition, logistic regression analyses were also performed in two steps separately for male and female subjects and stratified for each 10-year age group. In the first step, each of the anthropometric indices was included as independent variables, adjusted for all the covariates. In the second step, the statistical anthropometric indices in step one included as independent variables, also adjusted for all the covariates. The statistical significance in this paper was set at the 5% level.

3. Results

Table 1 shows the demographic data of the study subjects. The study included 7070 respondents with a mean age of 69.81 (± 6.445) years. The prevalence of hypertension was 44.19%, 39.0% in males and 47.7% in females. Generally, the majority of the participants had the following characteristics: female (59.7%), married (81.2%), a primary or below education level (73.5%), unemployed (69.2%), nonsmokers (71.1%), nondrinkers (75.9%), and non-empty nest (72.8%). As for health characteristics, most of the elderly reported moderate and above health (81.7%) and with no obvious dysfunction (77.3%).

Table 1. Characteristics between hypertensive and normotensive participants.

Characteristics	Total (<i>n</i> = 7070) <i>n</i> (%)	Hypertensive (<i>n</i> = 3124) <i>n</i> (%)	Normotensive (<i>n</i> = 3946) <i>n</i> (%)	χ^2	<i>p</i>
Gender				51.920	0.000
Male	2846 (40.3)	1110 (39.0)	1736 (61.0)		
Female	4224 (59.7)	2014 (47.7)	2210 (52.3)		
Age				83.129	0.000
60–64	1577 (22.3)	561 (35.6)	1016 (64.4)		
65–69	2129 (30.1)	941 (44.2)	1188 (55.8)		
70–74	1780 (25.2)	898 (50.4)	882 (49.6)		
75–80	975 (13.8)	469 (48.1)	506 (51.9)		
≥80	609 (8.6)	255 (41.9)	354 (58.1)		
Residence				0.385	0.825
Rural	4990 (70.6)	2203 (44.1)	2787 (55.9)		
Township	524 (7.4)	226 (43.1)	298 (56.9)		
City	1556 (22.0)	695 (44.7)	861 (55.3)		
Marital status				19.931	0.000
Single	1331 (18.8)	661 (49.7)	670 (50.3)		
Married	5739 (81.2)	2463 (42.9)	3276 (57.1)		
Education level				18.985	0.000
No formal education	2270 (32.1)	1083 (47.7)	1187 (52.3)		
Primary education	2924 (41.4)	1268 (43.4)	1656 (56.6)		
Secondary or above	1876 (26.5)	773 (41.2)	1103 (58.8)		
Employment				21.085	0.000
Employed	2176 (30.8)	873 (40.1)	1303 (59.9)		
Unemployed	4894 (69.2)	2251 (46.0)	2643 (54.0)		
Personal income				10.436	0.015
0–2500	1571 (22.2)	673 (42.8)	898 (57.2)		
2500–5000	1689 (23.9)	792 (46.9)	897 (53.1)		
5000–15,000	1908 (27.0)	860 (45.1)	1048 (54.9)		
≥15,000	1902 (26.9)	799 (42.0)	1103 (58.0)		
Smoking status				72.709	0.000
Nonsmoker	5029 (71.1)	2334 (46.4)	2695 (53.6)		
Ex-smoker	903 (12.8)	418 (46.3)	485 (53.7)		
Current smoker	1138 (16.1)	372 (32.7)	766 (67.3)		
Drinking status				57.511	0.000
Never drink	5365 (75.9)	2453 (45.7)	2912 (54.3)		
Past drink	552 (7.8)	275 (49.8)	277 (50.2)		
Drink seldom	377 (16.3)	126 (33.4)	251 (66.6)		
Drink occasionally	98 (1.4)	33 (33.7)	65 (66.3)		
Drink more often	678 (9.6)	237 (35.0)	441 (65.0)		
Exercise time (per day)				3.779	0.286
Never	2719 (38.5)	1182 (43.5)	1537 (56.5)		
Less than half hour	2160 (30.6)	966 (44.7)	1194 (55.3)		
More than half hour	1547 (21.9)	707 (45.7)	840 (54.3)		
More than one hour	644 (9.1)	269 (41.8)	375 (58.2)		
Empty nest				5.790	0.016
Empty nest	1923 (27.2)	805 (41.9)	1118 (58.1)		
Non-empty nest	5147 (72.8)	2319 (45.1)	2828 (54.9)		
Self-rated health				336.988	0.000
Very bad	127 (1.8)	74 (58.3)	53 (41.7)		
Bad	1169 (16.5)	667 (57.1)	502 (42.9)		
Moderate	1992 (28.2)	1051 (52.8)	941 (47.2)		
Very good	2639 (37.3)	1043 (39.5)	1596 (60.5)		
Excellent	1143 (16.2)	289 (25.3)	854 (74.7)		
ADL scores				57.648	0.000
14	5467 (77.3)	2291 (41.9)	3176 (58.1)		
15–21	1283 (18.1)	646 (50.4)	637 (49.6)		
22+	320 (4.5)	187 (58.4)	133 (41.6)		

Distributions of anthropometric variables in males and females were showed as follows. Comparing individuals with and without hypertension, there were statistically significant differences in anthropometric indices except height, ABSI, and HI in males, and except height in females (Tables 2 and 3).

The results of binary logistic regression by gender are shown in Table 4. The results of model 1 showed that the association of weight, WC, HC, WHR, WHtR, and BMI with hypertension were statistically significant in both males and females, and that two new developed indicators (ABSI and HI) were statistically significant in females. After adjusting for other anthropometric indices that have statistical significance in model 1, WC and BMI were still statistically significant in the association with hypertension in males, HC, BMI, and HI were still statistically significant in females. Compared with the reference group, individuals with a larger WC or a higher BMI were associated with a greater risk for having hypertension in male, and those with a larger HC, a higher BMI or a larger HI were associated with a greater risk for having hypertension in females.

Table 2. Distributions of anthropometric variables in males

Characteristics	Total (<i>n</i> = 2846) <i>n</i> (%)	Hypertension (<i>n</i> = 1110) <i>n</i> (%)	Normotensive (<i>n</i> = 1736) <i>n</i> (%)	χ^2	<i>p</i>
Height				0.256	0.968
Q1	162 (5.7)	61 (37.7)	101 (62.3)		
Q2	554 (19.5)	213 (38.4)	341 (61.6)		
Q3	701 (24.6)	276 (39.4)	425 (60.6)		
Q4	1429 (50.2)	560 (39.2)	869 (60.8)		
Weight				89.441	0.000
Q1	490 (17.2)	128 (26.1)	362 (73.9)		
Q2	630 (22.1)	212 (33.7)	418 (66.3)		
Q3	891 (31.3)	347 (38.9)	544 (61.1)		
Q4	835 (29.3)	423 (50.7)	412 (49.3)		
WC				106.522	0.000
<90	831 (29.2)	202 (24.3)	629 (75.7)		
≥90	2015 (70.8)	908 (45.1)	1107 (54.9)		
HC				91.743	0.000
Q1	701 (24.6)	187 (26.7)	514 (73.3)		
Q2	774 (27.2)	280 (36.2)	494 (63.8)		
Q3	640 (22.5)	274 (42.8)	366 (57.2)		
Q4	731 (25.7)	369 (50.5)	362 (49.5)		
WHR				27.46	0.000
<0.9	713 (25.1)	219 (30.7)	494 (69.3)		
≥0.9	2133 (74.9)	891 (41.8)	1242 (58.2)		
WHtR				81.806	0.000
<0.5	612 (21.5)	142 (23.2)	470 (76.8)		
≥0.5	2234 (78.5)	968 (43.3)	1266 (56.7)		
BMI				107.422	0.000
Underweight	122 (4.3)	23 (18.9)	99 (81.1)		
Normal weight	1684 (59.2)	568 (33.7)	1116 (66.3)		
Overweight	910 (32.0)	434 (47.7)	476 (52.3)		
Obese	130 (4.6)	85 (65.4)	45 (34.6)		
ABSI				5.513	0.138
Q1	693 (24.3)	253 (36.5)	440 (63.5)		
Q2	786 (27.6)	296 (37.7)	490 (62.3)		
Q3	735 (25.8)	294 (40.0)	441 (60.0)		
Q4	632 (22.2)	267 (42.2)	365 (57.8)		
HI				3.228	0.358
Q1	974 (34.2)	363 (37.3)	611 (62.7)		
Q2	913 (32.1)	353 (38.7)	560 (61.3)		
Q3	585 (20.6)	237 (40.5)	348 (59.5)		
Q4	374 (13.1)	157 (42.0)	217 (58.0)		

Note: WC = waist circumference (cm), HC = hip circumference (cm), WHR = waist to hip ratio, WHtR = waist to height ratio, BMI = body mass index (kg/m²), ABSI = a body shape index, HI = hip index.

Table 3. Distributions of anthropometric variables in females

Characteristics	Total (<i>n</i> = 4224) <i>n</i> (%)	Hypertension (<i>n</i> = 2014) <i>n</i> (%)	Normotensive (<i>n</i> = 2210) <i>n</i> (%)	χ^2	<i>p</i>
Height				2.326	0.508
Q1	1713 (40.6)	834 (48.7)	879 (51.3)		
Q2	1807 (42.8)	860 (47.6)	947 (52.4)		
Q3	560 (13.3)	252 (45.0)	308 (55.0)		
Q4	144 (3.4)	68 (47.2)	76 (52.8)		
Weight				79.705	0.000
Q1	1324 (31.3)	516 (39.0)	808 (61.0)		
Q2	1129 (26.7)	523 (46.3)	606 (53.7)		
Q3	1110 (26.3)	613 (55.2)	497 (44.8)		
Q4	661 (15.6)	362 (54.8)	299 (45.2)		
WC				83.932	0.000
<80	527 (12.5)	153 (29.0)	374 (71.0)		
≥80	3697 (87.5)	1861 (50.3)	1836 (49.7)		
HC				107.647	0.000
Q1	929 (22.0)	324 (34.9)	605 (65.1)		
Q2	1042 (24.7)	469 (45.0)	573 (55.0)		
Q3	939 (22.2)	482 (51.3)	457 (48.7)		
Q4	1314 (31.1)	739 (56.2)	575 (43.8)		
WHR				8.319	0.004
<0.8	74 (1.8)	23 (31.1)	51 (68.9)		
≥0.8	4150 (98.2)	1991 (48.0)	2159 (52.0)		
WHtR				77.75	0.000
<0.5	395 (9.4)	105 (26.6)	290 (73.4)		
≥0.5	3829 (90.6)	1909 (49.9)	1920 (50.1)		
BMI				112.593	0.000
Underweight	112 (2.7)	25 (22.3)	87 (77.7)		
Normal weight	2001 (47.4)	832 (41.6)	1169 (58.4)		
Overweight	1716 (40.6)	911 (53.1)	805 (46.9)		
Obese	395 (9.4)	246 (62.3)	149 (37.7)		
ABSI				14.533	0.002
Q1	1080 (25.6)	469 (43.4)	611 (56.6)		
Q2	1006 (23.8)	481 (47.8)	525 (52.2)		
Q3	991 (23.5)	513 (51.8)	478 (48.2)		
Q4	1147 (27.2)	551 (48.0)	596 (52.0)		
HI				10.582	0.014
Q1	792 (18.8)	348 (43.9)	444 (56.1)		
Q2	857 (20.3)	410 (47.8)	447 (52.2)		
Q3	1179 (27.9)	603 (51.1)	576 (48.9)		
Q4	1396 (33.0)	653 (46.8)	743 (53.2)		

Note: WC = waist circumference (cm), HC = hip circumference (cm), WHR = waist to hip ratio, WHtR = waist to height ratio, BMI = body mass index (kg/m²), ABSI = A body shape index, HI = hip index.

The logistic regression analyses stratified for age were conducted in two steps. After the first step (data not shown), anthropometric indices that had statistical significance were chosen from each age group in both males and females, and entered in the second step (Table 5). Adjustment for these anthropometric indices and all the covariates affected all associations. Among the male elderly, only WC and BMI were statistical significant in the association with hypertension in the 60–70 year old group, and only BMI was statistically significant in the 70–80 year old group. However, there were no statistical significance in the relationship between hypertension and anthropometric indices in the male elderly older than 80. Among the female elderly, only HC, WHtR, and BMI were statistically significant in the association with hypertension in the 60–70 year old group; only HC, HI, ABSI, and BMI were statistically significant in the 70–80 year old group; and only HC was statistically significant in the older than 80 year old group. In these older age groups, the associations of all anthropometric indices and hypertension became weaker, disappeared or even went in the opposite direction.

Table 4. The relationship between hypertension and anthropometric variables in both males and females

Characteristics	Model 1	<i>p</i>	Female (<i>n</i> = 4224)	<i>p</i>	Model 2	<i>p</i>	Female (<i>n</i> = 4224)	<i>p</i>
	Male (<i>n</i> = 2846)				Male (<i>n</i> = 2846)			
Weight ^a		0.000		0.000		0.224		0.205
Q2	1.609 (1.223, 2.115)	0.001	1.502 (1.268, 1.779)	0.000	1.120 (0.817, 1.534)	0.481	0.964 (0.778, 1.195)	0.738
Q3	2.125 (1.642, 2.750)	0.000	2.241 (1.884, 2.667)	0.000	1.117 (0.792, 1.575)	0.527	1.083 (0.810, 1.449)	0.589
Q4	3.590 (2.749, 4.688)	0.000	2.283 (1.861, 2.800)	0.000	1.436 (0.949, 2.174)	0.087	0.850 (0.586, 1.232)	0.390
WC	2.716 (2.240, 3.293)	0.000	2.497 (2.034, 3.066)	0.000	1.936 (1.360, 2.756)	0.000	1.257 (0.899, 1.757)	0.182
HC ^a		0.000		0.000		0.898		0.003
Q2	1.620 (1.283, 2.044)	0.000	1.600 (1.325, 1.933)	0.000	0.922 (0.679, 1.252)	0.605	1.356 (1.076, 1.708)	0.010
Q3	2.143 (1.681, 2.732)	0.000	2.103 (1.731, 2.555)	0.000	0.889 (0.618, 1.277)	0.524	1.682 (1.255, 2.253)	0.000
Q4	2.923 (2.305, 3.708)	0.000	2.584 (2.153, 3.101)	0.000	0.947 (0.642, 1.396)	0.782	1.972 (1.371, 2.836)	0.000
WHR	1.623 (1.343, 1.961)	0.000	1.863 (1.116, 3.110)	0.017	0.976 (0.774, 1.230)	0.837	0.884 (0.500, 1.565)	0.673
WHtR	2.509 (2.022, 3.113)	0.000	2.654 (2.090, 3.371)	0.000	1.166 (0.834, 1.631)	0.368	1.311 (0.905, 1.899)	0.152
BMI ^b		0.000		0.000		0.000		0.004
Normal weight	2.599 (1.600, 4.223)	0.000	2.888 (1.816, 4.593)	0.000	1.693 (0.991, 2.890)	0.054	1.713 (1.037, 2.828)	0.035
Overweight	4.617 (2.811, 7.583)	0.000	4.898 (3.068, 7.818)	0.000	2.026 (1.124, 3.652)	0.019	2.055 (1.188, 3.555)	0.010
Obese	10.682 (5.813, 19.631)	0.000	6.819 (4.123, 11.280)	0.000	4.327 (2.138, 8.758)	0.000	2.927 (1.584, 5.410)	0.001
ABSI ^c				0.036				0.388
Q2			1.136 (0.950, 1.360)	0.163			1.024 (0.841, 1.247)	0.813
Q3			1.292 (1.078, 1.549)	0.006			1.162 (0.942, 1.434)	0.161
Q4			1.063 (0.889, 1.271)	0.503			1.018 (0.813, 1.276)	0.875
HI ^c				0.031				0.028
Q2			1.170 (0.956, 1.432)	0.128			0.902 (0.714, 1.138)	0.384
Q3			1.281 (1.061, 1.547)	0.010			0.910 (0.707, 1.172)	0.464
Q4			1.052 (0.875, 1.263)	0.590			0.701 (0.521, 0.943)	0.019

Note: The bold indicate significance. WC = waist circumference (cm), HC = hip circumference (cm), WHR = waist to hip ratio, WHtR = waist to height ratio, BMI = body mass index (kg/m²), ABSI = A body shape index, HI = hip index; ^a Reference category is Q1. ^b Reference category is underweight. ^c Reference category is Q4.

Table 5. The relationship between hypertension and anthropometric variables in both males and females stratified by age

Characteristics	Male	70–80 (<i>n</i> = 1199)	80+ (<i>n</i> = 261)	Female	70–80 (<i>n</i> = 1556)	80+ (<i>n</i> = 348)
	60–70 (<i>n</i> = 1386)			60–70 (<i>n</i> = 2320)		
Weight ^a						
Q2	1.076 (0.634–1.827)	1.446 (0.918–2.276)	0.427 (0.145–1.255)	1.042 (0.786–1.383)	0.952 (0.672–1.349)	1.309 (0.663–2.581)
Q3	1.099 (0.621–1.944)	1.346 (0.816–2.220)	0.898 (0.261–3.083)	1.194 (0.829–1.718)	1.113 (0.689–1.799)	0.894 (0.335–2.389)
Q4	1.501 (0.774–2.911)	1.639 (0.885–3.036)	1.095 (0.230–5.217)	0.896 (0.575–1.394)	1.000 (0.533–1.876)	2.650 (0.324–21.683)
WC	2.454 (1.423–4.233) **	1.454 (0.862–2.452)	2.578 (0.802–8.284)	0.927 (0.576–1.492)	1.718 (0.980–3.011)	1.286 (0.504–3.280)
HC ^a						
Q2	0.873 (0.545–1.397)	0.821 (0.521–1.292)	2.094 (0.652–6.726)	1.263 (0.935–1.707)	1.507 (1.028–2.209) **	1.009 (0.532–1.912)
Q3	0.954 (0.554–1.644)	0.730 (0.422–1.265)	0.989 (0.226–4.323)	1.389 (1.001–1.928) **	1.992 (1.228–3.231) **	1.199 (0.517–2.781)
Q4	0.794 (0.442–1.429)	0.989 (0.549–1.781)	1.257 (0.237–6.676)	1.331 (0.935–1.896)	2.364 (1.287–4.342) **	2.655 (1.142–6.169) **
WHR	0.791 (0.562–1.114)	1.218 (0.854–1.738)		1.023 (0.491–2.131)		
WHtR	1.059 (0.637–1.761)	1.228 (0.748–2.018)	1.474 (0.424–5.126)	1.727 (1.033–2.887) **	0.766 (0.406–1.445)	1.457 (0.514–4.134)
BMI ^b						
Normal weight	2.032 (0.735–5.616)	1.665 (0.813–3.410)		1.628 (0.725–3.654)	2.181 (0.988–4.815)	1.172 (0.390–3.522)
Overweight	2.588 (0.876–7.644)	1.934 (0.862–4.341)		2.602 (1.103–6.139) **	2.077 (0.871–4.948)	0.855 (0.229–3.197)
Obese	6.745 (1.975–23.040) **	4.372 (1.582–12.082) **		3.614 (1.441–9.067) **	3.049 (1.123–8.281) **	3.563 (0.488–26.005)
ABSI ^c						
Q2					1.191 (0.841–1.686)	
Q3					1.466 (1.025–2.097) **	
Q4					1.179 (0.808–1.720)	
HI ^c						
Q2			1.233 (0.491–3.097)		0.792 (0.538–1.166)	
Q3			1.123 (0.406–3.108)		0.869 (0.572–1.321)	
Q4			2.322 (0.653–8.262)		0.553 (0.341–0.898) **	

Note: The bold indicate significance. WC = waist circumference (cm), HC = hip circumference (cm), WHR = waist to hip ratio, WHtR = waist to height ratio, BMI = body mass index (kg/m²), ABSI = A body shape index, HI = hip index. ^a Reference category is Q1. ^b Reference category is underweight. ^c Reference category is Q4. ***p* < 0.05.

4. Discussion

Our research showed that there were significant differences in most of the anthropometric indices between hypertensive patients and normotensive individuals, including both overall and abdominal obesity indicators. These results suggested that hypertension is linked to obesity in both male and female elderly, which is inconsistent with previous studies that both overall and abdominal obesity are significantly associated with hypertension [34,35]. The mechanisms of this association include structural arterial abnormalities, leptin, and the activation of the renin–angiotensin–aldosterone axis [13,36]. Furthermore, other studies showed that it may also be related to the variant of gene [37,38]. Fortunately, obesity being a modifiable factor, lifestyle modifications including increased physical activity, endurance and endurance-strength exercise, and dietary modifications can decrease the incidence of hypertension [39–41].

In the present study, we found that there were gender differences and age differences in the relationship between anthropometric indices and the prevalence of hypertension in Chinese elderly. In both genders, association was stronger between BMI and hypertension without considering the impact of age. These results are supported by numerous studies that the increase of BMI contributes to blood pressure increase [42–44]. After we took age into account, the association of BMI and hypertension in both males and females was weaker in the older elderly compared to those younger ones. However, the association of HI and hypertension in females was stronger with increased age.

After stratified by gender and age, BMI in men (except individuals older than 80) and HC in women remained significant in the relationship with hypertension in all age groups. Our findings in males are consistent with previous studies that BMI have the highest correlation with hypertension [45,46]. Besides, analyses of our findings revealed a positive relationship between HC and the incidence of hypertension in females. This relationship is inconsistent with most of the previous studies [46,47]. However, there are still some studies support us that high HC is a risk factor for the multi-metabolic disorders [48–50]. This result may be due to the following two reasons. First, HC carries some information on both overall obesity and abdominal obesity, since HC is positively correlated with BMI and WC [48]. Second, older Chinese women with a higher HC usually have experienced gravidity more times, studies have indicated that more gravidity was associated with a consistent increase in the risk of metabolic syndrome in older Chinese women [51,52]. In view of these results, it should be a priority to encourage those males with high BMI and females with high HC to pay more attention to guidelines of hypertension. Besides, health care facilities should strengthen the screening and monitoring for those at-risk subgroups to early identify and treat the elderly with hypertension.

The main results of the current study indicated that, overall, obesity was associated with hypertension in males, and partial obesity was associated with hypertension in females. This may be concluded that the amount of body fat mass in males and the distribution of body fat in females is closely related to hypertension [16]. In previous studies, general obesity and partial obesity usually showed the similar relationship with hypertension in different gender. For example, some studies indicated that general obesity has a stronger association with hypertension in both males and females [15,24]. Others found that the association of partial obesity with hypertension was stronger in both genders [14,19]. Our results indicated that the role of overall obesity and partial obesity in hypertension in separate gender may be different. Further studies are necessary to confirm this assumption.

This study has a number of strengths. First, we included more anthropometric indices than previous studies, which made our results more comprehensive. Second, to our best knowledge, this is the first study to analyze the gender and age difference of association between anthropometric indices and hypertension in Chinese elderly. Furthermore, the sample size in this study was relatively large, which makes our results more reliable.

However, there are some limitations that should be considered while interpreting the results. First, the cross-sectional design does not allow for causal inferences to be made about the relationships of obesity with the prevalence of hypertension. However, the largely causal association between obesity

and hypertension is well established [53]. Thus, it can be speculated that excess obesity precedes hypertension. Second, there may be more confounding factors than those available for consideration in our study. Third, using cut points for defining the metabolic disorders implies a loss of information, and more studies are needed in order to find out the relationships between anthropometric indices and continuous metabolic variables by use of linear regression.

5. Conclusions

In summary, there are age and gender differences in the association between anthropometric indices of obesity and hypertension in Chinese elderly. Moreover, BMI in males (except individuals aged ≥ 80 years) and HC in females were found to have a significant association with hypertension in all age groups.

These findings indicate a need to develop gender-specific strategies for the male and female elderly in the primary and secondary prevention of hypertension.

Acknowledgments: We would like to thank the center for health economics experiment and public policy, School of Public Health, Shandong University. This work was supported by two grants of National Natural Science Foundation of China (grant numbers: 71673169 and 71673170). The author is also grateful for the assistance in data collection processes of the master students from School of Public Health, Shandong University, they are: Yu Xia, Yali Wang, Lulu Ding, Yawei Guo, Anan Jiao, Yanran Huang, Xiang Jing, Rui Yu, Yan Hu, Hongjuan Liu, Lina Su, Ningze Xu, Xiuxiu Wu, Yaozu Li, Lu Lu, Yuheng Xin, Benyu Su, Qiangdong Guan, Dandan Ge, Li Zhang, Jing Zhang, Zhengyue Jing, and Qiongqiong Xu.

Author Contributions: Lingzhong Xu conceived the idea. Long Sun and Jiajia Li designed and supervised the analysis. Wenzhe Qin and Gan Ding participated in the statistical analysis and interpretation of the results. Qian Wang drafted the manuscript. Jiao Zhang, Jing Zhu, Su Xie, and Zihang Yu gave many valuable comments on the draft and also polished it. All authors read and approved the final manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

Abbreviations

BMI	body mass index
HC	hip circumference
CVD	cardiovascular disease
WC	waist circumference
WHR	waist-to-hip ratio
WHtR	waist-to-height
HI	hip index
ABSI	body shape index
PSUs	primary sampling units
SSUs	secondary sampling units
ADL	activities of daily living
ADLs	activities of daily living scale
IADL	instrumental activities of daily living scale
PSMS	physical self-maintenance scale
IADL	activities of daily living scale
ORs	odds ratios

References

1. Global Burden of Metabolic Risk Factors for Chronic Diseases, C. Cardiovascular disease, chronic kidney disease, and diabetes mortality burden of cardiometabolic risk factors from 1980 to 2010: A comparative risk assessment. *Lancet Diabetes Endocrinol.* **2014**, *2*, 634–647.
2. Lim, S.S.; Vos, T.; Flaxman, A.D.; Danaei, G.; Shibuya, K.; Adair-Rohani, H.; Amann, M.; Anderson, H.R.; Andrews, K.G.; Aryee, M.; et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: A systematic analysis for the global burden of disease study 2010. *Lancet* **2012**, *380*, 2224–2260. [[CrossRef](#)]

3. He, J.; Gu, D.; Chen, J.; Wu, X.; Kelly, T.N.; Huang, J.F.; Chen, J.C.; Chen, C.S.; Bazzano, L.A.; Reynolds, K.; et al. Premature deaths attributable to blood pressure in China: A prospective cohort study. *Lancet* **2009**, *374*, 1765–1772. [CrossRef]
4. Lu, J.; Lu, Y.; Wang, X.; Li, X.; Linderman, G.C.; Wu, C.; Cheng, X.; Mu, L.; Zhang, H.; Liu, J.; et al. Prevalence, awareness, treatment, and control of hypertension in china: Data from 1.7 million adults in a population-based screening study (China peace million persons project). *Lancet* **2017**, *390*, 2549–2558. [CrossRef]
5. Haslam, D.W.; James, W.P. Obesity. *Lancet* **2005**, *366*, 1197–1209. [CrossRef]
6. Obesity and Overweight. Available online: <http://www.Who.Int/mediacentre/factsheets/fs311/en/> (accessed on 20 February 2018).
7. Genelhu, V.A.; Celoria, B.M.; Duarte, S.F.; Cabello, P.H.; Francischetti, E.A. Not all obese subjects of multiethnic origin are at similar risk for developing hypertension and type 2 diabetes. *Eur. J. Intern. Med.* **2009**, *20*, 289–295. [CrossRef] [PubMed]
8. Hall, J.E. The kidney, hypertension, and obesity. *Hypertension* **2003**, *41*, 625–633. [CrossRef] [PubMed]
9. Klein, S.; Burke, L.E.; Bray, G.A.; Blair, S.; Allison, D.B.; Pi-Sunyer, X.; Hong, Y.; Eckel, R.H. Metabolism. Clinical implications of obesity with specific focus on cardiovascular disease: A statement for professionals from the american heart association council on nutrition, physical activity, and metabolism: Endorsed by the american college of cardiology foundation. *Circulation* **2004**, *110*, 2952–2967. [PubMed]
10. Brown, C.D.; Higgins, M.; Donato, K.A.; Rohde, F.C.; Garrison, R.; Obarzanek, E.; Ernst, N.D.; Horan, M. Body mass index and the prevalence of hypertension and dyslipidemia. *Obes. Res.* **2000**, *8*, 605–619. [CrossRef] [PubMed]
11. Colin-Ramirez, E.; Castillo-Martinez, L.; Orea-Tejeda, A.; Villa Romero, A.R.; Vergara Castaneda, A.; Asensio Lafuente, E. Waist circumference and fat intake are associated with high blood pressure in Mexican children aged 8 to 10 years. *J. Am. Diet. Assoc.* **2009**, *109*, 996–1003. [CrossRef] [PubMed]
12. Francischetti, E.A.; Genelhu, V.A. Obesity-hypertension: An ongoing pandemic. *Int. J. Clin. Pract.* **2007**, *61*, 269–280. [CrossRef] [PubMed]
13. Rahmouni, K.; Correia, M.L.; Haynes, W.G.; Mark, A.L. Obesity-associated hypertension: New insights into mechanisms. *Hypertension* **2005**, *45*, 9–14. [CrossRef] [PubMed]
14. Lee, C.M.; Huxley, R.R.; Wildman, R.P.; Woodward, M. Indices of abdominal obesity are better discriminators of cardiovascular risk factors than BMI: A meta-analysis. *J. Clin. Epidemiol.* **2008**, *61*, 646–653. [CrossRef] [PubMed]
15. Khashayar, P.; Aghaei Meybodi, H.; Rezaei Hemami, M.; Larijani, B. Role of obesity variables in detecting hypertension in an iranian population. *High Blood Press. Cardiovasc. Prev.* **2017**, *24*, 305–312. [CrossRef] [PubMed]
16. Cho, Y.G.; Song, H.J.; Kim, J.M.; Park, K.H.; Paek, Y.J.; Cho, J.J.; Caterson, I.; Kang, J.G. The estimation of cardiovascular risk factors by body mass index and body fat percentage in Korean male adults. *Metabolism* **2009**, *58*, 765–771. [CrossRef] [PubMed]
17. Frankenfield, D.C.; Rowe, W.A.; Cooney, R.N.; Smith, J.S.; Becker, D. Limits of body mass index to detect obesity and predict body composition. *Nutrition* **2001**, *17*, 26–30. [CrossRef]
18. Watanabe, J.; Tochikubo, O. Relationship between visceral fat accumulation and hypertension in obese men. *Clin. Exp. Hypertens.* **2003**, *25*, 199–208. [CrossRef] [PubMed]
19. Woo, J.; Ho, S.C.; Yu, A.L.; Sham, A. Is waist circumference a useful measure in predicting health outcomes in the elderly? *Int. J. Obes. Relat. Metab. Disord.* **2002**, *26*, 1349–1355. [CrossRef] [PubMed]
20. Hsieh, S.D.; Muto, T. Metabolic syndrome in japanese men and women with special reference to the anthropometric criteria for the assessment of obesity: Proposal to use the waist-to-height ratio. *Prev. Med.* **2006**, *42*, 135–139. [CrossRef] [PubMed]
21. Dobbeltsteyn, C.J.; Joffres, M.R.; MacLean, D.R.; Flowerdew, G. A comparative evaluation of waist circumference, waist-to-hip ratio and body mass index as indicators of cardiovascular risk factors. The Canadian heart health surveys. *Int. J. Obes. Relat. Metab. Disord.* **2001**, *25*, 652–661. [CrossRef] [PubMed]
22. Shiwaku, K.; Anuurad, E.; Enkhmaa, B.; Nogi, A.; Kitajima, K.; Yamasaki, M.; Yoneyama, T.; Oyunsuren, T.; Yamane, Y. Predictive values of anthropometric measurements for multiple metabolic disorders in Asian populations. *Diabetes Res. Clin. Pract.* **2005**, *69*, 52–62. [CrossRef] [PubMed]

23. Zimmet, P.; Magliano, D.; Matsuzawa, Y.; Alberti, G.; Shaw, J. The metabolic syndrome: A global public health problem and a new definition. *J. Atheroscler. Thromb.* **2005**, *12*, 295–300. [CrossRef] [PubMed]
24. Decoda Study, G.; Nyamdorj, R.; Qiao, Q.; Lam, T.H.; Tuomilehto, J.; Ho, S.Y.; Pitkanen, J.; Nakagami, T.; Mohan, V.; Janus, E.D.; et al. Bmi compared with central obesity indicators in relation to diabetes and hypertension in Asians. *Obesity* **2008**, *16*, 1622–1635. [CrossRef] [PubMed]
25. Krakauer, N.Y.; Krakauer, J.C. An anthropometric risk index based on combining height, weight, waist, and hip measurements. *J. Obes.* **2016**, *2016*, 8094275. [CrossRef] [PubMed]
26. Krakauer, N.Y.; Krakauer, J.C. A new body shape index predicts mortality hazard independently of body mass index. *PLoS ONE* **2012**, *7*, e39504. [CrossRef] [PubMed]
27. National Bureau of Statistic of China. The National Economic and Social Development Statistical Bulletin in 2015. Available online: http://www.Stats.Gov.Cn/tjsj/zxfb/201602/t20160229_1323991.html (accessed on 21 May 2017).
28. Yimin, C. Relationship of hypertension with BMI, waist circumference, and WHR in 6830 elderly people. *Zhejiang J. Integr. Tradit. Chin. Western Med.* **2015**, *24*, 3.
29. Ma, Y.; Zhang, B.; Wang, H.; Du, W.; Su, C.; Zhai, F. Relationship between adiposity indicators and blood pressure of urban and rural residents in China. *Wei Sheng Yan Jiu* **2012**, *41*, 70–74. [PubMed]
30. Alberti, K.G.; Zimmet, P.; Shaw, J. The metabolic syndrome—A new worldwide definition. *Lancet* **2005**, *366*, 1059–1062. [CrossRef]
31. Who, BMI Classification. 2015. Available online: http://apps.who.int/bmi/index.jsp?introPage=intro_3.html (accessed on 19 April 2018).
32. Guangdong Provincial Co-operation Group for Diabetes Epidemiological Study. Waist/height ratio: An effective index for abdominal obesity predicting diabetes and hypertension. *Chin. J. Endocrinol. Metab.* **2004**, *20*, 4.
33. Lawton, M.P.; Brody, E.M. Assessment of older people: Self-maintaining and instrumental activities of daily living. *Gerontologist* **1969**, *9*, 179–186. [CrossRef] [PubMed]
34. He, Y.H.; Jiang, G.X.; Yang, Y.; Huang, H.E.; Li, R.; Li, X.Y.; Ning, G.; Cheng, Q. Obesity and its associations with hypertension and type 2 diabetes among Chinese adults age 40 years and over. *Nutrition* **2009**, *25*, 1143–1149. [CrossRef] [PubMed]
35. Nguyen, N.T.; Magno, C.P.; Lane, K.T.; Hinojosa, M.W.; Lane, J.S. Association of hypertension, diabetes, dyslipidemia, and metabolic syndrome with obesity: Findings from the national health and nutrition examination survey, 1999 to 2004. *J. Am. Coll. Surg.* **2008**, *207*, 928–934. [CrossRef] [PubMed]
36. Nguyen, T.; Lau, D.C. The obesity epidemic and its impact on hypertension. *Can. J. Cardiol.* **2012**, *28*, 326–333. [CrossRef] [PubMed]
37. Baudrand, R.; Goodarzi, M.O.; Vaidya, A.; Underwood, P.C.; Williams, J.S.; Jeunemaitre, X.; Hopkins, P.N.; Brown, N.; Raby, B.A.; Lasky-Su, J.; et al. A prevalent caveolin-1 gene variant is associated with the metabolic syndrome in Caucasians and Hispanics. *Metabolism* **2015**, *64*, 1674–1681. [CrossRef] [PubMed]
38. Skrypnik, K.; Suliburska, J.; Skrypnik, D.; Pilarski, L.; Regula, J.; Bogdanski, P. The genetic basis of obesity complications. *Acta Sci. Pol. Technol. Aliment.* **2017**, *16*, 83–91. [CrossRef] [PubMed]
39. Skrypnik, D.; Bogdanski, P.; Madry, E.; Karolkiewicz, J.; Ratajczak, M.; Krysiak, J.; Pupek-Musialik, D.; Walkowiak, J. Effects of endurance and endurance strength training on body composition and physical capacity in women with abdominal obesity. *Obes. Facts* **2015**, *8*, 175–187. [CrossRef] [PubMed]
40. Skrypnik, D.; Ratajczak, M.; Karolkiewicz, J.; Madry, E.; Pupek-Musialik, D.; Hansdorfer-Korzon, R.; Walkowiak, J.; Jakubowski, H.; Bogdanski, P. Effects of endurance and endurance-strength exercise on biochemical parameters of liver function in women with abdominal obesity. *Biomed. Pharmacother.* **2016**, *80*, 1–7. [CrossRef] [PubMed]
41. Szulinska, M.; Skrypnik, D.; Ratajczak, M.; Karolkiewicz, J.; Madry, E.; Musialik, K.; Walkowiak, J.; Jakubowski, H.; Bogdanski, P. Effects of endurance and endurance-strength exercise on renal function in abdominally obese women with renal hyperfiltration: A prospective randomized trial. *Biomed. Environ. Sci.* **2016**, *29*, 706–712. [PubMed]
42. Droyvold, W.B.; Midthjell, K.; Nilsen, T.I.; Holmen, J. Change in body mass index and its impact on blood pressure: A prospective population study. *Int. J. Obes.* **2005**, *29*, 650–655. [CrossRef] [PubMed]
43. Wilsgaard, T.; Schirmer, H.; Arnesen, E. Impact of body weight on blood pressure with a focus on sex differences: The tromso study, 1986–1995. *Arch. Intern. Med.* **2000**, *160*, 2847–2853. [CrossRef] [PubMed]

44. Jousilahti, P.; Tuomilehto, J.; Vartiainen, E.; Valle, T.; Nissinen, A. Body mass index, blood pressure, diabetes and the risk of anti-hypertensive drug treatment: 12-year follow-up of middle-aged people in Eastern Finland. *J. Hum. Hypertens.* **1995**, *9*, 847–854. [[PubMed](#)]
45. Chen, S.C.; Lo, T.C.; Chang, J.H.; Kuo, H.W. Variations in aging, gender, menopause, and obesity and their effects on hypertension in Taiwan. *Int. J. Hypertens.* **2014**, *2014*, 515297. [[CrossRef](#)] [[PubMed](#)]
46. Zhang, Z.Q.; Deng, J.; He, L.P.; Ling, W.H.; Su, Y.X.; Chen, Y.M. Comparison of various anthropometric and body fat indices in identifying cardiometabolic disturbances in Chinese men and women. *PLoS ONE* **2013**, *8*, e70893. [[CrossRef](#)] [[PubMed](#)]
47. Snijder, M.B.; Zimmet, P.Z.; Visser, M.; Dekker, J.M.; Seidell, J.C.; Shaw, J.E. Independent and opposite associations of waist and hip circumferences with diabetes, hypertension and dyslipidemia: The ausdiab study. *Int. J. Obes. Relat. Metab. Disord.* **2004**, *28*, 402–409. [[CrossRef](#)] [[PubMed](#)]
48. Wang, Z.; Hoy, W.E. Waist circumference, body mass index, hip circumference and waist-to-hip ratio as predictors of cardiovascular disease in aboriginal people. *Eur. J. Clin. Nutr.* **2004**, *58*, 888–893. [[CrossRef](#)] [[PubMed](#)]
49. Liu, S.J.G.Z.; Hu, X.S.; Wu, M.; Chen, F.M.; Zhou, H.; Kang, G.D. Relationship between multi-metabolic disorders and hip circumference in Jiangsu community population. *Chin. J. Public Health* **2008**, *24*, 3.
50. Wang, S.L.; Pan, W.H.; Hwu, C.M.; Ho, L.T.; Lo, C.H.; Lin, S.L.; Jong, Y.S. Incidence of niddm and the effects of gender, obesity and hyperinsulinaemia in Taiwan. *Diabetologia* **1997**, *40*, 1431–1438. [[CrossRef](#)] [[PubMed](#)]
51. Lao, X.Q.; Thomas, G.N.; Jiang, C.Q.; Zhang, W.S.; Yin, P.; Schooling, M.; Heys, M.; Leung, G.M.; Adab, P.; Cheng, K.K.; et al. Parity and the metabolic syndrome in older Chinese women: The Guangzhou biobank cohort study. *Clin. Endocrinol.* **2006**, *65*, 460–469. [[CrossRef](#)] [[PubMed](#)]
52. Xu, F.H.; Huo, Y.N.; Wu, M.B.; Liu, J.D.; Zhang, Y.L. Relationship of parity and other factors with metabolic syndrome in middle-age and elderly women in Nanchang community. *Chin. J. Endocrinol. Metab.* **2014**, *30*, 4.
53. Taing, K.Y.; Farkouh, M.E.; Moineddin, R.; Tu, J.V.; Jha, P. Age and sex-specific associations of anthropometric measures of adiposity with blood pressure and hypertension in India: A cross-sectional study. *BMC Cardiovasc. Disord.* **2016**, *16*, 247. [[CrossRef](#)] [[PubMed](#)]



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).