



Table S1. Comparison of puff behavior of male and female smokers

	All (n = 300) (Median (IQR))	Male (n = 250) (Median (IQR))	Female (n = 50) (Median (IQR))	p-value (M:F)
No of Cigarette (w/ CReSS)	7.0 (5.0-11.0)	7.0 (5.0-11.0)	7.0 (5.0-10.0)	0.8143
Puff count	16 (13.5-19.0)	15.5 (13.0-19.0)	17.5 (15.0-21.0)	0.0123
Puff vol (mL)	60.6 (49.9-73.9)	62.7 (52.7-75.5)	53.5 (42.0-64.2)	0.0019
Duration (mSec)	1552.8 (1243.0-1908.7)	1575.7 (1243.0-1923.0)	1408.9 (1215.6-1847.0)	0.0635
AvgFlow (mL/sec)	40.4 (32.1-51.1)	41.2 (33.1-52.2)	36.4 (30.0-43.7)	0.0304
Peak Flow (mL/sec)	62.7 (49.4-82.4)	66.1 (50.4-85.1)	59.8 (44.3-75.1)	0.3534
Time to Peak (mSec)	493.9 (371.5-640.1)	503.4 (376.4-645.7)	451.1 (347.2-582.3)	0.122
InterPuff Interval (mSec)	8828.9 (6528.6- 11196.1)	8931.7 (6546.7- 11281.1)	8343.2 (6206.3- 10992.9)	0.122

* Mann-Whitney test to compare male's frequency with female's one

Table S2. Comparison of distribution of Fagerstrom score between flavored Composition of and non-flavored cigarette smokers

	NO	1	2	3	4	p-value*
		(Low)	(Low-Mod)	(Moderate)	(High)	
	n (%)	n (%)	n (%)	n (%)	n (%)	
Flavored	YES					0.0155
NO						
Capsule						
Additive	63(26.9)	98 (41.9)	67 (28.6)	6 (2.6)		
Menthol						
(n = 234)						
Flavored	YES					0.0155
NO						
Capsule						
Additive	28(42.4)	29 (44.0)	9 (13.6)	0 (0)		
or						
Menthol						
(n = 66)						


* p-value from Chi-square test

Table S3. Level of urinary cotinine according to the level of daily total puff volume

	Vol. per_day_<=4542.0mL	Vol. per_day_4542.1~7071.0mL	Vol. per_day_7071.1~11556.0mL	Vol. per_day_11556.1 + mL	p-value*	
	p50	(IQR)	p50	(IQR)	p50	(IQR)
Nicotine content in cigarette (mg)	0.4	(0.2~0.5)	0.4	(0.2~0.5)	0.4	(0.1~0.5)
Total vol (ml)/day	2840.0	(2034.6~3862.3)	5964.0	(5134.5~6448.0)	8672.5	(7728.5~10166.4)
Cotinine (ng/mL)	905.4	(432.5~1413.2)	933.1	(594.4~1486.2)	1089.9	(787.0~1650.9)
						<0.001
						0.0011

*p-value from Kruskal-Wallis Test

1



2 Figure S1. Comparison of puff measures (medina (interquartile range)) between m smokers of high-nicotine-dose cigarettes and low-nicotine-dose cigarettes (puff frequency per
3 cigarette, puff volume, puff duration, average flow rate, peak flow rate, inter puff interval)

4