

International Journal of *Environmental Research and Public Health*



Article Seasonal Fluctuation of Polycyclic Aromatic Hydrocarbons and Aerosol Genotoxicity in Long-Range Transported Air Mass Observed at the Western End of Japan

Takashi Kubo¹, Wenzhi Bai², Masaki Nagae¹ and Yuji Takao^{1,*}

- ¹ Faculty of Environmental Science, Nagasaki University, Nagasaki 852-8521, Japan; kubo-t@nagasaki-u.ac.jp (T.K.); nagae@nagasaki-u.ac.jp (M.N.)
- ² Graduate School of Fisheries and Environmental Sciences, Nagasaki University, Nagasaki 852-8521, Japan; wbbabjw@yahoo.co.jp
- * Correspondence: takao@nagasaki-u.ac.jp; Tel.: +81-95-819-2753

Received: 26 December 2019; Accepted: 9 February 2020; Published: 13 February 2020

Abstract: In order to clarify the level transboundary air pollution caused by polycyclic aromatic hydrocarbons (PAHs) and genotoxic substances, aerosols were collected from forest and suburban sites in Nagasaki, west Japan, for 6 years. The PAH concentration was measured, and the genotoxicity of the substances were evaluated using the umu test. The results showed no notable trends in the concentration or toxicity of either sites throughout the study period. The suburban and forest sites shared similar seasonal fluctuation patterns and quantitative values, suggesting that the western end of Japan might be affected by long-range transported pollutants, especially in winter. PAH concentration and genotoxicity showed the same seasonal patterns of increased levels in winter and lower levels in summer. This suggests that PAHs and genotoxic substances were correlated and share common sources. Back trajectory and source analyses were conducted using the diagnostic ratios of PAHs. It was predicted that air pollution by PAHs at the forest site arise predominantly as a result of biomass or coal combustion in continental regions, such as northern parts of China and the Korean Peninsula. This is particularly expected in winter. Therefore, genotoxic substances would also be strongly influenced by transboundary pollution from the continental region. In addition, it was estimated that the contribution of transboundary PAH pollution could reach 70% at the suburban site in winter.

Keywords: polycyclic aromatic hydrocarbons; genotoxicity; transboundary pollution; seasonal fluctuation; pollution sources

1. Introduction

Polycyclic aromatic hydrocarbons (PAHs) are typical pollutants that are unintentionally generated, mainly due to incomplete combustion of organic matter. In addition, since they are present in minute amounts in crude oil and petroleum, they are also released into the environment when these substances leak. Many PAHs have been reported as carcinogens and/or genotoxicants. In particular, benzo[a]pyrene (BaP) is a well-known carcinogen. The International Agency for Research on Cancer (IARC) classifies BaP as Group 1 (carcinogenic to humans). Several other PAHs are classified as Group 2A (probably carcinogenic to humans) and 2B (possibly carcinogenic to humans) [1]. Genotoxicity has been quantitatively evaluated for some PAHs. While metabolism reduces the toxicity of many chemicals, it has been reported that PAH genotoxicity tends to be enhanced after reaction with metabolic enzymes [2].

Up until the present, PAHs have been detected in various environmental samples such as river water [3–5], sediment [6–8], soil [9–11], and airborne particles [12–14], which indicates the spread of PAHs pollution. Various environmental emission sources have also been investigated, such as waste water [15–17], exhaust gas [18–20], biomass burning [21–23], and petroleum-related products [24–26]. Attempts have also been made to estimate emission sources from the measured concentration ratio of PAHs in the environmental samples [27–29].

Environmental pollution is not always restricted to the vicinity of the pollutants' emission sources. It can create challenges thousands of kilometers away, across the border. This so-called transboundary pollution is an old and new environmental problem. The trail smelter case that began in the 1920s, which led to the establishment of international environmental laws, was caused by transboundary air pollution [30]. The philosophy of "states have the responsibility to ensure that activities within their jurisdiction do not cause damage to the environment of other states" has been shared internationally. It has been affirmed repeatedly in international declarations and conventions, such as the Rio Declaration on Environment and Development [31] and the United Nations Framework Convention on Climate Change [32]. River pollution [33–35], acid rain [36–38], and aerosols [39–41] have been related to transboundary pollution. Since the late 1970s, investigations into PAH-related transboundary air pollution have also been reported [42].

PAHs exist as particulates or gaseous molecules in the atmosphere. It has been reported that some PAHs having four benzene rings, and all PAHs having more than four benzene rings exist in the particulate phase [43]. The concentrations of PAHs in the atmosphere have been measured in various regions, including Europe and Asia [44–46]; some of these investigations have indicated the probability of transboundary pollution [47–49]. Hayakawa et al. not only measured atmospheric PAHs across Japan for a long time [50–52], but also took measurements in foreign continents, including Africa [53–55]. Although some studies have compared PAHs by seasons [56–58], few studies have considered data over several years in combination with toxicity assays.

Genotoxicity is closely related to carcinogenicity. The Ames mutagenicity test is commonly used for evaluating the genotoxicity of environmental samples [59–61]. In recent years, however, there has been increased use of the umu test for testing DNA damage, particularly in aquatic environments [62–64]. Few studies have used the umu test to investigate the atmosphere, including seasonal changes in atmospheric genotoxicity. There seems to be no longitudinal studies on the use of umu tests in conjunction with PAH measurements. Unlike PAHs, no methods exist for using umu test results to estimate the source of genotoxicity. Therefore, there are few detailed reports on the origin of genotoxic substances in the atmosphere.

As mentioned above, since some PAHs are well-known carcinogens, their environmental behavior needs to be better understood. In addition to this, many things relating to genotoxic substances in the atmosphere remain unexplained, including their sources. If PAH measurements and umu tests are performed on air samples which have been collected at sites with different anthropogenic effects, their characteristics and sources could be estimated from their long-term fluctuation patterns. The objective of this study was to quantify seasonal fluctuation patterns and trends over several years using PAH measurements and umu test assays on airborne particles that were collected in forest and suburban sites. This was in order to gather information on transboundary pollution and its sources.

2. Materials and Methods

2.1. Sample Collection

The aerosol particles were collected at the forest site and at the suburban site in Nagasaki, Japan, as shown in Figure 1. Nagasaki is located on the western end of Japan. The forest site (32°54′32.4″ N, 129°44′33.5″ E, 533 m above sea level) was in Nagasaki Prefectural Forest Park (18 km northwest from Nagasaki city center), near the top of the Miharashi mountain in Nishisonogi peninsula and covered by trees. The peninsula facing the East China Sea has steep terrain, few flats, and low population density. The site is comparatively isolated from anthropogenic sources of pollution. The suburban

site (32°47′07.9″ N, 129°51′52.6″ E, 20 m above sea level) was located at Bunkyo Campus, Nagasaki University. Sampling was conducted on the roof of a four-storied campus building. The campus was surrounded by residential and business areas in Nagasaki city, which has a population of 0.4 million people. The suburban site is 5 km north of the industrial Nagasaki Port area.

Quartz fiber filters (254 mm × 203 mm, QR-100, Advantec, Tokyo, Japan) were preheated at 800 °C for 8 h and weighed after drying in a desiccator for >24 h. The quartz fiber filters were used for collecting aerosols with a high-volume sampler (HV-7000F, Sibata Scientific Technology Ltd., Saitama, Japan). The total suspended aerosol particles were collected, without size selection, by operating the high-volume sampler at a flow rate of 700 L/min. Collection of the aerosol particles were carried out three or four times in each season from 2012 to 2018, in spring (March–May), summer (June–August), fall (September–November), and winter (December–February). The sampling time was 1 week. Rainy days were avoided as much as possible when selecting sampling periods.



Figure 1. Location of the sampling sites.

2.2. PAHs Analysis

Fifteen types of PAHs, including acenaphthylene (Acy), acenaphthene (Ace), fluorene (Flu), phenanthrene (Phe), anthracene (Ant), fluoranthene (Flt), pyrene (Pyr), benzo[a]anthracene (BaA), chrysene (Chr), benzo[b]fluoranthene (BbF), benzo[k]fluoranthene (BkF), benzo[a]pyrene (BaP), dibenz[a,h]anthracene (DahA), indeno [1,2,3-cd]pyrene (IcdP), and benzo[ghi]perylene (BghiP) were selected for quantification in this study with reference to the 16 Priority Pollutant PAHs, as designated by the Environmental Protection Agency (EPA), excluding naphthalene. These standard reagents, apart from Ant, were purchased from AccuStandard, Inc. (New Haven, CT, USA). The Ant was purchased from Nacalai Tesque, Inc. (Kyoto, Japan). Acenaphthene-d₁₀, phenanthrene-d₁₀, pelyrene-d₁₂, and pyrene-d₁₀ were used as surrogate compounds for precise measurement. The former three were purchased from Wako Pure Chemical Industries, Ltd. (Osaka, Japan). Pyrene-d₁₀ was purchased from Kanto Chemical CO., Inc. (Tokyo, Japan). Silica gel C-200 was purchased from Wako Pure Chemical Industries, Ltd. for cleanup, and the analytical-grade solvents were purchased from Nacalai Tesque, Inc.

Filters were kept in a desiccator for several days, and then the filters were weighed and stored at -20 °C until extraction. The filter was chopped and put in a centrifuge tube, and then 100 ng of each of the four aforementioned surrogates (internal recovery standards) were added to it. Thirty milliliters of acetone was added to this, and ultrasonic extraction was performed for 10 min. After centrifugation, the extract was filtered through a cartridge with a pore size of 0.45 µm. This operation was repeated three times. The filtrate was concentrated to about 1 mL by nitrogen blowing. The

solution was fractionated using 5% H₂O deactivated silica gel column chromatography. PAHs were eluted with 50 mL of an acetone-dichloromethane mixture (3:2, v/v).

The eluted solution was dehydrated using anhydrous sodium sulfate and concentrated to 1 mL for analysis by nitrogen blowing. PAHs were determined with a 7890A gas chromatograph (Agilent Technologies, Santa Clara, CA, USA) equipped with a 7000A mass spectrometer (Agilent Technologies). One microliter aliquots were injected onto an HP-1MS capillary column (30 m × 0.25-mm i.d., 0.25 μ m) with 1.4 mL/min hexane gas. Injector temperature was kept at 300 °C. The column temperature was programmed as follows: 60 °C for 1 min, heat to 300 °C at 10 °C/min, and hold for 5 min. Quantification was conducted in selected ion monitoring (SIM) mode using the molecular weight of each PAH and deuterium-labelled PAH.

2.3. Umu Test

After collecting airborne particles, the filter was cut into small pieces, placed in a centrifuge tube and ultrasonically extracted with 30 mL of acetone for 10 min. Extraction, centrifugation, and filtration were repeated, as in the PAH analysis. Then, after the filtrate was almost dried by nitrogen blowing, 500 µL of Dimethyl sulfoxide (DMSO) was added to prepare the umu test solution.

The umu test performed as per the ISO 13829 with slight changes in the overnight culture conditions and the DMSO concentration of the test solution [65,66]. In this umu test using microplates, the test bacteria (*Salmonella typhimurium* TA1535/pSK1002) was brought into contact with a test solution to cause DNA damage. Then, the enzyme induced during the DNA repair processes was reacted with the chromogenic substrate. The intensity of DNA damage was measured from the amount of color development. The bacterial strain was provided by the Environmental Safety Management Laboratory, Faculty of Environment and Information Sciences, Yokohama National University, Japan. The S9 enzyme, obtained from Oriental Yeast Co., Ltd. (Tokyo, Japan), was used to determine genotoxicity in the presence of metabolic activation. The composition of S9 solution was 1.1% to the test solution described in the previous paragraph. The tests without and with the addition of the S9 solution are termed as the "–S9 test" and the "+S9 test", respectively. Positive control tests were performed each time. As a positive control substance, 4-nitroquinoline-1-oxide (4NQO) was used in the –S9 test, and 2-aminoanthracene (2AA) was used in the +S9 test. The tests were performed using four dose levels in triplicate for each level.

Induction ratio (IR) values were calculated following the ISO 13829. A scatter graph of added amount of air sample (horizontal axis) was plotted against the IR (vertical axis). The slope of the regression line was defined as genotoxicity. To correct for variations in the test sensitivity, genotoxicity was divided by the results of the positive control tests and represented as positive control equivalents, as previously reported [67,68].

2.4. Data Analysis

2.4.1. Back Trajectory Analysis

In order to grasp the long-range transport of air masses, back trajectory analysis is performed in many studies [39–41]. In this study, we used HYSPLIT Model-Trajectory Frequencies of the National Oceanic and Atmospheric Administration, USA [69,70]. Results from this model indicate the frequency that the trajectory passed over a grid cell. To investigate the typical transport behavior of air mass in each season, the analysis was conducted for 32 days in the middle of each season. This was the maximum number of days that could be set. The main analysis conditions were as follows: forest site source location; 32 days allowed to calculate trajectory frequencies; trajectory frequency grid resolution was set to 2.0 degree; trajectory starting interval was set to 12 h; and total run time was set to 72 h.

2.4.2. PAH Source Identification

Since relationships are thought to exist between the type of source and the PAH composition, the source's characteristics were estimated from the diagnostic ratio of PAHs. For the estimation,

5 of 16

various diagnostic ratios have been proposed [27–29]. In this study, Flt/(Flt+Pyr) and IcdP/(IcdP+BghiP) were selected as indicators that use PAHs with a relatively large number of rings and can distinguish three types of sources, namely, petroleum, petroleum combustion, and biomass/coal combustion. For example, when PAH Flt/(Flt+Pyr) levels are <0.4, the PAHs are thought to have petroleum sources; levels of 0.4–0.5 are thought to have come from liquid fossil fuel combustion pollution; and levels >0.5 indicate the grass, wood or coal combustion sources; levels of 0.2–0.5 are thought to have petroleum sources; levels of 0.2–0.5 are thought to have petroleum sources; levels of 0.2–0.5 are thought to have come from liquid fossil fuel combustion pollution; and levels >0.5 indicate the grass, wood or coal combustion sources; levels of 0.2–0.5 are thought to have come from liquid fossil fuel combustion pollution; and levels >0.5 indicate the grass, wood or coal combustion sources; levels of 0.2–0.5 are thought to have come from liquid fossil fuel combustion pollution; and levels >0.5 indicate the grass, wood or coal combustion sources; levels of 0.2–0.5 are thought to have come from liquid fossil fuel combustion pollution; and levels >0.5 indicate the grass, wood or coal combustion sources [71,72].

3. Results and Discussions

3.1. Back Trajectory Analysis

Figure 2 shows back trajectory analysis results for the forest site in 2018. These results were used to investigate seasonal trends relating to air mass advection. In spring, advection from the northwest was dominant, while some advection from the south side was also observed. In summer, many trajectory lines extended to the southeast side of Japan. In fall, there were many advections from the north and/or northwest sides. In winter, advection from the northwest was clearly dominant. In other words, air masses from the spring, fall, and winter seasons were transported from continental regions, such as northern China and the Korean peninsula, to the forest sampling site. Winter appeared to be strongly influenced by air masses that traveled through the continental region. Summer, on the other hand, was strongly influenced by air masses from the continental region. Although not shown in Figure 2, similar trends were seen in other years.



Figure 2. Frequency with which the back trajectories from the sampling site passed over a grid cell calculated by HYSPLIT Model of the National Oceanic and Atmospheric Administration, USA: (**a**) 29 March 2018–3 May 2018; (**b**) 29 June 2018–3 August 2018; (**c**) 29 September 2018–3 November 2018; (**d**) 30 December 2017–3 February 2018.

3.2. Concentration of PAHs and Genotoxicity

Table 1 shows PAH concentration and genotoxicity measurements at both sampling sites over 6 years, from 2012-2018. The total concentration of the 15 PAHs at the forest and suburban sites were $1.495 \pm 1.570 \text{ ng/m}^3$ and $1.996 \pm 2.209 \text{ ng/m}^3$, respectively. Although the measurement target was slightly different, it tended to be about two orders of magnitude lower than the previous reports on highly-polluted areas [73,74]. The increased amount of traffic at the suburban site is thought to give it a higher PAH concentration, compared to the forest site. Comparing each season, the values in spring, fall and winter were 2.4, 2.5 and 7.9 times higher than in summer at the forest site. At the suburban site, the values in spring, fall and winter were 3.8, 3.7 and 12.3 times higher than in summer. Both sites had the following relationship: winter > fall \Rightarrow spring > summer. The tendency to be higher in winter and lower in summer was consistent with the previous report [56-58]. Higher concentrations in winter might be explained by (1) increased burning of coal and biomass for thermal heating, especially in the cold northern parts of China; (2) the prevailing winter wind blowing from northwest by the stable atmospheric pressure chart pattern; (3) lower altitude for mixing of atmospheric zones in winter; (4) lower decomposition of chemical substances due to low temperatures and weak sunshine in winter; and (5) breakthrough of the PAHs with high vapor pressure from particulate phase to the gas phase due to high temperatures in the 1-week summer sampling period.

The genotoxicity of samples from the forest site (total average) was 0.16 ± 0.16 ng -4NQO/m³ for -S9 and 0.65 \pm 0.70 ng-2AA/m³ for +S9. The genotoxicity of samples from the suburban site was 0.16 ± 0.12 ng-4NQO/m³ for -S9 and 0.84 ± 0.74 ng-2AA/m³ for +S9. Comparing the two sites, the suburban site was higher in the +S9 condition. The main reason for this would be the higher levels of traffic around the suburban site, as in the case of PAHs. However, in the -S9 condition, the genotoxicity of samples from both sites were almost the same. The reason for this cannot be clarified at this time. We hope to investigate this further in the future. Since ±S9 values were converted to different positive control substances, the values cannot be compared directly. Comparing each season, the values in spring, fall and winter were 1.9, 3.5 and 9.8 times higher than in summer at the forest site in the -S9 condition, and 1.0, 2.5, and 7.6 times higher in the +S9 condition. At the suburban site, the values in spring, fall and winter were 2.6, 3.5 and 6.5 times higher than in summer in the -S9 condition, and 2.8, 5.8, and 12.9 times higher in the +S9 condition. Interestingly, these seasonal ratios were similar for both genotoxicity and PAH concentration. In all conditions, it was generally winter > fall > spring > summer. The tendency to be higher in winter and lower in summer was consistent with previous Ames test results on atmospheric genotoxicity [75–77]. Higher genotoxicity in winter might be explained in the same way as the PAH observations. This study would be the first longitudinal report on the atmospheric genotoxicity umu test results.

3.3. Seasonal Fluctuation

Figure 3 shows the results of continuous PAH concentration measurements and genotoxicity. Although the suburban site seemed to have higher PAH levels in the winter of 2018, there was no notable increase or decrease over the 6-year study duration. Both sites show higher PAH concentration and genotoxicity values in the winter, as indicated by the gray part of the figure. PAH concentration and genotoxicity showed the same seasonal patterns of increased levels in winter and lower levels in summer throughout the 6-year study period. This means that the PAH concentration and genotoxicity were correlated. Although they included the several effects that elevated winter concentrations and lowered summer concentrations, such as phase transition of some light PAHs, this correlation suggests that PAHs and genotoxic substances share common sources. Specifically, since there were several plots with different fluctuation trends in values even for the same sample, it was considered that there could be additional sources of genotoxic substances, other than PAHs. Despite seasonal changes in wind direction, the suburban site had almost the same seasonal fluctuation pattern and similar quantitative values as the forest site. This suggested that these areas might be affected by long-range transported pollutants, especially in winter.

Parameters	Forest site					Suburban				
	Total average	Spring	Summer	Fall	Winter	Total average	Spring	Summer	Fall	Winter
Concentration of										
PAHs (ng/m ³)	(n = 95)	(n = 21)	(n = 25)	(n = 26)	(n = 23)	(n = 97)	(n = 22)	(n = 25)	(n = 26)	(n = 24)
Асу	0.012 ± 0.021	0.004 ± 0.002	0.005 ± 0.008	0.007 ± 0.007	0.034 ± 0.034	0.024 ± 0.105	0.005 ± 0.003	0.003 ± 0.003	0.010 ± 0.012	$\begin{array}{c} 0.079 \pm \\ 0.203 \end{array}$
Ace	0.003 ± 0.005	0.001 ± 0.001	0.004 ± 0.007	0.004 ± 0.006	0.002 ± 0.002	0.004 ± 0.006	0.001 ± 0.001	0.005 ± 0.008	0.004 ± 0.008	0.003 ± 0.003
Flu	0.032 ± 0.086	0.006 ± 0.005	0.024 ± 0.082	0.009 ± 0.008	0.090 ± 0.140	0.046 ± 0.119	0.042 ± 0.151	0.005 ± 0.007	0.017 ± 0.036	0.125 ± 0.165
Phe	0.182 ± 0.257	0.101 ± 0.063	0.027 ± 0.037	0.081 ± 0.077	0.538 ± 0.302	0.198 ± 0.301	0.135 ± 0.082	0.046 ± 0.045	0.095 ± 0.070	0.526 ±
Ant	0.035 ± 0.162	0.015 ± 0.039	0.005 ± 0.008	0.018 ± 0.023	0.104 ± 0.321	0.047 ± 0.170	0.032 ± 0.074	0.013 ± 0.023	0.018 ± 0.025	0.129 ±
Flt	0.273 ± 0.322	0.204 ± 0.127	0.053 ± 0.053	0.198 ± 0.194	0.662 ± 0.399	0.342 ± 0.413	0.275 ± 0.155	0.047 ± 0.041	0.203 ± 0.157	0.861 ±
Pyr	0.177 ± 0.194	0.125 ± 0.063	0.054 ± 0.095	0.133 ± 0.127	0.409 ± 0.225	0.211 ± 0.229	0.173 ± 0.079	0.049 ± 0.047	0.141 ± 0.123	0.489 ±
BaA	0.065 ± 0.076	0.045 ± 0.025	0.021 ± 0.031	0.061 ± 0.061	0.136 ± 0.103	0.124 ± 0.368	0.093 ± 0.087	0.017 ± 0.014	0.061 ± 0.061	0.231 0.333 ±
Chr	0.168 ± 0.194	0.122 ± 0.084	0.034 ± 0.038	0.148 ± 0.134	0.378 ± 0.252	0.245 ± 0.296	0.178 ± 0.135	0.036 ± 0.031	0.248 ± 0.209	0.521 ±
BbF*	0.268 ± 0.390	0.161 ± 0.125	0.081 ± 0.063	0.167 ± 0.182	0.705 ± 0.600	0.467 ± 0.647	0.260 ± 0.248	0.081 ± 0.073	0.387 ± 0.429	0.412 1.173 ± 0.895

BF*	0.051 + 0.069	0.047 ± 0.063	0.026 ± 0.037	0.055 ± 0.037	0.079 ± 0.113	0 068 + 0 081	0.054 ± 0.046	0.018 ± 0.016	0.065 ± 0.057	$0.140 \ \pm$
DKI	0.001 ± 0.009	0.047 ± 0.005	0.020 ± 0.037	0.055 ± 0.057	0.077 ± 0.115	0.000 ± 0.001	0.034 ± 0.040	0.018 ± 0.010	0.005 ± 0.057	0.119
BaP	0.097 ± 0.097	0.088 ± 0.086	0.035 ± 0.043	0.098 ± 0.090	0.173 ± 0.108	0.094 ± 0.098	0.087 ± 0.052	0.035 ± 0.053	0.095 ± 0.074	$0.159\pm$
										0.144
Dah	0.022 + 0.054	0.022 + 0.058	0.017 ± 0.027	0.014 + 0.011	0.029 + 0.095	0.010 ± 0.025	0.015 + 0.014	0.005 ± 0.007	0.022 ± 0.010	$0.035 \pm$
IcdP	0.023 ± 0.034 0.092 ± 0.109	0.023 ± 0.038 0.069 ± 0.058	0.017 ± 0.037 0.044 ± 0.057	0.014 ± 0.011 0.095 ± 0.154	0.038 ± 0.083 0.163 ± 0.094	0.019 ± 0.023 0.117 ± 0.128	0.013 ± 0.014 0.084 ± 0.069	0.003 ± 0.007 0.025 ± 0.034	0.022 ± 0.019 0.108 ± 0.085	0.038
										$0.251\pm$
										0.162
D. 1 'D	0.004 + 0.125	0.002 + 0.007	0.040 + 0.052	0.065 + 0.040	0.104 + 0.224	0.1.40 + 0.100	0.100 + 0.110	0.025 + 0.025	0.100 ± 0.020	$0.313 \pm$
BghiP	0.094 ± 0.135	0.083 ± 0.087	0.040 ± 0.032	0.003 ± 0.049	0.194 ± 0.224	0.140 ± 0.190	0.109 ± 0.119	0.033 ± 0.033	0.109 ± 0.089	0.286
	1 405 - 1 550	1 0 2 0 1 0 5 0 7	0.420 + 0.471	1 117 - 0 705	2 405 + 1 0 4	1.000 + 2.200	1 4(0 + 0 ((2	0.200 + 0.202	1 426 + 0.001	$4.761 \pm$
ΣPAHs	$1.495 \pm 1.5/0$	1.039 ± 0.587	0.439 ± 0.471	$1.11/\pm 0./95$	3.485 ± 1.864	1.996 ± 2.209	1.469 ± 0.062	0.388 ± 0.283	1.436 ± 0.991	2.721
Genotoxicity	(n = 81)	(n = 17)	(n = 19)	(n = 25)	(n = 20)	(n = 82)	(n = 17)	(n = 19)	(n = 24)	(n = 22)
-S9 (ng-	0.16 ± 0.16	0.076 ± 0.040	0.040 ± 0.043	$0.14 \pm$	$0.39\pm$	0.16 ± 0.12	$0.12 \pm$	0.046 ± 0.037	$0.16 \pm$	0.30 ± 0.11
4NQO/m ³)				0.062	0.18		0.048		0.11	
+S9 (ng-2AA/m ³)	0.65 ± 0.70	$0.21 \pm$	$0.21 \pm$	0.53 ± 0.44	1.6 ± 0.64	0.84 ± 0.74	0.39 ± 0.21	$0.14 \pm$	$0.81 \ \pm$	1.8 ± 0.44
		0.15	0.22					0.13	0.48	

All data represent mean ± SD.

* Since BbF and BkF had been measured as a total value at the beginning of this study, the amount of data for BbF and BkF was about 60% of other PAHs.



Figure 3. Seasonal fluctuation of PAH concentrations and genotoxicity. The bar graph represents the total PAH concentrations. The circle and triangle plots in the line graph show genotoxicity under –S9 and +S9 conditions, respectively. The gray areas indicate winter data: (**a**) values of the forest site; (**b**) values of the suburban site.

3.4. PAH Source Identification

To analyze the source of the collected PAHs, Flt/(Flt+Pyr) and IcdP/(IcdP+BghiP) values were calculated using the measured values from the forest site, as shown in Figure 4. The summer plots varied relatively widely, suggesting that they were influenced by various sources for example, biomass or coal combustion, petroleum combustion, or petroleum itself. Figure 5 shows the distribution of values for each season. This was produced by statistically-processing the values on both axes in Figure 4 after removing outliers. Values exceeding 1.5 times the interquartile range were defined as outliers. In either index, the summer values tended to be lower than the other seasons. In particular, for the ratio IcdP/(IcdP+BghiP), the median value was in the area of liquid fossil fuel combustion only in summer. Considering the back trajectory analysis results, the influence of air masses from the Pacific Ocean side would be significant in summer. Ships might be a considerable source of summer PAHs. Compared to the IcdP/(IcdP+BghiP) ratio in summer, it was shown that the plots in spring, fall and winter shifted to the biomass or coal combustion side, with some variation. This may be due to the fact that PAHs would be generated from coal heating in the cold continental regions of northern China and the Korean peninsula. This is consistent with the back trajectory analysis results. According to a previous report, central heating systems are widely used in cities in north of China from October to April [78], including the spring and fall period of this study. Therefore, even in the spring and fall, the plots were considered to have shifted to the biomass or coal combustion side. The IcdP/(IcdP+BghiP) ratio in spring and fall were not much different from winter. This suggests that IcdP and BghiP were predominantly derived from biomass or coal combustion, rather than from petroleum combustion. Since the fluctuation patterns are correlated, as shown in Figure 3, it is suggested that PAHs and genotoxic substances share common sources. Therefore, biomass or coal combustion in the continental region during cold weather (especially in winter) would also be the main origin of genotoxic substances.



Figure 4. Cross plot of the diagnostic ratios of the forest site for PAH sources in each season.



Figure 5. Ranges of the diagnostic ratios of the forest site for each season, excluding outliers. The upper and lower ends of the bars represent the upper and lower limits of the observed values. The upper and lower ends of the boxes represent the third and first quartiles. The lines intersecting the boxes represents the median values: (**a**) ranges of IcdP/(IcdP+BghiP); (**b**) ranges of Flt/(Flt+Pyr).

3.5. Contribution of Transboundary Pollution.

At present, the impact of transboundary pollution cannot accurately be quantitatively evaluated. However, it is important to have an image of the magnitude of transboundary pollution effects. In order to estimate the extent of transboundary pollution, the forest and suburban sites were compared using the concentration of PAHs in winter (6-year average), when the impact of transboundary pollution was expected to increase. From Table 1, the concentration of PAHs at the forest and suburban sites were 3.485 ng/m³ and 4.761 ng/m³, respectively. Assuming that all the PAHs at the forest site were derived from transboundary pollution flying directly into the suburban site. Based on a three-dimensional chemical transport model, Ikeda et al. estimated relative contribution ratios of Northeast China, Central North China, Central South China, Korean Peninsula, and Japan to PM2.5 in the winter on Fukue Island, which is a remote island about 100 km west from the forest site. The relative contribution ratios were reported as 7.3%, 60.6%, 29.2%, 11.8%, and 1.6%, respectively [79]. Therefore, the contribution rate of 70% obtained in this study would not be unreasonable. Since the impact of transboundary pollution is considered to be at a level that cannot be ignored, it is necessary to continue related research and make international efforts to solve the problem as soon as possible.

4. Conclusions

- Throughout the 6 years studied, there was no notable increase or decrease in PAH concentration
 or genotoxicity at either the forest or suburban sites. This indicated that the air pollution around
 these sites caused by PAHs or genotoxic substances did not improve or worsen during this
 period.
- In terms of PAH concentration and genotoxicity, the suburban and forest sites shared similar seasonal fluctuation patterns and quantitative values, suggesting that the western end of Japan might be affected by long-range transported pollutants, especially in winter.
- Back trajectory analysis results strongly suggested the influence of transboundary pollution from continental regions, such as northern China and the Korean peninsula, especially in winter. Combined with analysis of the diagnostic ratios of PAHs, coal or biomass combustion in the continental region was considered one of the main sources of atmospheric PAHs at the sampling site.
- PAH concentration and genotoxicity were correlated, showing the same seasonal patterns of
 increased levels in winter and lower levels in summer throughout the 6-year study period. It
 was considered that PAHs and genotoxic substances share common sources. This suggested that

transboundary pollutants caused by coal and biomass combustion from the continental region were major sources of genotoxicity.

• It was estimated that the contribution of the transboundary PAH pollution at the suburban site could reach up to 70%.

Author Contributions: Conceptualization, Y.T., M.N. and T.K.; methodology, Y.T. and T.K.; formal analysis, T.K.; investigation, W.B.; Writing—Original draft preparation, T.K.; Writing—Review and editing, Y.T. and T.K.; supervision, Y.T.; funding acquisition, M.N.

Funding: This research was supported in part by JSPS KAKENHI Grant Number JP 24404007.

Acknowledgments: The authors gratefully acknowledge the NOAA Air Resources Laboratory (ARL) for the provision of the HYSPLIT transport and dispersion model and/or READY website (https://www.ready.noaa.gov) used in this publication. The authors would like to acknowledge professor Takashi Kameya, Faculty of Environment and Information Sciences, Yokohama National University, Japan for providing tester strain. The authors also acknowledge all members of Faculty of Environmental Science, Nagasaki University, Japan involved in this study.

Conflicts of Interest: The authors declare no conflict of interest.

References

- 1. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans;* International Agency for Research on Cancer: Lyon, France, 2010; Volume 92.
- Kubo, T.; Urano, K.; Utsumi, H. Mutagenicity characteristics of 255 environmental chemicals. J. Health Sci. 2002, 48, 545–554.
- 3. Fernandes, M.B.; Sicre, M.A.; Boireau, A.; Tronczynski, J. Polyaromatic hydrocarbon (PAH) distributions in the Seine River and its estuary. *Mar. Pollut. Bull.* **1997**, *34*, 857–867.
- 4. Shi, Z.; Tao, S.; Pan, B.; Fan, W.; He, X.C.; Zuo, Q.; Wu, S.P.; Li, B.G.; Cao, J.; Liu, W.X.; et al. Contamination of rivers in Tianjin, China by polycyclic aromatic hydrocarbons. *Environ. Pollut.* **2005**, *134*, 97–111.
- Wang, Y.; Zhang, S.; Cui, W.; Meng, X.; Tang, X. Polycyclic aromatic hydrocarbons and organochlorine pesticides in surface water from the Yongding River basin, China: Seasonal distribution, source apportionment, and potential risk assessment. *Sci. Total Environ.* 2018, *618*, 419–429.
- Zakaria, M.P.; Takada, H.; Tsutsumi, S.; Ohno, K.; Yamada, J.; Kouno, E.; Kumata, H. Distribution of polycyclic aromatic hydrocarbons (PAHs) in rivers and estuaries in Malaysia: A widespread input of petrogenic PAHs. *Environ. Sci. Technol.* 2002, *36*, 1907–1918.
- Mai, B.X.; Fu, J.M.; Sheng, G.Y.; Kang, Y.H.; Lin, Z.; Zhang, G.; Min, Y.S.; Zeng, E.Y. Chlorinated and polycyclic aromatic hydrocarbons in riverine and estuarine sediments from Pearl River Delta, China. *Environ. Pollut.* 2002, 117, 457–474.
- 8. Edokpayi, J.; Odiyo, J.; Popoola, O.; Msagati, T. Determination and distribution of polycyclic aromatic hydrocarbons in rivers, sediments and wastewater effluents in Vhembe District, South Africa. *Int. J. Environ. Res. Public Health* **2016**, *13*, 387.
- 9. Bucheli, T.D.; Blum, F.; Desaules, A.; Gustafsson, Ö. Polycyclic aromatic hydrocarbons, black carbon, and molecular markers in soils of Switzerland. *Chemosphere* **2004**, *56*, 1061–1076.
- 10. Peng, C.; Chen, W.; Liao, X.; Wang, M.; Ouyang, Z.; Jiao, W.; Bai, Y. Polycyclic aromatic hydrocarbons in urban soils of Beijing: Status, sources, distribution and potential risk. *Environ. Pollut.* **2011**, *159*, 802–808.
- 11. Gereslassie, T.; Workineh, A.; Liu, X.; Yan, X.; Wang, J. Occurrence and ecological and human health risk assessment of polycyclic aromatic hydrocarbons in soils from Wuhan, central China. *Int. J. Environ. Res. Public Health* **2018**, *15*, 2751.
- 12. Guo, H.; Lee, S.C.; Ho, K.F.; Wang, X.M.; Zou, S.C. Particle-associated polycyclic aromatic hydrocarbons in urban air of Hong Kong. *Atmos. Environ.* **2003**, *37*, 5307–5317.
- 13. Albinet, A.; Leoz-Garziandia, E.; Budzinski, H.; Villenave, E. Polycyclic aromatic hydrocarbons (PAHs), nitrated PAHs and oxygenated PAHs in ambient air of the Marseilles area (south of France): Concentrations and sources. *Sci. Total Environ.* **2007**, *384*, 280–292.

- Yang, L.; Suzuki, G.; Zhang, L.; Zhou, Q.; Zhang, X.; Xing, W.; Shima, M.; Yoda, Y.; Nakatsubo, R.; Hiraki, T.; et al. The characteristics of polycyclic aromatic hydrocarbons in different emission source areas in Shenyang, China. *Int. J. Environ. Res. Public Health* **2019**, *16*, 2817.
- Crisafully, R.; Milhome, M.A.L.; Cavalcante, R.M.; Silveira, E.R.; De Keukeleire, D.; Nascimento, R.F. Removal of some polycyclic aromatic hydrocarbons from petrochemical wastewater using low-cost adsorbents of natural origin. *Bioresour. Technol.* 2008, *99*, 4515–4519.
- Qi, W.; Liu, H.; Qu, J.; Hu, C.; Lan, H.; Berg, M.; Ren, H.; Xu, W. Polycyclic aromatic hydrocarbons in effluents from wastewater treatment plants and receiving streams in Tianjin, China. *Environ. Monit. Assess.* 2011, 177, 467–480.
- 17. Qi, W.; Liu, H.; Pernet-Coudrier, B.; Qu, J. Polycyclic aromatic hydrocarbons in wastewater, WWTPs effluents and in the recipient waters of Beijing, China. *Environ. Sci. Pollut. Res.* **2013**, *20*, 4254–4260.
- Lee, W.J.; Liow, M.C.; Tsai, P.J.; Hsieh, L.T. Emission of polycyclic aromatic hydrocarbons from medical waste incinerators. *Atmos. Environ.* 2002, *36*, 781–790.
- Yang, H.H.; Hsieh, L.T.; Liu, H.C.; Mi, H.H. Polycyclic aromatic hydrocarbon emissions from motorcycles. Atmos. Environ. 2005, 39, 17–25.
- Borrás, E.; Tortajada-Genaro, L.A.; Vázquez, M.; Zielinska, B. Polycyclic aromatic hydrocarbon exhaust emissions from different reformulated diesel fuels and engine operating conditions. *Atmos. Environ.* 2009, 43, 5944–5952.
- Pham, C.T.; Boongla, Y.; Nghiem, T.D.; Le, H.T.; Tang, N.; Toriba, A.; Hayakawa, K. Emission characteristics of polycyclic aromatic hydrocarbons and nitro-polycyclic aromatic hydrocarbons from open burning of rice straw in the north of vietnam. *Int. J. Environ. Res. Public Health* 2019, *16*, 2343.
- Cattaneo, A.; Fermo, P.; Urso, P.; Perrone, M.G.; Piazzalunga, A.; Tarlassi, J.; Carrer, P.; Cavallo, D.M. Particulate-bound polycyclic aromatic hydrocarbon sources and determinants in residential homes. *Environ. Pollut.* 2016, 218, 16–25.
- Piazzalunga, A.; Anzano, M.; Collina, E.; Lasagni, M.; Lollobrigida, F.; Pannocchia, A.; Fermo, P.; Pitea, D. Contribution of wood combustion to PAH and PCDD/F concentrations in two urban sites in Northern Italy. *J. Aerosol Sci.* 2013, *56*, 30–40.
- Marr, L.C.; Kirchstetter, T.W.; Harley, R.A.; Miguel, A.H.; Hering, S.V.; Hammond, S.K. Characterization of polycyclic aromatic hydrocarbons in motor vehicle fuels and exhaust emissions. *Environ. Sci. Technol.* 1999, 33, 3091–3099.
- Zoccolillo, L.; Babi, D.; Felli, M. Evaluation of polycyclic aromatic hydrocarbons in gasoline by HPLC and GC-MS. *Chromatographia* 2000, *52*, 373–376.
- Zhao, Y.; Hong, B.; Fan, Y.; Wen, M.; Han, X. Accurate analysis of polycyclic aromatic hydrocarbons (PAHs) and alkylated PAHs homologs in crude oil for improving the gas chromatography/mass spectrometry performance. *Ecotoxicol. Environ. Saf.* 2014, 100, 242–250.
- Yunker, M.B.; Macdonald, R.W.; Vingarzan, R.; Mitchell, R.H.; Goyette, D.; Sylvestre, S. PAHs in the Fraser River basin: A critical appraisal of PAH ratios as indicators of PAH source and composition. *Org. Geochem.* 2002, 33, 489–515.
- Zhang, X.L.; Tao, S.; Liu, W.X.; Yang, Y.; Zuo, Q.; Liu, S.Z. Source diagnostics of polycyclic aromatic hydrocarbons based on species ratios: A multimedia approach. *Environ. Sci. Technol.* 2005, 39, 9109–9114.
- De La Torre-Roche, R.J.; Lee, W.Y.; Campos-Díaz, S.I. Soil-borne polycyclic aromatic hydrocarbons in El Paso, Texas: Analysis of a potential problem in the United States/Mexico border region. *J. Hazard. Mater.* 2009, 163, 946–958.
- Wirth, J.D. The Trail Smelter Dispute: Canadians and Americans Confront Transboundary Pollution, 1927– 41. In *Environmental History*; Oxford University Press: Oxford, UK, 1996; Volume 1, pp. 34–51.
- REPORT OF THE UNITED NATIONS CONFERENCE ON ENVIRONMENT AND DEVELOPMENT. Available online: https://sustainabledevelopment.un.org/content/documents/1709riodeclarationeng.pdf (accessed on 21 December 2019).
- UNITED NATIONS FRAMEWORK CONVENTIONON CLIMATE CHANGE. Available online: https://unfccc.int/files/essential_background/background_publications_htmlpdf/application/pdf/conveng .pdf (accessed on 21 December 2019).
- Johnson, A.; Norton, D.; Yake, B.; Twiss, S. Transboundary metal pollution of the Columbia River (Franklin D. Roosevelt Lake). *Bull. Environ. Contam. Toxicol.* **1990**, 45, 703–710.

- 34. Suleymanov, B.; Ahmedov, M.; Safarova, K.; Steinnes, E. Metals in main rivers of Azerbaijan: Influence of transboundary pollution. *Water Air Soil Pollut*. **2010**, *213*, 301–310.
- 35. Schudel, G.; Miserendino, R.A.; Veiga, M.M.; Velasquez-López, P.C.; Lees, P.S.J.; Winland-Gaetz, S.; Davée Guimarães, J.R.; Bergquist, B.A. An investigation of mercury sources in the Puyango-Tumbes River: Using stable Hg isotopes to characterize transboundary Hg pollution. *Chemosphere* 2018, 202, 777–787.
- 36. Berge, E.; Bartnicki, J.; Olendrzynski, K.; Tsyro, S.G. Long-term trends in emissions and transboundary transport of acidifying air pollution in Europe. *J. Environ. Manag.* **1999**, *57*, 31–50.
- Chang, K.H.; Jeng, F.T.; Tsai, Y.L.; Lin, P.L. Modeling of long-range transport on Taiwan's acid deposition under different weather conditions. *Atmos. Environ.* 2000, 34, 3281–3295.
- 38. Murano, K.; Mukai, H.; Hatakeyama, S.; Suk Jang, E.; Uno, I. Trans-boundary air pollution over remote islands in Japan: Observed data and estimates from a numerical model. *Atmos. Environ.* **2000**, *34*, 5139–5149.
- Moreno, T.; Kojima, T.; Amato, F.; Lucarelli, F.; De La Rosa, J.; Calzolai, G.; Nava, S.; Chiari, M.; Alastuey, A.; Querol, X.; et al. Daily and hourly chemical impact of springtime transboundary aerosols on Japanese air quality. *Atmos. Chem. Phys.* 2013, *13*, 1411–1424.
- Masiol, M.; Benetello, F.; Harrison, R.M.; Formenton, G.; De Gaspari, F.; Pavoni, B. Spatial, seasonal trends and transboundary transport of PM_{2.5} inorganic ions in the Veneto region (Northeastern Italy). *Atmos. Environ.* 2015, 117, 19–31.
- Itahashi, S.; Uno, I.; Osada, K.; Kamiguchi, Y.; Yamamoto, S.; Tamura, K.; Wang, Z.; Kurosaki, Y.; Kanaya, Y. Nitrate transboundary heavy pollution over East Asia in winter. *Atmos. Chem. Phys.* 2017, *17*, 3823–3843.
- Lunde, G.; Bjorseth, A. Polycyclic aromatic hydrocarbons in long-range transported aerosols. *Nature* 1977, 268, 518–519.
- Yagishita, M.; Kageyama, S.; Ohshima, S.; Matsumoto, M.; Aoki, Y.; Goto, S.; Nakajima, D. Atmospheric concentration and carcinogenic risk of polycyclic aromatic hydrocarbons including benzo[c]fluorene, cyclopenta[c,d]pyrene, and benzo[j]fluoranthene in Japan. *Atmos. Environ.* 2015, *115*, 263–268.
- 44. Odabasi, M.; Vardar, N.; Sofuoglu, A.; Tasdemir, Y.; Holsen, T.M. Polycyclic aromatic hydrocarbons (PAHs) in Chicago air. *Sci. Total Environ.* **1999**, 227, 57–67.
- Bari, M.A.; Baumbach, G.; Kuch, B.; Scheffknecht, G. Particle-phase concentrations of polycyclic aromatic hydrocarbons in ambient air of rural residential areas in southern Germany. *Air Qual. Atmos. Health* 2010, 3, 103–116.
- Singh, D.P.; Gadi, R.; Mandal, T.K. Characterization of particulate-bound polycyclic aromatic hydrocarbons and trace metals composition of urban air in Delhi, India. *Atmos. Environ.* 2011, 45, 7653– 7663.
- Vestenius, M.; Leppänen, S.; Anttila, P.; Kyllönen, K.; Hatakka, J.; Hellén, H.; Hyvärinen, A.P.; Hakola, H. Background concentrations and source apportionment of polycyclic aromatic hydrocarbons in southeastern Finland. *Atmos. Environ.* 2011, 45, 3391–3399.
- Chen, Y.C.; Chiang, H.C.; Hsu, C.Y.; Yang, T.T.; Lin, T.Y.; Chen, M.J.; Chen, N.T.; Wu, Y.S. Ambient PM₂₅bound polycyclic aromatic hydrocarbons (PAHs) in Changhua county, central Taiwan: Seasonal variation, source apportionment and cancer risk assessment. *Environ. Pollut.* **2016**, *218*, 372–382.
- Yang, L.; Tang, N.; Matsuki, A.; Takami, A.; Hatakeyama, S.; Kaneyasu, N.; Nagato, E.G.; Sato, K.; Yoshino, A.; Hayakawa, K. A comparison of particulate-bound polycyclic aromatic hydrocarbons long-range transported from the Asian continent to the Noto Peninsula and Fukue Island, Japan. *Asian J. Atmos. Environ.* 2018, 12, 369–376.
- Kakimoto, H.; Kitamura, M.; Matsumoto, Y.; Sakai, S.; Kanoh, F.; Murahashi, T.; Akutsu, K.; Kizu, R.; Hayakawa, K. Comparison of atmospheric polycyclic aromatic hydrocarbons and nitropolycyclic aromatic hydrocarbons in Kanazawa, Sapporo and Tokyo. J. Health Sci. 2000, 46, 5–15.
- Tang, N.; Hattori, T.; Taga, R.; Igarashi, K.; Yang, X.; Tamura, K.; Kakimoto, H.; Mishukov, V.F.; Toriba, A.; Kizu, R.; et al. Polycyclic aromatic hydrocarbons and nitropolycyclic aromatic hydrocarbons in urban air particulates and their relationship to emission sources in the Pan-Japan Sea countries. *Atmos. Environ.* 2005, 39, 5817–5826.
- Hayakawa, K.; Tang, N.; Nagato, E.G.; Toriba, A.; Sakai, S.; Kano, F.; Goto, S.; Endo, O.; Arashidani, K.; Kakimoto, H. Long term trends in atmospheric concentrations of polycyclic aromatic hydrocarbons and nitropolycyclic aromatic hydrocarbons: A study of Japanese cities from 1997 to 2014. *Environ. Pollut.* 2018, 233, 474–482.

- Nassar, H.F.; Tang, N.; Kameda, T.; Toriba, A.; Khoder, M.I.; Hayakawa, K. Atmospheric concentrations of polycyclic aromatic hydrocarbons and selected nitrated derivatives in Greater Cairo, Egypt. *Atmos. Environ.* 2011, 45, 7352–7359.
- 54. Chuesaard, T.; Chetiyanukornkul, T.; Kameda, T.; Hayakawa, K.; Toriba, A. Influence of biomass burning on the levels of atmospheric polycyclic aromatic hydrocarbons and their nitro derivatives in Chiang Mai, Thailand. *Aerosol Air Qual. Res.* **2014**, *14*, 1247–1257.
- Tang, N.; Suzuki, G.; Morisaki, H.; Tokuda, T.; Yang, X.; Zhao, L.; Lin, J.; Kameda, T.; Toriba, A.; Hayakawa, K. Atmospheric behaviors of particulate-bound polycyclic aromatic hydrocarbons and nitropolycyclic aromatic hydrocarbons in Beijing, China from 2004 to 2010. *Atmos. Environ.* 2017, *152*, 354–361.
- 56. Manoli, E.; Kouras, A.; Samara, C. Profile analysis of ambient and source emitted particle-bound polycyclic aromatic hydrocarbons from three sites in northern Greece. *Chemosphere* **2004**, *56*, 867–878.
- Li, Z.; Porter, E.N.; Sjödin, A.; Needham, L.L.; Lee, S.; Russell, A.G.; Mulholland, J.A. Characterization of PM_{2.5}-bound polycyclic aromatic hydrocarbons in Atlanta–Seasonal variations at urban, suburban, and rural ambient air monitoring sites. *Atmos. Environ.* **2009**, *43*, 4187–4193.
- Rehwagen, M.; Müller, A.; Massolo, L.; Herbarth, O.; Ronco, A. Polycyclic aromatic hydrocarbons associated with particles in ambient air from urban and industrial areas. *Sci. Total Environ.* 2005, 348, 199– 210.
- Matsushita, H.; Endo, O.; Goto, S.; Shimizu, H.; Matsumoto, H.; Tamakawa, K.; Endo, T.; Sakabe, Y.; Tokiwa, H.; Ando, M. Collaborative study using the preincubation Salmonella typhimurium mutation assay for airborne particulate matter in Japan. A trial to minimize interlaboratory variation. *Mutat. Res. Environ. Mutagen. Relat. Subj.* 1992, 271, 1–12.
- De Aragão Umbuzeiro, G.; Roubicek, D.A.; Rech, C.M.; Sato, M.I.Z.; Claxton, L.D. Investigating the sources of the mutagenic activity found in a river using the Salmonella assay and different water extraction procedures. *Chemosphere* 2004, 54, 1589–1597.
- Traversi, D.; Degan, R.; De Marco, R.; Gilli, G.; Pignata, C.; Villani, S.; Bono, R. Mutagenic properties of PM2.5 urban pollution in the Northern Italy: The nitro-compounds contribution. *Environ. Int.* 2009, 35, 905– 910.
- 62. Wang, L.S.; Hu, H.Y.; Wang, C. Effect of ammonia nitrogen and dissolved organic matter fractions on the genotoxicity of wastewater effluent during chlorine disinfection. *Environ. Sci. Technol.* **2007**, *41*, 160–165.
- Kubo, T.; Wang, L.; Kameya, T.; Kobayashi, T.; Hu, H.; Urano, K.; Shigeoka, T. Quantitative evaluation of the genotoxic activity of Japanese tap water using the umu test. J. Water Environ. Technol. 2015, 13, 291–300.
- 64. Chai, Q.; Hu, A.; Qian, Y.; Ao, X.; Liu, W.; Yang, H.; Xie, Y.F. A comparison of genotoxicity change in reclaimed wastewater from different disinfection processes. *Chemosphere* **2018**, *191*, 335–341.
- International Standard Organisation. Water quality—Determination of the Genotoxicity of Water and Waste Water Using the Umu-Test, 1st ed.; ISO 13829; International Standard Organisation: Geneva, Switzerland, 2000; pp. 1–18.
- 66. Sugimura, Y.; Kubo, T.; Murata, N.; Urano, K.; Shigeoka, T. Development of the efficient umu-test system for evaluation of genotoxicity of water samples. *Jpn. J. Environ. Toxicol.* **2008**, *11*, 21–32.
- 67. Kameya, T.; Nagato, T.; Nakagawa, K.; Yamashita, D.; Kobayashi, T.; Fujie, K. Quantification of *umu* genotoxicity level of urban river water. *Water Sci. Technol.* **2011**, *63*, 410–415.
- Bai, W.; Takao, Y.; Kubo, T. Evaluation of genotoxicity potential of household effluents from onsite wastewater treatment systems using *umu* test. J. Toxicol. Environ. Health Part A 2020, doi:10.1080/15287394.2020.1719447.
- 69. Stein, A.F.; Draxler, R.R.; Rolph, G.D.; Stunder, B.J.B.; Cohen, M.D.; Ngan, F. Noaa's hysplit atmospheric transport and dispersion modeling system. *Bull. Am. Meteorol. Soc.* **2015**, *96*, 2059–2077.
- Rolph, G.; Stein, A.; Stunder, B. Real-time Environmental Applications and Display sYstem: READY. Environ. Model. Softw. 2017, 95, 210–228.
- Tobiszewski, M.; Namieśnik, J. PAH diagnostic ratios for the identification of pollution emission sources. Environ. Pollut. 2012, 162, 110–119.
- Huang, L.; Chernyak, S.M.; Batterman, S.A. PAHs (polycyclic aromatic hydrocarbons), nitro-PAHs, and hopane and sterane biomarkers in sediments of southern Lake Michigan, USA. *Sci. Total Environ.* 2014, 487, 173–186.

- Shi, G.L.; Feng, Y.C.; Wu, J.H.; Li, X.; Wang, Y.Q.; Xue, Y.H.; Zhu, T. Source identification of polycyclic aromatic hydrocarbons in urban particulate matter of Tangshan, China. *Aerosol Air Qual. Res.* 2009, *9*, 309– 315.
- Byambaa, B.; Yang, L.; Matsuki, A.; Nagato, E.G.; Gankhuyag, K.; Chuluunpurev, B.; Banzragch, L.; Chonokhuu, S.; Tang, N.; Hayakawa, K. Sources and characteristics of polycyclic aromatic hydrocarbons in ambient total suspended particles in Ulaanbaatar City, Mongolia. *Int. J. Environ. Res. Public Health* 2019, 16, 442.
- Matsumoto, Y.; Sakai, S.; Kato, T.; Nakajima, T.; Satoh, H. Long-term trends of particulate mutagenic activity in atmosphere of Sapporo. 1. Determination of mutagenic activity by the conventional tester strains TA98 and TA100 during an 18-year period (1974-1992). *Environ. Sci. Technol.* 1998, 32, 2665–2671.
- Zwodziak, J.; Jadczyk, P.; Kucharczyk, J. Seasonal variability of the mutagenicity of airborne particles in the town center. J. Aerosol Sci. 2001, 32, 409–423.
- Coulibaly, S.; Minami, H.; Abe, M.; Hasei, T.; Sera, N.; Yamamoto, S.; Funasaka, K.; Asakawa, D.; Watanabe, M.; Honda, N.; et al. Seasonal fluctuations in air pollution in Dazaifu, Japan, and effect of long-range transport from mainland East Asia. *Biol. Pharm. Bull.* 2015, *38*, 1395–1403.
- Yang, X.Y.; Okada, Y.; Tang, N.; Matsunaga, S.; Tamura, K.; Lin, J.M.; Kameda, T.; Toriba, A.; Hayakawa, K. Long-range transport of polycyclic aromatic hydrocarbons from China to Japan. *Atmos. Environ.* 2007, 41, 2710–2718.
- Ikeda, K.; Yamaji, K.; Kanaya, Y.; Taketani, F.; Pan, X.; Komazaki, Y.; Kurokawa, J.; Ohara, T. Sensitivity analysis of source regions to PM_{2.5} concentration at Fukue Island, Japan. *J. Air Waste Manag. Assoc.* 2014, 64, 445–452.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).