

Table S1. Mean scores, standard deviations and one way analysis of variance results accounted for by gender and country of residence.

Gender	Traumatic distress (M, SD)	Anxiety (M, SD)	Depression (M, SD)	Resilience (M, SD)	Hope (M, SD)
Female	16.1 (14.8)	5.57 (7.04)	8.99 (9.10)	71.8 (10.0)	50.6 (9.02)
Male	12.7 (14.1)	3.67 (5.02)	7.20 (7.80)	70.1 (10.0)	52.1 (7.69)
ANOVA ($n=455$)	$F(1, 454) = 6.06, p = 0.014, \eta^2 = 0.01$	$F(1, 454) = 9.50, p = 0.002, \eta^2 = 0.02$	$F(1, 454) = 4.53, p = 0.03, \eta^2 = 0.009$	$F(1, 455) = 0.76, p = 0.38, \eta^2 = 0.001$	$F(1, 455) = 3.15, p = 0.08, \eta^2 = 0.006$
Country of residence					
Ireland	18.7 (16.2)	6.24 (7.3)	11.1 (9.59)	69.2 (10.6)	48.4 (9.12)
U.S.	11.9 (12.4)	3.85 (5.45)	6.19 (7.29)	73.1 (9.27)	53.1 (7.62)
ANOVA ($n = 456$)	$F(1, 455) = 26.3, p < 0.001, \eta^2 = 0.05$	$F(1, 455) = 15.9, p < 0.001, \eta^2 = 0.03$	$F(1, 455) = 39.1, p < 0.001, \eta^2 = 0.07$	$F(1, 455) = 17.0, p < 0.001, \eta^2 = 0.03$	$F(1, 455) = 35.4, p < 0.001, \eta^2 = 0.07$

M, mean scores; SD, standard deviation; U.S., United States