

SUPPLEMENTARY INFORMATION

Efficient removal of nonylphenol isomers from water by use of organo-hydrotalcites

Daniel Cosano*, Dolores Esquivel, Francisco J. Romero-Salguero, César Jiménez-Sanchidrián and José Rafael Ruiz*

Departamento de Química Orgánica, Instituto Universitario de Investigación en Química Fina y Nanoquímica IUIQFN, Facultad de Ciencias, Universidad de Córdoba, Campus de Rabanales, Edificio Marie Curie, E-14071 Córdoba, Spain

*Corresponding authors. E-mail address: q92cohid@uco.es (D. Cosano); qo1ruarj@uco.es (J.R. Ruiz), Tel.: 34 957218638; Fax: 34 957212066.

Figure S1. Kinetic models for 15 ppm concentration: a) Pseudo-first-order kinetic model, b) Pseudo-second-order kinetic model, c) Elovich kinetic model, d) and Interparticle diffusion kinetic model.




Figure S2. Kinetic models for 25 ppm concentration: a) Pseudo-first-order kinetic model, b) Pseudo-second-order kinetic model, c) Elovich kinetic model, d) and Interparticle diffusion kinetic model.




Figure S3. Kinetic models for 30 ppm concentration: a) Pseudo-first-order kinetic model, b) Pseudo-second-order kinetic model, c) Elovich kinetic model, d) and Interparticle diffusion kinetic model.

