



Article Big Five Traits as Predictors of a Healthy Lifestyle during the COVID-19 Pandemic: Results of a Russian Cross-Sectional Study

Alena Zolotareva ^{1,2,*}, Sergei Shchebetenko ¹, Svetlana Belousova ³, Irina Danilova ^{4,5}, Vadim Tseilikman ⁶, Maxim Lapshin ², Lilia Sarapultseva ⁷, Svetlana Makhniova ⁷, Maria Sarapultseva ^{5,8}, Maria Komelkova ², Desheng Hu ⁹, Shanshan Luo ¹⁰, Ekaterina Lisovskaya ³ and Alexey Sarapultsev ²

- ¹ School of Psychology, HSE University, 101000 Moscow, Russia
- ² Russian-Chinese Education and Research Center of System Pathology, South Ural State University, 454080 Chelyabinsk, Russia
- ³ Department of Psychology, Chelyabinsk State University, 454001 Chelyabinsk, Russia
- ⁴ Institute of Natural Sciences and Mathematics, Ural Federal University Named after the First President of Russia B.N. Yeltsin, 620002 Ekaterinburg, Russia
- ⁵ Institute of Immunology and Physiology, Ural Branch of the Russian Academy of Science, 620049 Ekaterinburg, Russia
- ⁶ School of Medical Biology, South Ural State University, 454080 Chelyabinsk, Russia
- ⁷ Department of Mathematics and Natural Sciences, Russian State Vocational Pedagogical University, 620012 Ekaterinburg, Russia
- ⁸ Department of Pediatric Dentistry, Medical Firm Vital EBB, 620144 Ekaterinburg, Russia
- ⁹ Department of Integrated Traditional Chinese and Western Medicine, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430022, China
- ¹ Institute of Hematology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430022, China
- Correspondence: alena.a.zolotareva@gmail.com

Abstract: The healthy lifestyle of people around the world has changed dramatically during the COVID-19 pandemic. The personality risk factors for these processes from around the world remain understudied. This study aimed to examine the associations of the Big Five traits with a healthy lifestyle during the COVID-19 pandemic. In a cross-sectional study, data from 1215 Russian university students were analyzed. Participants completed the Big Five Inventory-10 and Short Multidimensional Inventory Lifestyle Evaluation. The results showed that personality traits predicted many dimensions of a healthy lifestyle during the COVID-19 pandemic. Diet and nutrition were positively predicted by extraversion, agreeableness, and conscientiousness, and it was negatively predicted by neuroticism. Substance abuse was positively predicted by agreeableness and conscientiousness, and it was negatively predicted by extraversion. Physical activity was positively predicted by extraversion and conscientiousness, and it was negatively predicted by neuroticism. Stress management was positively predicted by extraversion and conscientiousness, and it was negatively predicted by neuroticism. Restorative sleep was positively predicted by extraversion and conscientiousness, and it was negatively predicted by neuroticism. Social support for healthy practices was positively predicted by extraversion, agreeableness, and conscientiousness. Environmental exposures were positively predicted by extraversion, and neuroticism was positively and negatively predicted by conscientiousness. Our findings may be useful for further exploration of personality risk factors for healthy practices in challenging life circumstances.

Keywords: healthy lifestyles; personality traits; COVID-19 pandemic

1. Introduction

A healthy lifestyle can be defined as complex behavior patterns and routines that include healthy practices and exclude harmful habits. Healthy practices imply balanced



Citation: Zolotareva, A.; Shchebetenko, S.; Belousova, S.; Danilova, I.; Tseilikman, V.; Lapshin, M.; Sarapultseva, L.; Makhniova, S.; Sarapultseva, M.; Komelkova, M.; et al. Big Five Traits as Predictors of a Healthy Lifestyle during the COVID-19 Pandemic: Results of a Russian Cross-Sectional Study. *Int. J. Environ. Res. Public Health* **2022**, 19, 10716. https://doi.org/10.3390/ ijerph191710716

Academic Editors: Paul B. Tchounwou and Karlijn Massar

Received: 27 July 2022 Accepted: 26 August 2022 Published: 28 August 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). nutrition and high consumption of fruits and vegetables, regular physical exercises and activities [1], rest, and restful sleep [2]. On the contrary, harmful habits involve poor eating [3], physical inactivity and sedentary behavior [4], sleep deficiency and deprivation [5], alcohol consumption [6], excessive smartphone use [7], and active, passive, and electronic smoking [8]. A healthy lifestyle has been associated with increased life expectancy [9], mental health [10], and psychological well-being [11], as well as beneficial effects on the hormonal [12], immune [13], and hematopoietic systems [14].

Many people went through significant transformations in their healthy lifestyles throughout COVID-19. At the beginning of the pandemic, the average number of steps dropped from 10,000 to 4600 steps per day, sleep increased by 25–30 min per night, social time increased by more than half to less than 30 min, and screen time increased by more than double to more than 5 h per day [15]. Previous studies revealed unfavorable immediate and long-term impacts of the pandemic on self-reported physical activities and sedentary behaviors among people [16], including changes in eating habits [17], increased substance use [18], and problematic smartphone use [19]. These led to subsequent weight gain in more than a third of the people surveyed in France [20], Italy [21], Poland [22], UAE [23], and many other countries.

Importantly, individuals with unhealthy lifestyle patterns have been reported to have more severe forms of COVID-19 [24]. The mechanisms that underpin these relations are not fully understood. They may include the activation of the immune and stress systems, as well as the detrimental influence of obesity on the course of the disease [11,25,26]. Unhealthy lifestyles can also lead to reduced immunity and organ injury, predisposing people to diseases, and their severity, commonly defined as 'communicable' such as SARS-CoV-2 [27].

Lifestyle changes during lockdown occurred in both unfavorable and favorable directions. According to the results of the French NutriNet-Santé cohort study, 19.8% of the participants reported positive changes, such as increased consumption of fruits, vegetables, and fish [20], while increased physical activity, especially for bodyweight training, was reported by 38.3% of respondents from Italy [21]. In children and adolescents, the adverse effects of SARS-CoV-2 include an unbalanced diet with an increased risk of overweight or nutritional deficiencies, a sedentary lifestyle, lack of schooling, social isolation, and deteriorating mental health [28].

Although a healthy lifestyle was critical during the COVID-19 pandemic, personality risk factors have not yet been investigated. In pre-pandemic studies, personality traits have been associated with health [29–31]. Physical activity and a vegetable diet were associated with increased extraversion, adherence to the Mediterranean diet, and lower alcohol and cigarette intake, and excessive use of screens and social networking sites was associated with greater conscientiousness. Lower fruit consumption, sleep issues, alcohol and tobacco consumption, smartphone use, and excessive use of the internet were all associated with increased neuroticism. Likewise, according to Zhu et al. (2021), greater perceived vulnerability, feeling stressed, apprehensive, and helpless during the pandemic was associated with more reports of positive lifestyle changes, including increased social/family support, increased awareness of mental health, and a positive lifestyle [32].

Given the above, this study aimed to examine possible associations of the Big Five traits with a healthy lifestyle during the COVID-19 pandemic.

2. Materials and Methods

2.1. Procedure and Participants

The data were collected in March and April 2021. We invited students from four Russian universities in Yekaterinburg and Chelyabinsk to participate in this study. We conducted the study in a psychology class, with 1236 students volunteering. Subsequently, 21 questionnaires were excluded due to incomplete completion, and data from different universities were combined because the procedure and methods of diagnosing COVID-19 disease, the patients' routing protocols in case of positive results, and the algorithms for

3 of 8

inpatient and outpatient treatment were the same throughout the country (in accordance with the orders of the Ministry of Health).

The total sample consisted of 1215 university students (71.9% females) aged 18–26 (M = 19.37, SD = 1.32). All volunteers completed a written informed consent form, disclosing the purposes of this study and stating that they could withdraw from participation at any time. The study was approved by an Institutional Review Board at Chelyabinsk State University (# 6, 17 November 2020) and carried out in accordance with the Declaration of Helsinki [33].

2.2. Measures

2.2.1. Personality Traits

To measure personality traits, we used a Russian version [34] of the Big Five Inventory-10 (BFI-10) [35]. The BFI-10 consists of 10 items evaluating 5 personality traits: extraversion (e.g., "... is outgoing, sociable"), agreeableness (e.g., "... is generally trusting"), conscientiousness (e.g., "... does a thorough job"), neuroticism (e.g., "... gets nervous easily"), and openness to experience (e.g., "... has an active imagination"). Respondents rated their agreement using a five-point Likert scale. The internal consistency was evaluated using Spearman–Brown coefficients. In the current study, the Spearman–Brown coefficients were 0.67, 0.25, 0.47, 0.64, and 0.30 for extraversion, agreeableness, conscientiousness, neuroticism, and openness to experience, respectively.

2.2.2. A Healthy Lifestyle

To measure a healthy lifestyle, we used a translated Russian version of the Short Multidimensional Inventory Lifestyle Evaluation (SMILE-C) [36]. SMILE-C consists of 27 items that examined 7 dimensions of a healthy lifestyle during the COVID-19 pandemic: diet and nutrition (e.g., "Do you eat healthy foods such as fresh fruits, fresh vegetables, whole grains, legumes, or nuts?"), substance use (e.g., "Do you use marijuana or hashish?"), physical activity (e.g., "Do you exercise for at least 30 min daily (or 150 min a week)?"), stress management (e.g., "Do you make time to relax?"), restorative sleep (e.g., "Do you manage to sleep between 7 and 9 h per night?"), social support (e.g., "Do you spend time on a computer/smartphone within one hour of going to sleep?"). Respondents rated their agreement using a four-point Likert scale. Higher scores indicated a healthier lifestyle. In the current study, Cronbach's alpha was 0.47, 0.56, 0.60, 0.53, and 0.80 for diet and nutrition, substance use, stress management, restorative sleep, and social support, respectively. The physical activity and environmental exposure scores were measured using a single question.

2.3. Data Analysis

All analyses were performed using software IBM SPSS version 27 (IBM Corporation, Armonk, NY, USA). In the first stage, descriptive statistics and correlation matrix for all study variables were calculated. In the second stage, multivariate regression analyzes were used to examine personality risk factors for a healthy lifestyle during the COVID-19 pandemic.

3. Results

3.1. Preliminary Analyses

Table 1 summarizes the correlation matrix for all study variables.

Table 1. Correlation matrix for study variables.

Model	Ext	Agr	Con	Neu	Ope
Diet and nutrition	0.18 ^a	0.13 ^a	0.29 ^a	-0.16 ^a	0.04
Substance abuse	-0.08 ^b	0.12 ^a	0.13 ^a	0.05	0.02
Physical activity	0.19 ^a	0.01	0.25 ^a	-0.12 ^a	0.07 ^c

Model	Ext	Agr	Con	Neu	Ope
Stress management	0.34 ^a	0.06	0.32 ^a	-0.19 ^a	0.10 ^a
Restorative sleep	0.14 ^a	0.05	0.20 ^a	-0.17^{a}	0.01
Social support	0.40 ^a	0.15 ^a	0.20 ^a	-0.10^{a}	0.10 ^a
Environmental exposures	0.05	0.01	-0.11^{a}	0.10 ^a	-0.02

Table 1. Cont.

Note: ^a p < 0.05, ^b p < 0.01, ^c p < 0.001. Ext = Extraversion, Agr = Agreebleness, Con = Conscientiousness, Neu = Neuroticism, Ope = Openness to experience.

Personality traits were correlated with many healthy lifestyles. Therefore, extraversion was negatively associated with substance use and positively associated with diet and nutrition, physical activity, stress management, restorative sleep, and social support for healthy practices. Agreeableness was positively related to diet and nutrition, substance use, and social support for healthy practices. Consciousness was negatively correlated with environmental exposures and positively correlated with diet and nutrition, substance use, physical activity, stress management, restorative sleep, and social support for healthy practices. Because higher SMILE-C scores indicated a healthier lifestyle, individuals with increased conscientiousness and agreeableness showed decreased substance use. Neuroticism was positively related to environmental exposures and negatively related to diet and nutrition, physical activity, stress management, restorative sleep, and social support for healthy practices. Openness to experience was positively correlated with physical activity, stress management, restorative sleep, and social support for healthy practices. Openness to experience was positively correlated with physical activity, stress management, restorative sleep, and social support for healthy practices.

3.2. Regression Analyses

We tested linear regression models that predict changes in COVID-19 on seven lifestyle indicators separately (Table 2).

Model	В	SE	β	t	p
Me	odel 1: Risk fa	ctors for diet	and nutrition		
Extraversion	0.11	0.03	0.09	3.27	0.001
Agreeableness	0.13	0.04	0.10	3.48	0.001
Conscientiousness	0.34	0.04	0.25	8.82	< 0.001
Neuroticism	-0.13	0.03	-0.11	-4.09	< 0.001
Openness to experience	0.01	0.04	0.01	0.21	0.838
Μ	lodel 2: Risk fa	actors for sub	stance abuse		
Extraversion	-0.12	0.03	-0.12	-4.05	< 0.001
Agreeableness	0.15	0.03	0.13	4.41	< 0.001
Conscientiousness	0.17	0.03	0.15	5.05	< 0.001
Neuroticism	0.05	0.03	0.05	1.70	0.089
Openness to experience	0.02	0.03	0.01	0.49	0.627
М	lodel 3: Risk fa	actors for phy	vsical activity		
Extraversion	0.07	0.01	0.13	4.67	< 0.001
Agreeableness	-0.02	0.02	-0.03	-1.19	0.234
Conscientiousness	0.12	0.02	0.22	7.59	< 0.001
Neuroticism	-0.04	0.01	-0.08	-2.73	0.007
Openness to experience	0.02	0.02	0.04	1.38	0.168
Мо	del 4: Risk fac	tors for stres	s management	:	
Extraversion	0.43	0.04	0.26	9.79	< 0.001
Agreeableness	-0.01	0.05	-0.01	-0.12	0.907
Conscientiousness	0.46	0.05	0.24	9.02	< 0.001
Neuroticism	-0.20	0.04	-0.13	-4.83	< 0.001
Openness to experience	0.08	0.05	0.05	1.82	0.070

Table 2. Linear regression models that predict a healthy lifestyle during the COVID-19 pandemic.

Model	В	SE	β	t	p
Ν	/lodel 5: Risk fa	actors for res	torative sleep		
Extraversion	0.09	0.03	0.08	2.66	0.008
Agreeableness	0.03	0.04	0.02	0.72	0.474
Conscientiousness	0.23	0.04	0.17	5.86	<0.001
Neuroticism	-0.15	0.03	-0.14	-4.88	<0.001
Openness to experience	-0.02	0.04	-0.02	-0.53	0.597
	Model 6: Risk	factors for so	cial support		
Extraversion	0.62	0.05	0.35	12.94	<0.001
Agreeableness	0.20	0.05	0.10	3.67	<0.001
Conscientiousness	0.22	0.05	0.11	4.10	<0.001
Neuroticism	-0.05	0.05	-0.03	-1.00	0.317
Openness to experience	0.08	0.05	0.04	1.51	0.131
Mode	l 7: Risk factor	s for environ	mental exposu	ires	
Extraversion	0.03	0.01	0.09	3.08	0.002
Agreeableness	0.01	0.01	0.01	0.16	0.870
Conscientiousness	-0.05	0.01	-0.11	-3.83	<0.001
Neuroticism	0.04	0.01	0.11	3.68	<0.001
Openness to experience	-0.01	0.01	-0.02	-0.85	0.397

Table 2. Cont.

Note: Significant predictors are marked in bold.

Extraversion, agreeableness, and conscientiousness were positively predicted while neuroticism negatively predicted diet and nutrition, $R^2 = 0.11$, F (5, 1209) = 32.14, p < 0.001. Substance abuse was positively predicted by agreeability and conscientiousness and was negatively predicted by extraversion, $R^2 = 0.05$, F (5, 1209) = 11.64, p < 0.001. Extraversion and conscientiousness positively predicted while neuroticism negatively predicted physical activity, $R^2 = 0.09$, F (5, 1209) = 24.40, p < 0.001. Stress management was positively predicted by extraversion and conscientiousness, and negatively predicted by neuroticism, $R^2 = 0.20$, F (5, 1209) = 58.61, p < 0.001. Extraversion and conscientiousness positively predicted restorative sleep, $R^2 = 0.07$, F (5, 1209) = 17.77, p < 0.001. Social support for healthy practices was positively predicted by extraversion, agreeableness, and conscientiousness, $R^2 = 0.18$, F (5, 1209) = 54.06, p < 0.001. Extraversion and neuroticism were positively predicted while conscientiousness negatively predicted environmental exposures, $R^2 = 0.03$, F (5, 1209) = 6.82, p < 0.001. Therefore, openness to experience was the only variable that did not incrementally predict a healthy lifestyle during the COVID-19 pandemic.

4. Discussion

This study reveals that personality traits predicted various indicators of a healthy lifestyle during the COVID-19 pandemic. Extraversion, agreeableness, and conscientiousness were found to be favorable predictors of numerous aspects of a healthy lifestyle, while neuroticism was found to be a negative predictor. There was one exception to these findings. Extraversion was found to be a negative predictor of substance use. In other words, the higher the extraversion score, the less likely people were to use alcohol, tobacco, or psychoactive substances. Similar findings have previously been obtained in the areas of stimulant use, smoking, alcohol consumption, and opioid dependency [37–39]. Higher extroversion was also associated with a perceived decrease in alcohol consumption and a shift to a healthier diet among Finnish women during the COVID-19 pandemic [40]. Furthermore, during the COVID-19 pandemic, receptivity to new experiences did not predict a healthy lifestyle. This finding supports previous research that found no significant links between openness to experience and hedonistic lifestyles among university students [41] or health behaviors among young people [42].

Personality traits largely determine the degree to which people adapt to the pandemic, although it appears that personality characteristics can be the basis for healthy lifestyles

regardless of COVID-19. Thus, lower neuroticism and higher extraversion, agreeableness, and consciousness were associated with improved mood, decreased perceived stress, and increased participation in health promotion among college students both before and after the onset of the pandemic [43]. A recent large-scale panel study of the British population (UK Household Longitudinal Study, UKHLS) found that during the pandemic, people with more extroverted and open personality traits reported greater deterioration in mental health, while those with higher agreeableness scores suffered less [44]. Neuroticism significantly predicted cross-sectional mental health deterioration but did not significantly worsen during the pandemic. Similar to previous findings, our study demonstrated a sustained role of personality traits in healthy lifestyles and mental health, which is intensified in times of crisis such as COVID-19.

This study has several strengths and limitations. In contrast to previous studies, our findings have expanded ideas about the relationship between personality traits and healthy lifestyles during the COVID-19 pandemic. In addition to alcohol consumption [40], our study covered many other aspects of a healthy lifestyle. As for limitations, the nature of the study's self-report could distort the objective picture of healthy practices and harmful habits. Previous studies showed that self-reporting underestimated sedentary behavior compared to device measurements [45], underreported energy intake, especially when the body mass index of the respondents' increases [46], and underestimated smoking behavior compared to biomarkers [47]. Next, this study is cross-sectional, which limits the possibility of causal inferences from the findings obtained, although the regression analysis is more suitable for longitudinal data [48]. Moreover, we employed a convenience sample of Russian undergraduates, which could distort revealed associations to a more global population. Some concerns are also raised by the fact that the study sample was only students from four Russian universities in two nearby cities. In this regard, we understand that our findings may be explained to some extent by geographic characteristics. Finally, we did not control for the experience of COVID-19 disease, but this factor could have influenced our findings.

5. Conclusions

This study highlighted the relationship between the five main characteristics and a healthy lifestyle during the COVID-19 pandemic. Our findings extend previous research on the relationship between personality traits and healthy practices by describing this relationship in challenging life circumstances. Although this relationship needs to be tested in longitudinal studies, we anticipate that our findings will help scientists and practitioners better understand and adjust personality risk variables for healthy behaviors. Specifically, we suggest that preventive measures and psychoeducational activities may be useful for young people with personality traits responsible for unhealthy lifestyles.

Author Contributions: Conceptualization, A.Z. and A.S.; methodology, A.Z., S.S., S.B., D.H. and S.L.; formal analysis, A.Z. and S.S.; investigation, S.B., I.D., V.T., M.L., L.S., S.M., M.S., M.K. and E.L.; resources, S.B. and I.D.; data curation, A.Z. and D.H.; writing—original draft preparation, A.Z.; writing—review and editing, S.S. and D.H. All authors have read and agreed to the published version of the manuscript.

Funding: This study was funded by RFBR (project No. 20-515-55003) and partly by the Government contract of the Institute of Immunology and Physiology (AAAA-A21-121012090090-9).

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki and approved by the Institutional Review of Chelyabinsk State University (# 6, 17 November 2020).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: The datasets analyzed during the current study are available from the corresponding author upon reasonable request.

Conflicts of Interest: The authors declare no conflict of interest.

References

- 1. Daskalopoulou, C.; Stubbs, B.; Kralj, C.; Koukounari, A.; Prince, M.; Prina, A.M. Physical Activity and Healthy Ageing: A Systematic Review and Meta-Analysis of Longitudinal Cohort Studies. *Ageing Res. Rev.* **2017**, *38*, 6–17. [CrossRef] [PubMed]
- Lamp, A.; Cook, M.; Soriano Smith, R.N.; Belenky, G. Exercise, Nutrition, Sleep, and Waking Rest? Sleep 2019, 42, zsz138. [CrossRef]
- Afshin, A.; Sur, P.J.; Fay, K.A.; Cornaby, L.; Ferrara, G.; Salama, J.S.; Mullany, E.C.; Abate, K.H.; Abbafati, C.; Abebe, Z.; et al. Health Effects of Dietary Risks in 195 Countries, 1990–2017: A Systematic Analysis for the Global Burden of Disease Study 2017. *Lancet* 2019, 393, 1958–1972. [CrossRef]
- 4. González, K.; Fuentes, J.; Márquez, J.L. Physical Inactivity, Sedentary Behavior and Chronic Diseases. *Korean J. Fam. Med.* 2017, 38, 111–115. [CrossRef] [PubMed]
- 5. Kohansieh, M.; Makaryus, A.N. Sleep Deficiency and Deprivation Leading to Cardiovascular Disease. *Int. J. Hypertens.* 2015, 2015, 615681. [CrossRef] [PubMed]
- Griswold, M.G.; Fullman, N.; Hawley, C.; Arian, N.; Zimsen, S.R.M.; Tymeson, H.D.; Venkateswaran, V.; Tapp, A.D.; Forouzanfar, M.H.; Salama, J.S.; et al. Alcohol Use and Burden for 195 Countries and Territories, 1990–2016: A Systematic Analysis for the Global Burden of Disease Study 2016. *Lancet* 2018, 392, 1015–1035. [CrossRef]
- Sohn, S.Y.; Rees, P.; Wildridge, B.; Kalk, N.J.; Carter, B. Prevalence of Problematic Smartphone Usage and Associated Mental Health Outcomes amongst Children and Young People: A Systematic Review, Meta-Analysis and GRADE of the Evidence. *BMC Psychiatry* 2019, 19, 356. [CrossRef]
- 8. Kim, S.Y.; Sim, S.; Choi, H.G. Active, Passive, and Electronic Cigarette Smoking Is Associated with Asthma in Adolescents. *Sci. Rep.* 2017, 7, 17789. [CrossRef]
- Chudasama, Y.V.; Khunti, K.; Gillies, C.L.; Dhalwani, N.N.; Davies, M.J.; Yates, T.; Zaccardi, F. Healthy Lifestyle and Life Expectancy in People with Multimorbidity in the UK Biobank: A Longitudinal Cohort Study. *PLoS Med.* 2020, 17, e1003332. [CrossRef]
- 10. Velten, J.; Bieda, A.; Scholten, S.; Wannemüller, A.; Margraf, J. Lifestyle Choices and Mental Health: A Longitudinal Survey with German and Chinese Students. *BMC Public Health* **2018**, *18*, 632. [CrossRef]
- Sapranaviciute-Zabazlajeva, L.; Luksiene, D.; Virviciute, D.; Bobak, M.; Tamosiunas, A. Link between Healthy Lifestyle and Psychological Well-Being in Lithuanian Adults Aged 45–72: A Cross-Sectional Study. *BMJ Open* 2017, 7, e014240. [CrossRef] [PubMed]
- 12. Armamento-Villareal, R.; Aguirre, L.E.; Qualls, C.; Villareal, D.T. Effect of Lifestyle Intervention on the Hormonal Profile of Frail, Obese Older Men. J. Nutr. Health Aging 2016, 20, 334–340. [CrossRef]
- Chastin, S.F.M.; Abaraogu, U.; Bourgois, J.G.; Dall, P.M.; Darnborough, J.; Duncan, E.; Dumortier, J.; Pavón, D.J.; McParland, J.; Roberts, N.J.; et al. Effects of Regular Physical Activity on the Immune System, Vaccination and Risk of Community-Acquired Infectious Disease in the General Population: Systematic Review and Meta-Analysis. *Sports Med.* 2021, *51*, 1673–1686. [CrossRef] [PubMed]
- Nahrendorf, M.; Swirski, F.K. Lifestyle Effects on Hematopoiesis and Atherosclerosis. Circ. Res. 2015, 116, 884–894. [CrossRef] [PubMed]
- 15. Giuntella, O.; Hyde, K.; Saccardo, S.; Sadoff, S. Lifestyle and Mental Health Disruptions During COVID-19. *Proc. Natl. Acad. Sci.* USA 2021, 118, e2016632118. [CrossRef] [PubMed]
- 16. Zhou, J.; Xie, X.; Guo, B.; Pei, R.; Pei, X.; Yang, S.; Jia, P. Impact of COVID-19 Lockdown on Physical Activity Among the Chinese Youths: The COVID-19 Impact on Lifestyle Change Survey (COINLICS). *Front. Public Health* **2021**, *9*, 592795. [CrossRef] [PubMed]
- 17. Prete, M.; Luzzetti, A.; Augustin, L.S.A.; Porciello, G.; Montagnese, C.; Calabrese, I.; Ballarin, G.; Coluccia, S.; Patel, L.; Vitale, S.; et al. Changes in Lifestyle and Dietary Habits during COVID-19 Lockdown in Italy: Results of an Online Survey. *Nutrients* **2021**, *13*, 1923. [CrossRef]
- 18. Horigian, V.E.; Schmidt, R.D.; Feaster, D.J. Loneliness, Mental Health, and Substance Use among US Young Adults during COVID-19. J. Psychoact. Drugs 2021, 53, 1–9. [CrossRef]
- 19. Jin, L.; Hao, Z.; Huang, J.; Akram, H.R.; Saeed, M.F.; Ma, H. Depression and Anxiety Symptoms Are Associated with Problematic Smartphone Use under the COVID-19 Epidemic: The Mediation Models. *Child. Youth Serv. Rev.* **2021**, *121*, 105875. [CrossRef]
- Deschasaux-Tanguy, M.; Druesne-Pecollo, N.; Esseddik, Y.; de Edelenyi, F.S.; Allès, B.; Andreeva, V.A.; Baudry, J.; Charreire, H.; Deschamps, V.; Egnell, M.; et al. Diet and Physical Activity during the Coronavirus Disease 2019 (COVID-19) Lockdown (March–May 2020): Results from the French NutriNet-Santé Cohort Study. Am. J. Clin. Nutr. 2021, 113, 924–938. [CrossRef]
- 21. Di Renzo, L.; Gualtieri, P.; Pivari, F.; Soldati, L.; Attinà, A.; Cinelli, G.; Leggeri, C.; Caparello, G.; Barrea, L.; Scerbo, F.; et al. Eating Habits and Lifestyle Changes during COVID-19 Lockdown: An Italian Survey. J. Transl. Med. 2020, 18, 229. [CrossRef] [PubMed]
- Górnicka, M.; Drywień, M.E.; Zielinska, M.A.; Hamułka, J. Dietary and Lifestyle Changes During COVID-19 and the Subsequent Lockdowns among Polish Adults: A Cross-Sectional Online Survey PLifeCOVID-19 Study. *Nutrients* 2020, 12, 2324. [CrossRef] [PubMed]
- Radwan, H.; Al Kitbi, M.; Hasan, H.; Al Hilali, M.; Abbas, N.; Hamadeh, R.; Saif, E.R.; Naja, F. Indirect Health Effects of COVID-19: Unhealthy Lifestyle Behaviors during the Lockdown in the United Arab Emirates. *Int. J. Environ. Res. Public Health* 2021, 18, 1964. [CrossRef] [PubMed]

- Tavakol, Z.; Ghannadi, S.; Tabesh, M.R.; Halabchi, F.; Noormohammadpour, P.; Akbarpour, S.; Alizadeh, Z.; Nezhad, M.H.; Reyhan, S.K. Relationship between Physical Activity, Healthy Lifestyle and COVID-19 Disease Severity; a Cross-Sectional Study. Z. Gesundh. Wiss. 2021, 1–9. [CrossRef]
- 25. Lange, K.W.; Nakamura, Y. Lifestyle Factors in the Prevention of COVID-19. Glob. Health J. 2020, 4, 146–152. [CrossRef]
- 26. Zhao, X.; Gang, X.; He, G.; Li, Z.; Lv, Y.; Han, Q.; Wang, G. Obesity Increases the Severity and Mortality of Influenza and COVID-19: A Systematic Review and Meta-Analysis. *Front. Endocrinol.* **2020**, *11*, 595109. [CrossRef]
- 27. Doraiswamy, S.; Cheema, S.; Mamtani, R. The COVID-19 and Lifestyle Nexus: Settling the Debate. *Glob. Health Promot.* 2022, 29, 96–100. [CrossRef]
- 28. Scapaticci, S.; Neri, C.R.; Marseglia, G.L.; Staiano, A.; Chiarelli, F.; Verduci, E. The Impact of the COVID-19 Pandemic on Lifestyle Behaviors in Children and Adolescents: An International Overview. *Ital. J. Pediatr.* **2022**, *48*, 22. [CrossRef]
- Cheng, C.-H.E.; Weiss, J.W.; Siegel, J.M. Personality Traits and Health Behaviors as Predictors of Subjective Wellbeing among a Multiethnic Sample of University-Attending Emerging Young Adults. *Int. J. Wellbeing* 2015, *5*, 21–43. [CrossRef]
- 30. Peterka-Bonetta, J.; Sindermann, C.; Elhai, J.D.; Montag, C. Personality Associations with Smartphone and Internet Use Disorder: A Comparison Study Including Links to Impulsivity and Social Anxiety. *Front. Public Health* **2019**, *7*, 127. [CrossRef]
- Yañez, A.M.; Bennasar-Veny, M.; Leiva, A.; García-Toro, M. Implications of Personality and Parental Education on Healthy Lifestyles among Adolescents. *Sci. Rep.* 2020, *10*, 7911. [CrossRef] [PubMed]
- 32. Zhu, S.; Zhuang, Y.; Ip, P. Impacts on Children and Adolescents' Lifestyle, Social Support and Their Association with Negative Impacts of the COVID-19 Pandemic. *Int. J. Environ. Res. Public Health* **2021**, *18*, 4780. [CrossRef] [PubMed]
- 33. World Medical Association. World Medical Association Declaration of Helsinki: Ethical Principles for Medical Research Involving Human Subjects. *JAMA* 2013, *310*, 2191–2194. [CrossRef]
- 34. Shchebetenko, S. "The best man in the world": Attitudes toward personality traits. Psychology. J. High. Sch. Econ. 2014, 11, 129–148. (In Russian)
- 35. Rammstedt, B.; John, O.P. Measuring Personality in One Minute or Less: A 10-Item Short Version of the Big Five Inventory in English and German. *J. Res. Personal.* 2007, *41*, 203–212. [CrossRef]
- Balanzá-Martínez, V.; Kapczinski, F.; de Azevedo Cardoso, T.; Atienza-Carbonell, B.; Rosa, A.R.; Mota, J.C.; De Boni, R.B. The Assessment of Lifestyle Changes during the COVID-19 Pandemic Using a Multidimensional Scale. *Rev. Psiquiatr. Salud Ment. Engl. Ed.* 2021, 14, 16–26. [CrossRef] [PubMed]
- 37. Hokm Abadi, M.E.; Bakhti, M.; Nazemi, M.; Sedighi, S.; Mirzadeh Toroghi, E. The Relationship between Personality Traits and Drug Type among Substance Abuse. *J. Res. Health* **2018**, *8*, 531–540. [CrossRef]
- 38. Kornør, H.; Nordvik, H. Five-Factor Model Personality Traits in Opioid Dependence. BMC Psychiatry 2007, 7, 37. [CrossRef]
- 39. Turiano, N.A.; Whiteman, S.D.; Hampson, S.E.; Roberts, B.W.; Mroczek, D.K. Personality and Substance Use in Midlife: Conscientiousness as a Moderator and the Effects of Trait Change. *J. Res. Personal.* **2012**, *46*, 295–305. [CrossRef]
- Kekäläinen, T.; Hietavala, E.-M.; Hakamäki, M.; Sipilä, S.; Laakkonen, E.K.; Kokko, K. Personality Traits and Changes in Health Behaviors and Depressive Symptoms during the COVID-19 Pandemic: A Longitudinal Analysis from Pre-pandemic to Onset and End of the Initial Emergency Conditions in Finland. *Int. J. Environ. Res. Public Health* 2021, 18, 7732. [CrossRef]
- 41. Taufik, T.; Prihartanti, N.; Hamid, H.S.A. Neuroticism, Extraversion and Conscientiousness as Predictors of the Hedonistic Lifestyle. *N. Am. J. Psychol.* **2019**, *21*, 645–660.
- 42. Obara-Gołębiowska, M.; Michałek-Kwiecień, J. Personality Traits, Dieting Self-Efficacy and Health Behaviors in Emerging Adult Women: Implications for Health Promotion and Education. *Health Promot. Perspect.* **2020**, *10*, 230–237. [CrossRef]
- Rettew, D.C.; McGinnis, E.W.; Copeland, W.; Nardone, H.Y.; Bai, Y.; Rettew, J.; Devadenam, V.; Hudziak, J.J. Personality Trait Predictors of Adjustment During the COVID Pandemic Among College Students. *PLoS ONE* 2021, 16, e0259431. [CrossRef] [PubMed]
- Proto, E.; Zhang, A. COVID-19 and Mental Health of Individuals with Different Personalities. *Proc. Natl. Acad. Sci. USA* 2021, 118, e2109282118. [CrossRef] [PubMed]
- Prince, S.A.; Cardilli, L.; Reed, J.L.; Saunders, T.J.; Kite, C.; Douillette, K.; Fournier, K.; Buckley, J.P. A Comparison of Self-Reported and Device Measured Sedentary Behaviour in Adults: A Systematic Review and Meta-Analysis. *Int. J. Behav. Nutr. Phys. Act.* 2020, 17, 31. [CrossRef] [PubMed]
- Ravelli, M.N.; Schoeller, D.A. Traditional Self-Reported Dietary Instruments Are Prone to Inaccuracies and New Approaches Are Needed. *Front. Nutr.* 2020, 7, 90. [CrossRef] [PubMed]
- Khouja, J.N.; Munafò, M.R.; Relton, C.L.; Taylor, A.E.; Gage, S.H.; Richmond, R.C. Investigating the Added Value of Biomarkers Compared with Self-Reported Smoking in Predicting Future e-Cigarette Use: Evidence from a Longitudinal UK Cohort Study. *PLoS ONE* 2020, 15, e0235629. [CrossRef]
- Yao, F.; Müller, H.-G.; Wang, J.-L. Functional Linear Regression Analysis for Longitudinal Data. Ann. Stat. 2005, 33, 2873–2903. [CrossRef]