



Article

Cross-Sectional Associations of Self-Reported Social/Emotional Support and Life Satisfaction with Smoking and Vaping Status in Adults

Zidian Xie ^{1,*}, Francisco Cartujano-Barrera ², Paula Cupertino ² and Dongmei Li ^{1,*}

¹ Department of Clinical and Translational Research, University of Rochester Medical Center, Rochester, NY 14642, USA

² Department of Public Health Sciences and Cancer Center, University of Rochester Medical Center, Rochester, NY 14642, USA

* Correspondence: zidian_xie@urmc.rochester.edu (Z.X.); dongmei_li@urmc.rochester.edu (D.L.)

Abstract: This study aimed to examine the cross-sectional association of self-reported social/emotional support and life satisfaction with smoking/vaping status in US adults. The study included 47,163 adult participants who self-reported social/emotional support, life satisfaction, and smoking/vaping status in the 2016 and 2017 BRFSS national survey data. We used multivariable weighted logistic regression models to measure the cross-sectional association of self-reported social/emotional support and life satisfaction with smoking/vaping status. Compared to never users, dual users and exclusive smokers were more likely to have low life satisfaction, with an adjusted odds ratio (aOR) = 1.770 (95% confidence interval [CI]: 1.135, 2.760) and an aOR = 1.452 (95% CI: 1.121, 1.880) respectively, especially for the age group 18–34. Exclusive cigarette smokers were more likely to have low life satisfaction compared to ex-smokers (aOR = 1.416, 95% CI: 1.095, 1.831). Exclusive cigarette smokers were more likely to have low social/emotional support (aOR = 1.193, 95% CI: 1.030, 1.381) than never users, especially those aged 65 and above. In addition, exclusive cigarette smokers were more likely to have low social/emotional support than ex-smokers, with an aOR = 1.279 (95% CI: 1.097, 1.492), which is more pronounced among the age group 18–34, as well as 65 and above. Our results suggest that life satisfaction and social/emotional support may play important roles in smoking and vaping, which should be incorporated into behavioral interventions to reduce tobacco use.

Keywords: smoking; vaping; social/emotional support; life satisfaction; smoking cessation



Citation: Xie, Z.; Cartujano-Barrera, F.; Cupertino, P.; Li, D.

Cross-Sectional Associations of Self-Reported Social/Emotional Support and Life Satisfaction with Smoking and Vaping Status in Adults. *Int. J. Environ. Res. Public Health* **2022**, *19*, 10722. <https://doi.org/10.3390/ijerph191710722>

Academic Editor: Paul B. Tchounwou

Received: 13 July 2022

Accepted: 26 August 2022

Published: 28 August 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Tobacco smoking kills more than 8 million people each year worldwide [1]. In 2019, approximately 34.1 million adults were cigarette smokers in the United States (US) [2]. While there has been a decline in smoking prevalence in recent years, e-cigarette use (vaping) has become increasingly popular, especially among youths (ages 12 to 18) and young adults [3,4]. In 2019, 4.5% of adults in the US were current e-cigarette users [2]. While the long-term health effects of vaping remain to be determined, vaping is associated with many health effects, including respiratory and cardiovascular diseases [5–10]. Therefore, it is of utmost importance to understand psychosocial constructs related to smoking and vaping to inform future smoking and vaping cessation interventions.

Smoking and vaping are related to poor mental health, including quality of life, depression, and anxiety [11–14]. Nicotine triggers the release of dopamine, which plays a significant role in emotional responses and reward-motivated behavior [15]. Therefore, tobacco use by smoking or vaping might be related to the relaxation reduction of stress and anxiety. On the other hand, smoking and vaping have both been associated with psychiatric symptoms (such as depression) [16–19]. Defined as an individual's cognitive evaluation of their life, life satisfaction has strong interactive effects on mental health, as one of the main

dimensions [20,21]. Several previous studies have shown that low life satisfaction is highly associated with substance use, including marijuana, alcohol, cannabis, and smoking [22,23], as well as other addictive behaviors [24].

While most smokers are willing to quit (55.1% of adult smokers in 2018), the success rates are very low (7.5% in 2018), even with different treatments, including nicotine replacement therapy [3]. Once addicted to nicotine, it is hard to stop, due to the negative emotional responses involved in the withdrawal effect of nicotine [25]. Social and emotional support is defined as the social resources that one receives as a result of empathetic, caring, and reassuring communication from non-professionals [26]. As moderators of stress, social and emotional support are usually regarded as an essential factor for smoking cessation, although their exact role remains controversial [27,28]. Several studies have shown that positive social and emotional support are associated with successful smoking cessation and prevention of relapse [29–31]. There is one study that showed that adult smokers in South Korean rural areas had high levels of social support [32]. However, whether life satisfaction and social/emotional support might be associated with vaping remains undetermined.

This study aimed to examine the cross-sectional association of self-reported life satisfaction and social/emotional support with smoking and vaping status, using the US nationally represented survey data. Because vaping is more prevalent among young adults [33], the present study will also expand the understanding of the association of smoking and vaping with social/emotional support and life satisfaction across age groups. Results from the present study are intended to inform the development of future interventions designed to promote both smoking and vaping prevention and/or cessation among US adults.

2. Materials and Methods

2.1. Study Participants

The Behavioral Risk Factor Surveillance System (BRFSS) is an annual interview survey, conducted through landlines and cell phones by the Centers for Disease Control and Prevention (CDC) in the US, which collects the information about basic demographics (such as age, gender, education, and income), health-related risk behaviors, and chronic health conditions among adult participants in all 50 US states, as well as the District of Columbia and the three US territories (including Puerto Rico, Guam, and U.S. Virgin Islands) [34]. The BRFSS survey was designed in a way such that data from different years could be easily combined to study the association of e-cigarette use with respiratory diseases (such as COPD and asthma) and other health conditions [10,35–39]. Since the survey data in 2016 and 2017 contain the most recent and similar interview questions related to smoking and vaping, especially life satisfaction and emotional support, we decided to combine the 2016 (486,303 participants) and 2017 (450,016 participants) BRFSS data in our analysis. The 2016 and 2017 BRFSS data are publicly available from the CDC website: https://www.cdc.gov/brfss/annual_data/annual_data.htm (accessed on 16 February 2021).

2.2. Vaping and Smoking Categories

Based on their current smoking and vaping status, as well as past smoking experience, we grouped the adult participants into six smoking and vaping categories: (1) exclusive smokers: currently established smokers (have smoked at least 100 cigarettes in their entire life and now smoke every day or some days) who were not currently established vapers; (2) ex-smokers: previous smokers (have smoked at least 100 cigarettes in their entire life and now do not smoke cigarettes at all) who were not currently established vapers; (3) current vapers who were ex-smokers: currently established vapers (using e-cigarettes every day or some days) who were ex-smokers; (4) exclusive vapers: currently established vapers who were never smokers; (5) dual users: currently established smokers and currently established vapers; (6) never users: never smokers (have smoked less than 100 cigarettes in their entire life, and now do not smoke cigarettes at all) who also were not currently established vapers.

2.3. Main Measures

The primary outcome variables of interest in this study included self-reported life satisfaction and social/emotional support. The variable for life satisfaction came from the survey question, “In general, how satisfied are you with your life?”. We classified it into two levels, one as “low life satisfaction”, with the response of “dissatisfied” or “very dissatisfied”, and the other as “high life satisfaction”, with the response of “satisfied” or “very satisfied.” The variable for social/emotional support came from the survey question, “How often do you get the social and emotional support you need?”. We classified it into two levels: one as “high emotional support”, with the answer of “always” or “usually,” and the other as “low emotional support”, with the response of “sometimes” and “rarely”.

2.4. Covariates

Using the purposeful variable selection method [10,40], covariates that significantly contributed to the models were selected and controlled for our statistical models, including age, gender, employment status, self-reported general health categories, and self-reported mental health (including stress, depression, and problems with emotions). To examine if there is an age effect on the association of smoking and vaping with life satisfaction and social/emotional support, with the consideration of the sample size, we divided adult participants into three age groups, including “18–34,” “35–64,” and “65+.” It has been shown that compared to adults within other age categories, adults in the 18–34 category have higher odds of e-cigarette use, as well as different socioeconomic statuses, for example, poverty and marital status, which might affect their life satisfaction [33,41,42]. Except for mental health, all other covariates are categorical variables.

2.5. Statistical Analysis

Weighted frequency distributions were calculated to determine the association of smoking and vaping status with each covariate. After adjusting for those selected covariates, multivariate weighted logistic regression models were used to assess the association of tobacco and vaping status with the outcome variables (life satisfaction and social/emotional support). To account for the complex sampling design in the BRFSS survey study, the variable _LLCPWT as the final weight, the stratification variable _STSTR, and the clustering variable _PSU for each adult participant were included in the statistical models. The final weight was half of the weight from each year (2016 and 2017) to ensure the weighted sample size was equivalent to the population size [10]. To measure the association of smoking and vaping status with life satisfaction and social/emotional support, multivariable weighted logistic regression models with PROC SURVEY procedures in SAS V9.4 (SAS Institute Inc., Cary, NC, USA) were used to calculate the adjusted odds ratios (aORs) and 95% confidence intervals (CIs). The Taylor series linearization method was used to calculate the standard deviations. All significance tests were two-sided, with a significance level of 5%.

3. Results

3.1. Demographic Characteristics of Survey Participants

From the combined 2016 and 2017 BRFSS survey data, there were 47,163 out of 936,319 adult participants included in our study who indicated their smoking and vaping status, as well as provided valid responses to our main measures about life satisfaction and social/emotional support. Among them, 6121 (12.98%) were exclusive smokers, 806 (1.71%) were dual users, 13,376 (28.36%) were ex-smokers, 26,198 (55.55%) were never users, 454 (0.96%) were current vapers who were ex-smokers, and 208 (0.44%) were exclusive vapers.

As shown in Table 1, the majority of participants aged between 35 and 64 years in all smoking/vaping categories except exclusive vapers. The majority of exclusive vapers were young, aged 18–34 (90.38%) and never married (76.75%). While there was a similar proportion of males and females in other categories (such as dual users, exclusive smokers, ex-smokers, and never users), there were more males than females in current vapers who

were ex-smokers (60.87% vs. 39.13%) or exclusive vapers (67.87% vs. 32.14%). Among exclusive vapers, 28.80% were students, which was higher than in other smoking/vaping categories. Of ex-smokers, 30.32% were retired, higher than in other smoking/vaping categories.

Table 1. The distribution of co-variables within smoking/vaping status.

		Current Vaping and Smoking Status (% with 95% CI)						p
Variables	Levels	Exclusive Smokers (n = 6121)	Dual Users (n = 806)	Current Vapers Who Were Ex-Smokers (n = 454)	Exclusive Vapers (n = 208)	Ex-Smokers (n = 13,376)	Never Users (n = 26,198)	
Age								<0.0001
	18–34	28.56 (27.12, 30.07)	44.52 (40.91, 48.45)	35.44 (31.64, 39.70)	90.38 (89.26, 91.52)	13.05 (12.08, 14.11)	31.63 (30.86, 32.42)	
	35–64	58.22 (57.53, 58.93)	50.46 (49.00, 51.97)	57.85 (54.90, 60.96)	9.17 (5.98, 14.03)	51.73 (51.08, 52.39)	49.60 (49.09, 50.11)	
Gender	65+	13.22 (12.59, 13.88)	5.01 (4.31, 5.83)	6.71 (5.59, 8.05)	0.45 (0.13, 1.66)	35.22 (34.80, 35.65)	18.77 (18.42, 19.13)	<0.0001
	Male	51.36 (50.29, 52.44)	54.60 (52.21, 57.10)	60.87 (57.22, 64.76)	67.87 (65.04, 70.82)	54.94 (54.30, 55.60)	43.47 (42.87, 44.07)	
	Female	48.64 (47.87, 49.42)	45.40 (43.11, 47.82)	39.13 (37.15, 41.21)	32.14 (27.31, 37.81)	45.06 (44.47, 45.65)	56.53 (56.04, 57.02)	
Marital Status								<0.0001
	Married	39.09 (38.07, 40.14)	35.36 (33.15, 37.72)	54.45 (50.96, 58.20)	8.02 (4.86, 13.27)	60.85 (60.40, 61.30)	55.59 (55.16, 56.02)	
	Divorced	19.36 (18.41, 20.37)	13.60 (11.05, 16.75)	10.58 (8.70, 12.86)	5.50 (3.03, 10.00)	12.82 (12.16, 13.52)	8.39 (7.93, 8.86)	
	Widowed	6.05 (5.47, 6.69)	3.80 (2.71, 5.32)	1.62 (1.02, 2.56)	1.59 (0.67, 3.79)	10.22 (9.77, 10.69)	6.30 (6.00, 6.62)	<0.0001
	Separated	3.68 (2.95, 4.60)	4.31 (2.93, 6.34)	1.06 (0.37, 3.04)	0.00 (1.16, 1.86)	1.47 (1.16, 1.86)	1.19 (0.98, 1.45)	
	Never Married	26.09 (24.90, 27.34)	35.74 (32.40, 39.41)	23.26 (20.17, 26.84)	76.75 (74.05, 79.53)	10.80 (9.97, 11.70)	25.55 (24.76, 26.35)	
Employment	A member of an unmarried couple	5.72 (4.97, 6.59)	7.19 (5.37, 9.63)	9.03 (5.82, 13.99)	8.13 (5.41, 12.18)	3.84 (3.28, 4.50)	2.98 (2.67, 3.34)	<0.0001
	Employed for wages	50.32 (49.30, 51.36)	51.66 (48.93, 54.55)	62.79 (59.98, 65.73)	59.15 (55.10, 63.50)	44.19 (43.42, 44.98)	53.44 (52.91, 53.96)	
	Self-employed	7.99 (7.14, 8.94)	7.40 (5.65, 9.69)	9.16 (5.60, 14.96)	3.82 (1.71, 8.51)	9.28 (8.68, 9.91)	8.63 (8.14, 9.16)	
	Out of work for 1 year or more	4.09 (3.43, 4.88)	5.63 (3.91, 8.12)	1.99 (1.20, 3.31)	1.28 (0.43, 3.81)	1.90 (1.63, 2.21)	1.77 (1.49, 2.09)	<0.0001
	Out of work for less than 1 year	4.13 (3.50, 4.89)	3.71 (2.45, 5.62)	5.04 (2.63, 9.64)	2.57 (1.52, 4.34)	2.01 (1.49, 2.70)	2.28 (1.99, 2.61)	
	A homemaker	5.18 (4.43, 6.05)	4.88 (3.14, 7.58)	3.77 (2.26, 6.30)	1.56 (0.61, 4.03)	4.06 (3.63, 4.53)	5.63 (5.25, 6.03)	
	A student	1.56 (1.14, 2.12)	3.97 (2.39, 6.60)	3.08 (1.77, 5.37)	28.80 (23.90, 34.71)	1.17 (0.83, 1.64)	7.09 (6.52, 7.72)	<0.0001
	Retired	12.23 (11.62, 12.87)	5.54 (4.62, 6.65)	6.78 (5.80, 7.93)	0.95 (0.34, 2.67)	30.32 (29.86, 30.79)	17.02 (16.66, 17.39)	
	Unable to work	14.51 (13.48, 15.61)	17.20 (14.78, 20.00)	7.40 (5.73, 9.57)	1.86 (0.63, 5.42)	7.08 (6.50, 7.70)	4.14 (3.78, 4.54)	
General Health								<0.0001
	Excellent	10.60 (9.66, 11.63)	10.79 (7.75, 15.02)	8.62 (7.22, 10.28)	29.92 (24.42, 36.67)	14.86 (14.17, 15.59)	21.45 (20.83, 22.10)	
	Very good	26.86 (25.76, 28.01)	22.72 (20.18, 25.58)	35.52 (31.39, 40.19)	35.70 (30.87, 41.28)	32.93 (32.20, 33.68)	37.30 (36.74, 37.88)	
	Good	34.55 (33.59, 35.53)	38.32 (35.64, 41.20)	32.81 (29.28, 36.77)	25.47 (21.14, 30.67)	32.44 (31.71, 33.18)	29.25 (28.64, 29.87)	<0.0001
	Fair	19.30 (18.13, 20.53)	16.28 (14.19, 18.67)	16.25 (12.72, 20.75)	7.96 (5.93, 10.68)	13.93 (13.29, 14.60)	9.18 (8.72, 9.66)	
	Poor	8.70 (7.85, 9.64)	11.90 (9.77, 14.48)	6.81 (4.76, 9.74)	0.95 (0.40, 2.25)	5.84 (5.32, 6.41)	2.82 (2.48, 3.19)	
Mental Health	Average days mental health not good	6.09	8.06	5.16	5.45	2.76	2.49	<0.0001
Emotional Support								<0.0001
	Low	30.45 (29.18, 31.77)	32.91 (29.92, 36.21)	22.15 (18.40, 26.66)	19.80 (15.88, 24.72)	19.17 (18.36, 20.01)	17.40 (16.75, 18.06)	
	High	69.55 (69.04, 70.06)	67.09 (65.58, 68.63)	77.85 (76.42, 79.33)	80.20 (77.96, 82.50)	80.83 (80.59, 81.08)	82.60 (82.37, 82.84)	
Life Satisfaction								<0.0001
	Low	12.18 (11.04, 13.44)	15.88 (13.37, 18.86)	7.59 (5.34, 10.79)	8.05 (5.13, 12.63)	4.43 (3.99, 4.93)	3.40 (3.08, 3.75)	
	High	87.82 (87.62, 88.03)	84.12 (83.22, 85.03)	92.41 (91.74, 93.09)	91.95 (90.99, 92.92)	95.57 (95.49, 95.64)	96.60 (96.55, 96.65)	

Note: CI denotes confidence interval.

By comparison, never users and ex-smokers had the best overall self-reported mental health, with 2.49 days and 2.76 days (out of 30 days) of having bad mental health. Relatively, exclusive vapers and current vapers who were ex-smokers had worse self-reported mental health, with 5.45 and 5.16 days of having bad mental health. Exclusive smokers and dual users had the worst self-reported mental health, with 6.09 and 8.06 days of having bad mental health.

3.2. Association of Life Satisfaction with Smoking/Vaping Status

Exclusive smokers and dual users were more likely to receive lower social/emotional support and life satisfaction than other groups (Table 1). For example, 15.88% of dual users and 12.18% of exclusive smokers had low life satisfaction, while it was only 4.43% for ex-smokers and 3.40% for never users. From the weighted multivariate logistic regression models, the adjusted odds ratios of the other five smoking and vaping categories relative to never users for different covariates were calculated (Supporting information, Table S1). As shown in Table 2, compared to never users, while other smoking/vaping categories did not show any significant difference, dual users and exclusive smokers had significantly higher odds ratios of low life satisfaction with an aOR = 1.770 (95% CI: 1.135, 2.760) and an aOR = 1.452 (95% CI: 1.254, 5.896). In addition, exclusive smokers had a higher odds ratio of low life satisfaction than ex-smokers (aOR = 1.416, 95% CI: 1.095, 1.831).

Table 2. Association of life satisfaction with smoking/vaping status.

Low Life Satisfaction	Adjusted Odds Ratio (95% Confidence Interval)			
	All	Age: 18–34 (n = 7459)	Age: 35–64 (n = 24,141)	Age: 65+ (n = 15,563)
Dual users vs. Never users	1.770 (1.135, 2.760)	2.719 (1.254, 5.896)	1.280 (0.763, 2.146)	1.882 (0.682, 5.190)
Ex-smokers vs. Never users	1.025 (0.806, 1.303)	1.516 (0.806, 2.852)	0.915 (0.674, 1.244)	0.795 (0.535, 1.183)
Exclusive smokers vs. Never users	1.452 (1.121, 1.880)	2.457 (1.439, 4.193)	1.127 (0.829, 1.532)	1.170 (0.747, 1.832)
Current vapers who were ex-smokers vs. Never users	1.335 (0.708, 2.517)	1.334 (0.387, 4.594)	1.474 (0.679, 3.201)	0.104 (0.014, 0.786)
Exclusive vapers vs. Never users	1.230 (0.440, 3.440)	1.599 (0.544, 4.701)	0.190 (0.016, 2.206)	0.013 (0.001, 0.161)
Exclusive smokers vs. Ex-smokers	1.416 (1.095, 1.831)	1.621 (0.788, 3.332)	1.231 (0.908, 1.670)	1.471 (0.947, 2.285)

Note: *p*-values below 0.05 are in bold.

As shown in Table 2, similar to all adults, in the age group 18–34, dual users and exclusive smokers had higher odds ratios of low life satisfaction than never users, with an aOR = 2.719 (95% CI: 1.254, 5.896) and an aOR = 2.457 (95% CI: 1.439, 4.193), respectively. Compared to never users, in age group 65+, exclusive vapers and current vapers who were ex-smokers had significantly lower odds ratios of low life satisfaction, with an aOR = 0.013 (95% CI: 0.001, 0.161) and an aOR = 0.104 (95% CI: 0.014, 0.786). However, the sample size for these categories was very small. For example, there were 62 subjects for current vapers who were ex-smokers and in the age group 65+, and only two subjects for exclusive vapers in the age group 65+ (Supporting information, Table S2).

3.3. Association of Social/Emotional Support with Smoking/Vaping Status

To understand if social and emotional support might be associated with smoking/vaping status, we calculated the odds ratios of having low social/emotional support between different smoking/vaping categories. As shown in Table 3, compared to never users, only exclusive cigarette smokers showed a significantly higher odds ratio of low social/emotional support, with an aOR = 1.193 (95% CI: 1.030, 1.381). Moreover, exclusive smokers showed a significantly higher odds ratio of low social/emotional support than ex-smokers (aOR = 1.279, 95% CI: 1.097, 1.492). In addition, we examined the association of social/emotional support with smoking/vaping status in different age groups. Exclusive smokers in the age group 65+ showed a higher odds ratio of low social/emotional support than never users (aOR = 1.385, 95% CI: 1.061, 1.808). Compared to ex-smokers, exclusive smokers in the age

groups 18–34 and 65+ showed a higher odds ratio of low social/emotional support, with an aOR = 1.542 (95% CI: 1.003, 2.372) and an aOR = 1.470 (95% CI: 1.125, 1.920). Compared to never users, exclusive vapers age 65+ (only two subjects) showed a lower odds ratio of low social/emotional support (aOR = 0.014, 95% CI: 0.003, 0.071).

Table 3. Association of social/emotional support with smoking/vaping status.

Low social/Emotional Support	Adjusted Odds Ratio (95% Confidence Interval)			
	All	Age: 18–34 (n = 7459)	Age: 35–64 (n = 24,141)	Age: 65+ (n = 15,563)
Dual users vs. Never users	1.075 (0.768, 1.505)	0.923 (0.498, 1.710)	1.169 (0.801, 1.708)	1.061 (0.563, 2.000)
Ex-smokers vs. Never users	0.933 (0.835, 1.042)	0.806 (0.557, 1.165)	0.944 (0.809, 1.102)	0.942 (0.801, 1.108)
Exclusive smokers vs. Never users	1.193 (1.030, 1.381)	1.243 (0.909, 1.698)	1.100 (0.906, 1.335)	1.385 (1.061, 1.808)
Current vapers who were ex-smokers vs. Never users	1.148 (0.713, 1.847)	0.939 (0.437, 2.017)	1.354 (0.697, 2.629)	0.896 (0.306, 2.626)
Exclusive vapers vs. Never users	0.762 (0.412, 1.407)	0.757 (0.390, 1.466)	0.708 (0.209, 2.397)	0.014 (0.003, 0.071)
Exclusive smokers vs. Ex-smokers	1.279 (1.097, 1.492)	1.542 (1.003, 2.372)	1.165 (0.949, 1.430)	1.470 (1.125, 1.920)

Note: *p*-values below 0.05 are in bold.

4. Discussion

This study aimed to examine possible cross-sectional associations between smoking/vaping status and life satisfaction and social/emotional support among US adults. Our results showed that dual users and exclusive smokers were more likely to have low life satisfaction than never users, especially among young adults (age group 18–34). In contrast, vapers in the age group 65+ were more likely to be satisfied with their life, but the sample size was relatively small. Exclusive smokers were more likely to have low life satisfaction than ex-smokers. Compared to never users, exclusive smokers were more likely to have low social/emotional support, especially among the elders (age group 65+). Exclusive smokers were more likely to have lower social/emotional support than ex-smokers among age groups 18–34 and 65+.

In this study, we showed that, compared to never users, smokers and dual users had significantly higher odds of low life satisfaction among young adults but not among old adults. Previous studies showed that low life satisfaction is significantly associated with smoking in adolescents [22,43]. Hoping to improve their life quality, low life satisfaction might motivate individuals (especially young adults) to engage in health risk-related behaviors, such as tobacco use, alcohol, and drug abuse [44]. On the other hand, these health risk behaviors might lead to low life satisfaction [45]. However, our cross-sectional study does not determine the causal direction between low life satisfaction and smoking or vaping. In addition, our results showed that exclusive smokers were more likely to have low life satisfaction than ex-smokers. Several studies showed that ex-smokers were much happier with their lives or had a better quality of life than when they were smokers [46,47], supporting a notion that quitting smoking could improve life satisfaction. Our study showed that vapers (exclusive vapers and current vapers who were ex-smokers) had lower odds of low life satisfaction than never users, but the differences were not statistically significant. A lack of association of vaping with life satisfaction could be due to either the relatively small sample size or the possibility that there is no association between vaping with life satisfaction. The association between vaping and life satisfaction should be further validated in the future.

Our results showed that current smokers received significantly lower social and emotional support than never users, especially among the age group 65+. More importantly, current smokers had lower social and emotional support than ex-smokers. These results might suggest that low social and emotional support might be related to smoking. More social and emotional support might help with smoking cessation, which is consistent with previous findings [29,48–50]. A proposed model suggests that social and emotional

support might help with the smoking cessation by alleviating stress (also known as stress buffering) [51,52]. When facing daily stress or withdrawal symptoms, with social and emotional support, one does not need to seek substance use for coping with stresses and therefore continue the smoking cessation process [53]. It has been shown that online help is popular for providing social and emotional support for smokers [49,54]. While it was reported that most smokers have attempted to quit smoking on their own [55], social and emotional support might be an effective strategy to help this process, such as social support from their family members, especially their romantic partners [56]. With the development of the internet and electronic technologies, social and emotional support from friends and peers through social media (such as Twitter, Reddit, and Facebook) might be another effective approach for smoking cessation, which awaits further exploration and validation in the future [57]. While several social and emotional support approaches (such as friends, family members, and the community) could help with smoking/vaping cessation, it is important to determine which ones might be more effective in further studies. In our study, we did not observe a significant association of social/emotional support with current vaping status, which could be partially due to the relatively small sample size. Another possible explanation is that unlike smoking, vaping is not associated with social/emotional support. Considering the current vaping epidemic [58], it is important to determine its potential association with social/emotional support in future studies with a large sample size.

There were several limitations to our study. First, some of the smoking and vaping groups in our study had a relatively small sample size. An example would be the exclusive vapers and current vapers who were ex-smokers, especially in the old age group (65+). Therefore, the associations of life satisfaction and social/emotional support with vaping were not conclusive, especially among the youth, which requires further investigation. Second, our study was based on cross-sectional data, which did not determine the causal effect between life satisfaction or social/emotional support and smoking/vaping status. We also could not determine the longitudinal change in life satisfaction when smokers quit smoking. Third, some other important information about smoking and vaping (such as if they ever attempted to quit smoking, if there are current smokers or vapers in their family/friends, the number of cigarettes per day, the number of days used in the past month, and how long they have been smoking or vaping) were not included in our models, due to the unavailability of these data in the BRFSS survey, which might introduce some biases into our final results. In addition, in this study, there was no information on other tobacco product use, such as cigars or the waterpipe in the BRFSS data. Thus, our study only considered smoking and vaping status, which could bias our results. There was a possibility that some participants might have answered both the 2016 and 2017 BRFSS survey questionnaire. However, we were not able to identify those participants due to the anonymous property of the BRFSS survey, which might bring some bias to our analysis results. Finally, in this study we grouped survey participants into three large age groups. However, the level of life satisfaction and social/emotional support depended on the age [59,60]. Even with the same age group (such as age group 18–34), participants with different ages might have different levels of life satisfaction and social/emotional support [59]. Therefore, our age grouping method might introduce some biases.

5. Conclusions

In this study, using the US national survey data with 47,163 subjects, we showed that adult smokers are more likely to have low life satisfaction and low social/emotional support compared to never users or ex-smokers, suggesting the potential role of life satisfaction and social/emotional support in the current smoking status. Together, our study provided some preliminary but valuable information on potential smoking maintenance, which could be sources for interventions for smoking or even vaping cessation.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/ijerph191710722/s1>, Table S1: The estimated adjusted odds ratios of smoking and vaping status for covariates; Table S2: Number of adult participants in each smoking/vaping category and age group.

Author Contributions: Conceptualization, Z.X. and D.L.; methodology, Z.X.; software, Z.X.; validation, Z.X.; formal analysis, Z.X. and D.L.; investigation, Z.X.; resources, Z.X.; data curation, Z.X.; writing—original draft preparation, Z.X. and D.L.; writing—review and editing, Z.X., F.C.-B., P.C. and D.L.; visualization, Z.X. All authors have read and agreed to the published version of the manuscript.

Funding: Research reported in this publication was supported by the National Cancer Institute of the National Institutes of Health (NIH) and the Food and Drug Administration (FDA) Center for Tobacco Products under Award Number U54CA228110.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The 2016 and 2017 BRFSS data are publicly available from the CDC website and can be downloaded from https://www.cdc.gov/brfss/annual_data/annual_data.htm (accessed on 16 February 2021).

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

- World Health Organization. 2022. Available online: <https://www.who.int/news-room/fact-sheets/detail/tobacco> (accessed on 11 August 2022).
- Cornelius, M.E.; Wang, T.W.; Jamal, A.; Loretan, C.G.; Neff, L.J. Tobacco Product Use among Adults—United States, 2019. *MMWR Morb. Mortal Wkly. Rep.* **2020**, *69*, 1736–1742. [[CrossRef](#)] [[PubMed](#)]
- Creamer, M.R.; Wang, T.W.; Babb, S.; Cullen, K.A.; Day, H.; Willis, G.; Jamal, A.; Neff, L. Tobacco product use and cessation indicators among adults—United States, 2018. *MMWR Morb. Mortal Wkly. Rep.* **2019**, *68*, 1013–1019. [[CrossRef](#)] [[PubMed](#)]
- Cullen, K.A.; Gentzke, A.S.; Sawdey, M.D.; Chang, J.T.; Anic, G.M.; Wang, T.W.; Creamer, M.R.; Jamal, A.; Ambrose, B.K.; King, B.A. e-Cigarette use among youth in the United States, 2019. *JAMA* **2019**, *322*, 2095–2103. [[CrossRef](#)]
- Li, D.; Sundar, I.K.; McIntosh, S.; Ossip, D.J.; Goniewicz, M.L.; O'Connor, R.J.; Rahman, I. Association of smoking and electronic cigarette use with wheezing and related respiratory symptoms in adults: Cross-sectional results from the Population Assessment of Tobacco and Health (PATH) study, wave 2. *Tob. Control* **2020**, *29*, 140–147. [[CrossRef](#)] [[PubMed](#)]
- Li, D.; Xie, Z. Cross-sectional association of lifetime electronic cigarette use with wheezing and related respiratory symptoms in U.S. Adults. *Nicotine Tob. Res.* **2020**, *22*, S85–S92. [[CrossRef](#)]
- Wills, T.A.; Pagano, I.; Williams, R.J.; Tam, E.K. E-cigarette use and respiratory disorder in an adult sample. *Drug Alcohol Depend.* **2019**, *194*, 363–370. [[CrossRef](#)] [[PubMed](#)]
- Bhatta, D.N.; Glantz, S.A. Association of E-cigarette use with respiratory disease among adults: A longitudinal analysis. *Am. J. Prev. Med.* **2020**, *58*, 182–190. [[CrossRef](#)]
- Qasim, H.; Karim, Z.A.; Rivera, J.O.; Khasawneh, F.T.; Alshbool, F.Z. Impact of Electronic Cigarettes on the Cardiovascular System. *J. Am. Heart Assoc.* **2017**, *6*, e006353. [[CrossRef](#)]
- Xie, Z.; Ossip, D.J.; Rahman, I.; Li, D. Use of electronic cigarettes and self-reported chronic obstructive pulmonary disease diagnosis in adults. *Nicotine Tob. Res.* **2020**, *22*, 1155–1161. [[CrossRef](#)]
- Patten, S.B. Vaping and Mental Health. *J. Can. Acad. Child. Adolesc. Psychiatry* **2021**, *30*, 3–5.
- Javed, S.; Usmani, S.; Sarfraz, Z.; Sarfraz, A.; Hanif, A.; Firoz, A.; Baig, R.; Sharath, M.; Walia, N.; Cherrez-Ojeda, I.; et al. A scoping review of vaping, E-cigarettes and mental health impact: Depression and suicidality. *J. Community Hosp. Intern. Med. Perspect* **2022**, *12*, 33–39. [[CrossRef](#)] [[PubMed](#)]
- Kastaun, S.; Brose, L.S.; Scholz, E.; Viechtbauer, W.; Kotz, D. Mental health symptoms and associations with tobacco smoking, dependence, motivation, and attempts to quit: Findings from a population survey in germany (DEBRA Study). *Eur. Addict. Res.* **2022**, *28*, 287–296. [[CrossRef](#)]
- Farrell, K.R.; Weitzman, M.; Karey, E.; Lai, T.K.Y.; Gordon, T.; Xu, S. Passive exposure to E-cigarette emissions is associated with worsened mental health. *BMC Public Health* **2022**, *22*, 1138. [[CrossRef](#)] [[PubMed](#)]
- Cross, A.J.; Anthenelli, R.; Li, X. Metabotropic glutamate receptors 2 and 3 as targets for treating nicotine addiction. *Biol. Psychiatry* **2018**, *83*, 947–954. [[CrossRef](#)] [[PubMed](#)]
- Riehm, K.E.; Young, A.S.; Feder, K.A.; Krawczyk, N.; Tormohlen, K.N.; Pacek, L.R.; Mojtabai, R.; Crum, R.M. Mental health problems and initiation of E-cigarette and combustible cigarette use. *Pediatrics* **2019**, *144*, e20182935. [[CrossRef](#)]

17. Becker, T.D.; Arnold, M.K.; Ro, V.; Martin, L.; Rice, T.R. Systematic review of electronic cigarette use (vaping) and mental health comorbidity among adolescents and young adults. *Nicotine Tob. Res.* **2021**, *23*, 415–425. [\[CrossRef\]](#) [\[PubMed\]](#)
18. Weinberger, A.H.; Zhu, J.; Barrington-Trimis, J.L.; Wyka, K.; Goodwin, R.D. Cigarette use, E-cigarette use, and dual product use are higher among adults with serious psychological distress in the United States: 2014–2017. *Nicotine Tob. Res.* **2020**, *22*, 1875–1882. [\[CrossRef\]](#)
19. Grant, J.E.; Lust, K.; Fridberg, D.J.; King, A.C.; Chamberlain, S.R. E-cigarette use (vaping) is associated with illicit drug use, mental health problems, and impulsivity in university students. *Ann. Clin. Psychiatry* **2019**, *31*, 27–35.
20. Ruggeri, K.; Garcia-Garzon, E.; Maguire, A.; Matz, S.; Huppert, F.A. Well-being is more than happiness and life satisfaction: A multidimensional analysis of 21 countries. *Health Qual. Life Outcomes* **2020**, *18*, 192. [\[CrossRef\]](#)
21. Margolis, S.; Schwitzgebel, E.; Ozer, D.J.; Lyubomirsky, S. A new measure of life satisfaction: The riverside life satisfaction scale. *J. Pers. Assess.* **2019**, *101*, 621–630. [\[CrossRef\]](#)
22. Lew, D.; Xian, H.; Qian, Z.; Vaughn, M.G. Examining the relationships between life satisfaction and alcohol, tobacco and marijuana use among school-aged children. *J. Public Health* **2019**, *41*, 346–353. [\[CrossRef\]](#) [\[PubMed\]](#)
23. Maccagnan, A.; Taylor, T.; White, M.P. Valuing the relationship between drug and alcohol use and life satisfaction: Findings from the crime survey for England and Wales. *J. Happiness Stud.* **2020**, *21*, 877–898. [\[CrossRef\]](#)
24. Claydon, E.A.; DeFazio, C.; Lilly, C.L.; Zullig, K.J. Life satisfaction among a clinical eating disorder population. *J. Eat. Disord* **2020**, *8*, 53. [\[CrossRef\]](#) [\[PubMed\]](#)
25. Farris, S.G.; Zvolensky, M.J.; Schmidt, N.B. Difficulties with emotion regulation and psychopathology interact to predict early smoking cessation lapse. *Cognit. Ther. Res.* **2016**, *40*, 357–367. [\[CrossRef\]](#)
26. Uchino, B.N.; Bowen, K.; Kent de Grey, R.; Mikel, J.; Fisher, E.B. *Social Support. and Physical Health: Models, Mechanisms, and Opportunities*; Springer: New York, NY, USA, 2018.
27. Ioakeimidis, N.; Vlachopoulos, C.; Katsi, V.; Tousoulis, D. Smoking cessation strategies in pregnancy: Current concepts and controversies. *Hellenic J. Cardiol* **2019**, *60*, 11–15. [\[CrossRef\]](#) [\[PubMed\]](#)
28. Twyman, L.; Cowles, C.; Walsberger, S.C.; Baker, A.L.; Bonevski, B.; Tackling Tobacco Mental Health Advisory Group. ‘They’re going to smoke anyway’: A qualitative study of community mental health staff and consumer perspectives on the role of social and living environments in tobacco use and cessation. *Front. Psychiatry* **2019**, *10*, 503. [\[CrossRef\]](#)
29. van den Brand, F.A.; Nagtzaam, P.; Nagelhout, G.E.; Winkens, B.; van Schayck, C.P. The association of peer smoking behavior and social support with quit success in employees who participated in a smoking cessation intervention at the workplace. *Int. J. Environ. Res. Public Health* **2019**, *16*, 2831. [\[CrossRef\]](#)
30. Liao, Y.; Wu, Q.; Kelly, B.C.; Zhang, F.; Tang, Y.Y.; Wang, Q.; Ren, H.; Hao, Y.; Yang, M.; Cohen, J.; et al. Effectiveness of a text-messaging-based smoking cessation intervention (“Happy Quit”) for smoking cessation in China: A randomized controlled trial. *PLoS Med.* **2018**, *15*, e1002713. [\[CrossRef\]](#)
31. de Dios, M.A.; Stanton, C.A.; Cano, M.A.; Lloyd-Richardson, E.; Niaura, R. The influence of social support on smoking cessation treatment adherence among HIV+ smokers. *Nicotine Tob Res.* **2016**, *18*, 1126–1133. [\[CrossRef\]](#)
32. Yun, E.H.; Kang, Y.H.; Lim, M.K.; Oh, J.K.; Son, J.M. The role of social support and social networks in smoking behavior among middle and older aged people in rural areas of South Korea: A cross-sectional study. *BMC Public Health* **2010**, *10*, 78. [\[CrossRef\]](#)
33. Stallings-Smith, S.; Ballantyne, T. Ever use of E-cigarettes among adults in the United States: A cross-sectional study of sociodemographic factors. *Inquiry* **2019**, *56*, 46958019864479. [\[CrossRef\]](#)
34. Iachan, R.; Pierannunzi, C.; Healey, K.; Greenlund, K.J.; Town, M. National weighting of data from the Behavioral Risk Factor Surveillance System (BRFSS). *BMC Med. Res. Methodol.* **2016**, *16*, 155. [\[CrossRef\]](#) [\[PubMed\]](#)
35. Osei, A.D.; Mirbolouk, M.; Orimoloye, O.A.; Dzaye, O.; Uddin, S.M.I.; Benjamin, E.J.; Hall, M.E.; DeFilippis, A.P.; Bhatnagar, A.; Biswal, S.S.; et al. Association between E-cigarette use and chronic obstructive pulmonary disease by smoking status: Behavioral risk factor surveillance system 2016 and 2017. *Am. J. Prev Med.* **2020**, *58*, 336–342. [\[CrossRef\]](#) [\[PubMed\]](#)
36. Atuegwu, N.C.; Oncken, C.; Laubenbacher, R.C.; Perez, M.F.; Mortensen, E.M. Factors associated with E-cigarette use in U.S. young adult never smokers of conventional cigarettes: A machine learning approach. *Int. J. Environ. Res. Public Health* **2020**, *17*, 7271. [\[CrossRef\]](#)
37. Bircan, E.; Bezirhan, U.; Porter, A.; Fagan, P.; Orloff, M.S. Electronic cigarette use and its association with asthma, chronic obstructive pulmonary disease (COPD) and asthma-COPD overlap syndrome among never cigarette smokers. *Tob. Induc. Dis.* **2021**, *19*, 23. [\[CrossRef\]](#)
38. Al Rifai, M.; Mirbolouk, M.; Obisesan, O.H.; Jia, X.; Nasir, K.; Merchant, A.T.; Blaha, M.; Virani, S. The association of electronic cigarette use and the subjective domains of physical and mental health: The behavioral risk factor surveillance system survey. *Cureus* **2020**, *12*, e7088. [\[CrossRef\]](#) [\[PubMed\]](#)
39. Xie, Z.; Ossip, D.J.; Rahman, I.; O’Connor, R.J.; Li, D. Electronic cigarette use and subjective cognitive complaints in adults. *PLoS ONE* **2020**, *15*, e0241599. [\[CrossRef\]](#)
40. Bursac, Z.; Gauss, C.H.; Williams, D.K.; Hosmer, D.W. Purposeful selection of variables in logistic regression. *Source Code Biol. Med.* **2008**, *3*, 17. [\[CrossRef\]](#) [\[PubMed\]](#)
41. Bernstein, S.F.; Rehkopf, D.; Tuljapurkar, S.; Horvitz, C.C. Poverty dynamics, poverty thresholds and mortality: An age-stage Markovian model. *PLoS ONE* **2018**, *13*, e0195734. [\[CrossRef\]](#)

42. Lo, C.C.; Cheng, T.C.; Simpson, G.M. Marital status and work-related health limitation: A longitudinal study of young adult and middle-aged Americans. *Int. J. Public Health* **2016**, *61*, 91–100. [\[CrossRef\]](#)
43. Heshmat, R.; Qorbani, M.; Safiri, S.; Eslami-Shahr Babaki, A.; Matin, N.; Motamed-Gorji, N.; Motlagh, M.E.; Djalalinia, S.; Ardalan, G.; Mansourian, M.; et al. Association of passive and active smoking with self-rated health and life satisfaction in Iranian children and adolescents: The CASPIAN IV study. *BMJ Open* **2017**, *7*, e012694. [\[CrossRef\]](#)
44. Kroencke, L.; Kuper, N.; Bleidorn, W.; Denissen, J.J.A. How does substance use affect personality development? Disentangling between- and within-person effects. *Soc. Psychol. Pers. Sci.* **2021**, *12*, 517–527. [\[CrossRef\]](#)
45. Stenlund, S.; Koivumaa-Honkanen, H.; Sillanmaki, L.; Lagstrom, H.; Rautava, P.; Suominen, S. Health behavior of working-aged Finns predicts self-reported life satisfaction in a population-based 9-years follow-up. *BMC Public Health* **2021**, *21*, 1815. [\[CrossRef\]](#) [\[PubMed\]](#)
46. Shahab, L.; West, R. Do ex-smokers report feeling happier following cessation? Evidence from a cross-sectional survey. *Nicotine Tob. Res.* **2009**, *11*, 553–557. [\[CrossRef\]](#) [\[PubMed\]](#)
47. Piper, M.E.; Kenford, S.; Fiore, M.C.; Baker, T.B. Smoking cessation and quality of life: Changes in life satisfaction over 3 years following a quit attempt. *Ann. Behav. Med.* **2012**, *43*, 262–270. [\[CrossRef\]](#) [\[PubMed\]](#)
48. Soulakova, J.N.; Tang, C.Y.; Leonardo, S.A.; Taliaferro, L.A. Motivational Benefits of Social Support and Behavioural Interventions for Smoking Cessation. *J. Smok. Cessat.* **2018**, *13*, 216–226. [\[CrossRef\]](#)
49. Qian, Y.; Gui, W.; Ma, F.; Dong, Q. Exploring features of social support in a Chinese online smoking cessation community: A multidimensional content analysis of user interaction data. *Health Inform. J.* **2021**, *27*, 14604582211021472. [\[CrossRef\]](#)
50. Ochsner, S.; Knoll, N.; Stadler, G.; Luszczynska, A.; Hornung, R.; Scholz, U. Interacting effects of receiving social control and social support during smoking cessation. *Ann. Behav. Med.* **2015**, *49*, 141–146. [\[CrossRef\]](#)
51. Mayne, S.L.; Auchincloss, A.H.; Moore, K.A.; Michael, Y.L.; Tabb, L.P.; Echeverria, S.E.; Diez Roux, A.V. Cross-sectional and longitudinal associations of neighbourhood social environment and smoking behaviour: The multiethnic study of atherosclerosis. *J. Epidemiol. Community Health* **2017**, *71*, 396–403. [\[CrossRef\]](#)
52. Westmaas, J.L.; Bontemps-Jones, J.; Bauer, J.E. Social support in smoking cessation: Reconciling theory and evidence. *Nicotine Tob. Res.* **2010**, *12*, 695–707. [\[CrossRef\]](#)
53. Creswell, K.G.; Cheng, Y.; Levine, M.D. A test of the stress-buffering model of social support in smoking cessation: Is the relationship between social support and time to relapse mediated by reduced withdrawal symptoms? *Nicotine Tob. Res.* **2015**, *17*, 566–571. [\[CrossRef\]](#) [\[PubMed\]](#)
54. Graham, A.L.; Amato, M.S. Twelve million smokers look online for smoking cessation help annually: Health information national trends survey data, 2005–2017. *Nicotine Tob. Res.* **2019**, *21*, 249–252. [\[CrossRef\]](#)
55. Larabie, L.C. To what extent do smokers plan quit attempts? *Tob. Control* **2005**, *14*, 425–428. [\[CrossRef\]](#) [\[PubMed\]](#)
56. Whitton, S.W.; McLeish, A.C.; Godfrey, L.M.; James-Kangal, N.; Rhoades, G.K. Partner assisted smoking cessation treatment: A randomized clinical trial. *Subst. Use Misuse* **2020**, *55*, 1228–1236. [\[CrossRef\]](#) [\[PubMed\]](#)
57. Thrul, J.; Tormohlen, K.N.; Meacham, M.C. Social media for tobacco smoking cessation intervention: A review of the literature. *Curr. Addict. Rep.* **2019**, *6*, 126–138. [\[CrossRef\]](#)
58. *Surgeon General's Advisory on E-Cigarette Use among Youth*; Centers for Disease Control and Prevention: Atlanta, GA, USA, 2018.
59. Blanchflower, D.G. Is happiness U-shaped everywhere? Age and subjective well-being in 145 countries. *J. Popul. Econ.* **2021**, *34*, 575–624. [\[CrossRef\]](#)
60. Franssen, T.; Stijnen, M.; Hamers, F.; Schneider, F. Age differences in demographic, social and health-related factors associated with loneliness across the adult life span (19–65 years): A cross-sectional study in the Netherlands. *BMC Public Health* **2020**, *20*, 1118. [\[CrossRef\]](#)