

Table S6. Descriptive data from the OHIP dimensions orofacial appearance and psychosocial impact from the four dimensional OHIP scale. SD: Standard deviation.

OHIP Dimension		Orofacial Appearance						Psychosocial Impact					
Time of Assessment		T0		T1		T2		T0		T1		T2	
Parameter	Groups	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Total	-	10.0	5.7	7.3	4.9	6.2	5.3	22.8	12.3	17.1	13.3	15.6	13.2
Stage	I	9.6	5.5	6.2	3.9	4.9	4.0	21.9	16.0	15.8	12.4	13.1	10.8
	II	11.0	6.1	9.8	6.2	9.1	6.9	24.8	17.5	20.0	15.4	21.5	16.5
Pain	no	10.1	4.8	6.5	4.1	7.0	4.9	20.0	14.2	15.1	10.5	16.1	11.2
	yes	10.0	6.3	7.9	5.5	5.6	5.7	24.9	17.7	18.5	15.1	15.3	14.7
Primary disease	osteoporosis	10.7	6.7	6.8	5.5	5.6	5.0	22.8	17.0	14.5	11.3	15.1	9.2
	malignoma	9.9	5.4	7.5	4.8	6.3	5.5	22.8	16.4	17.9	13.9	15.8	14.3
Risk evaluation	low risk	11.6	7.6	7.1	6.5	6.7	6.6	27.5	21.1	17.5	16.5	19.9	17.2
	high risk	9.3	4.4	7.4	4.1	5.9	4.7	20.5	13.2	16.9	11.8	13.5	10.4
Duration of intake	short	10.3	5.2	7.7	4.4	6.2	5.2	25.6	15.7	19.7	14.0	14.9	12.2
	long	20.9	16.8	7.0	5.4	6.1	5.6	20.8	16.8	15.2	12.8	16.2	14.1
Defect size	small	10.5	5.9	6.7	4.2	5.5	3.8	23.7	15.6	16.8	11.8	15.3	11.2
	medium	9.4	5.4	7.5	5.0	6.0	6.2	19.3	14.8	14.2	12.9	13.5	12.0
	large	10.1	5.9	7.6	5.7	6.9	5.9	24.9	18.5	19.6	15.1	17.6	15.9
Need for prosthodontics	yes	11.2	5.7	8.2	5.6	6.9	6.8	26.5	16.2	18.6	13.9	18.1	16.1
	no	8.9	5.5	6.5	4.2	5.4	3.7	19.5	16.0	15.8	13.0	13.4	9.8