

Table S5. Descriptive data from the OHIP dimensions oral function and orofacial pain from the four dimensional OHIP scale. SD: Standard deviation.

OHIP Dimension		Oral Function						Orofacial Pain					
Time of Assessment		T0		T1		T2		T0		T1		T2	
Parameter	Groups	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Total	-	13.3	8.9	12.5	8.4	10.2	7.8	13.1	6.7	8.2	4.9	6.9	5.3
Stage	I	12.5	8.8	11.1	7.6	8.6	7.1	13.1	5.8	8.1	4.0	5.8	4.9
	II	14.9	9.4	15.7	9.3	14.0	8.1	13.4	8.8	8.3	6.7	9.6	5.4
Pain	no	12.9	7.7	10.4	6.6	10.2	7.3	14.1	6.1	7.9	4.6	8.3	6.1
	yes	13.5	9.9	14.0	9.4	10.2	8.3	12.6	7.2	8.4	5.2	5.9	4.5
Primary disease	osteoporosis	14.6	11.2	12.3	9.3	9.6	9.1	11.1	6.3	5.8	4.2	5.1	4.6
	malignoma	12.9	8.4	12.6	8.3	10.4	7.5	13.8	6.8	8.6	4.9	7.5	5.4
Risk evaluation	low risk	16.6	11.8	14.7	9.8	12.5	10.0	12.5	6.6	7.5	6.5	6.8	5.2
	high risk	11.6	6.7	11.9	7.8	9.1	6.3	13.5	6.9	8.5	4.0	7.0	5.4
Duration of intake	short	15.0	6.1	13.9	8.1	10.7	6.9	12.1	6.5	7.9	4.9	6.9	4.6
	long	12.0	10.4	11.6	8.7	9.9	8.5	14.0	6.7	11.6	8.7	9.9	8.5
Defect size	small	14.0	10.0	12.5	8.6	10.9	7.4	11.8	4.6	8.5	3.8	6.6	3.1
	medium	12.7	8.4	9.6	9.7	7.7	7.5	12.4	7.0	7.7	4.5	7.5	5.8
	large	13.1	9.1	14.8	7.0	11.6	8.3	14.9	7.9	8.3	6.1	6.7	6.5
Need for prosthodontics	yes	15.6	9.9	14.4	9.0	12.8	8.4	13.4	6.7	7.8	5.6	6.8	6.2
	no	11.2	7.6	10.8	7.7	7.9	6.5	13.1	6.9	8.5	4.2	7.1	4.5