



Article

Enhanced Biodegradation of Phenylurea Herbicides by *Ochrobactrum anthropi* CD3 Assessment of Its Feasibility in Diuron-Contaminated Soils

Lara-Moreno Alba , Morillo Esmeralda and Villaverde Jaime *

Department of Agrochemistry, Environmental Microbiology and Soil Conservation, Institute of Natural Resources and Agrobiological of Seville, Spanish National Research Council (IRNAS-CSIC), 41012 Seville, Spain; lara@irnas.csic.es (L.-M.A.); morillo@irnas.csic.es (M.E.)

* Correspondence: j.villaverde@csic.es

Abstract: The phenylurea herbicides are persistent in soil and water, making necessary the development of techniques for their removal from the environment. To identify new options in this regard, bacterial strains were isolated from a soil historically managed with pesticides. *Ochrobactrum anthropi* CD3 showed the ability to remove completely herbicides such as diuron, linuron, chlorotoluron and fluometuron from aqueous solution, and up to 89% of isoproturon. In the case of diuron and linuron, their main metabolite, 3,4-dichloroaniline (3,4-DCA), which has a higher toxicity than the parent compounds, was formed, but remained in solution without further degradation. *O. anthropi* CD3 was also tested for bioremediation of two different agricultural soils artificially contaminated with diuron, employing bioremediation techniques: (i) biostimulation, using a nutrient solution (NS), (ii) bioaugmentation, using *O. anthropi* CD3, and (iii) bioavailability enhancement using 2-hydroxypropyl- β -cyclodextrin (HPBCD). When bioaugmentation and HPBCD were jointly applied, 50% of the diuron initially added to the soil was biodegraded in a range from 4.7 to 0.7 d. Also, 3,4-DCA was degraded in soil after the strain was inoculated. At the end of the soil biodegradation assay an ecotoxicity test confirmed that after inoculating *O. anthropi* CD3 the toxicity was drastically reduced.

Keywords: phenylurea herbicides; 3,4-dichloroaniline; soil; bioremediation; bioaugmentation; biostimulation; 2-hydroxypropyl- β -cyclodextrin



Citation: Alba, L.-M.; Esmeralda, M.; Jaime, V. Enhanced Biodegradation of Phenylurea Herbicides by *Ochrobactrum anthropi* CD3 Assessment of Its Feasibility in Diuron-Contaminated Soils. *Int. J. Environ. Res. Public Health* **2022**, *19*, 1365. <https://doi.org/10.3390/ijerph19031365>

Academic Editor: Andrew S. Hursthouse

Received: 29 December 2021

Accepted: 22 January 2022

Published: 26 January 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Over the few last decades, the use of pesticides in agriculture has become more extended. Studies have proved the presence of high levels of pesticides in soils, leachates and river water, higher than the limit values established by the local, national and European legislation [1]. Phenylurea herbicides are among the most widely pesticides used in agriculture worldwide [2]. They are used for pre- and post-emergence pest control in different crops such as cotton, cereals, and fruits [3]. These herbicides act as photosynthesis inhibitors by blocking electron transfer at photosystem II resulting in necrosis and plant death [4].

Their solubilities in water range generally from moderately to relatively high and they are expected to have low to moderate adsorptivity to soil, based on their octanol-water partition coefficient, K_{ow} [5] (Table S1). This makes them mobile in soils and, once applied to the target field, irrigation and rainfall can deliver them through runoff and leaching from the ground to rivers, groundwaters, seawater and lakes [6]. However, the properties of phenylurea herbicides (Table S1) significantly affect the possibility of their adsorption in soils, and, consequently, their mobility and their possibility of reaching water environments. Blondel et al. and Agbaogun et al. [7,8] observed that the K_{ow} values of these herbicides were as high as the number of chlorine atoms on the phenyl group, increasing their adsorption to soils. Consequently, diuron and linuron, with two chlorine

atoms, present the highest adsorption to soils and are considered only as slightly mobile, and, therefore, they are the most frequently detected in contaminated soils. The rest of phenylurea herbicides are classified as mobile, being more frequently detected in water environments [9].

The degradation of phenylurea herbicides, both in water and in soil gives, as a result, the production of metabolites, such as N-(3,4-dichlorophenyl)urea (DCPU), N-(3,4-dichlorophenyl)-N-methylurea (DCPMU) and 3,4-dichloroaniline (3,4-DCA), which are more toxic than their parent contaminant, and together with them can lead to serious environmental and public health problems [10]. For these reasons, many phenylurea herbicides have been banned in European countries [11], where their presence in soils and water has to be progressively reduced, but they continue to be used in numerous regions of the world. Therefore, the removal of these herbicides from the environment is a central issue.

In the last decades, eco-friendly methods to remediate polluted environments have emerged. One of them is the use of microorganisms which can remove a wide variety of organic pollutants and can adapt to different bleak environments [12]. However, the natural degradation of these herbicides in soil is relatively slow. One of the reasons could be the insufficient number of endogenous degrading microorganisms specific for these herbicides, or that the endogenous microbiota does not have the catabolic capacities to metabolise and effectively degrade the target contaminants [13]. With the aim of improving degradation in soil, studies have been focused on isolation and identification of different specific degrading microbial populations from adapted soils which have showed capabilities to eliminate these organic contaminants from environments [13,14].

Diuron biodegradation in aqueous solution through microbial inoculation (bioaugmentation) has been studied by several authors, inoculating different microorganisms in the form of microbial consortia [15,16] or employing pure bacterial cultures, such as, *Arthrobacter* sp. N2 [17]; *Variovorax* sp. SRS16 [18] and *Micrococcus* sp. PS [19]. However, as far as we know, few studies have been described in literature about the use of bioaugmentation for diuron biodegradation in contaminated soils. Bacterial strains belonging to the genus *Bacillus* and *Stenotrophomonas* were employed in bioaugmentation [20,21]). In another work, the effect of the fungus *Neurospora intermedia* to degrade diuron in soil was studied by Villaverde et al. [22]. The remaining studies in soils were performed by Villaverde et al., directly using the endogenous soil microbiota as degrader [23], microbial consortia isolated from different soils [24] or an artificial consortium composed of *Arthrobacter sulfonivorans*, *Variovorax soli* and *Advenella* sp. JRO [25]. However, to the best of our knowledge, the bacterial strain *O. anthropi* has never been described as a phenylurea degrader.

Although bioavailability of phenylurea herbicides is not considered as the main issue concerning biodegradation [26], the bioavailability in soils of most hydrophobic compounds is reduced with their ageing [27]. Cyclodextrins (CDs) have been suggested as an option for the removal of hydrophobic contaminants from soils because they are capable of increasing water solubility of hydrophobic organic compounds [28]. This ability to form inclusion complexes with different contaminant guest molecules has been employed in soil bioremediation processes [29] since CDs enhance the water solubility of low polarity compounds and hence, their bioavailability [30].

In this work, the strain *O. anthropi* CD3 was isolated in our laboratory, using diuron enrichment cultures, from an agricultural soil treated for decades with a variety of herbicides, including phenylureas. As previously mentioned, the species *O. anthropi* has never been described as a phenylurea degrader and degrades very high percentages of the five phenylureas studied, but, in addition, few studies have described bacterial strains belonging to this genus as pesticide degrader. *Ochrobactrum* sp. JAS2 was able to degrade 300 mg L⁻¹ of chlorpyrifos after 12 h [31] or *Ochrobactrum intermedium* G3.48 was described as diuron-degrading (33% after 120 h) [32]. In the present study, CD3 strain was tested for diuron, linuron, chlorotoluron, isoproturon and fluometuron biodegradation in solution. Furthermore, the investigated strain was inoculated in two soils artificially contaminated with diuron, with the aim of finding an effective bioremediation tool based

on biostimulation (using nutrients solution (NS)), bioaugmentation (inoculating with *O. anthropi* CD3) and the use of 2-hydroxypropyl- β -cyclodextrin (HPBCD) which is able to improve water solubility of the herbicide diuron, increasing it up to 23-fold [25], and hence, its bioavailability to be degraded. An ecotoxicology study was also carried out before and after soil diuron bioremediation by *O. anthropi* CD3 to determine if the treatment achieved the desired decrease in soil toxicity.

2. Materials and Methods

2.1. Materials

Analytical grade (99%) chlorotoluron (3-(3-chloro-4-methyl)-1,1-dimethylurea), isoproturon (3-(4-isopropylphenyl)-1,1-dimethylurea), fluometuron (1,1-dimethyl-3-[3-(trifluoromethyl)]phenylurea), linuron (3-(3,4-dichlorophenyl)-1-methoxy-1-methylurea), and the metabolite 3,4-dichloroaniline were purchased from Sigma-Aldrich (Madrid, Spain). Technical grade (98%) diuron (N-(3, 4-dichlorophenyl)-N, N-dimethylurea) was supplied by Presmar S.L. (Seville, Spain). HPBCD was provided by Cyclolab Ltd (Budapest, Hungary).

Three agricultural soil samples (LL, PLD, R) from the South of Spain were used, classified as an Alfisol, Inceptisol and Alfisol, respectively [33]. LL soil is located at Vejer de la Frontera—Cádiz (36°17'52.6" N 5°52'45.2" W), dedicated to the intensive agriculture of carrots, avocados, grapefruits, leeks, sunflowers, etc. This soil has been treated for many years with phenylurea herbicides and other organic pesticides such as organochlorine and organophosphate pesticides. PLD soil was taken from a farm located in Los Palacios y Villafranca—Seville (37°10'20.0" N 5°55'21.9" W) and has been treated with organohalogen herbicides for years, and wheat, cereals and vineyard has been grown on this soil. R soil was collected from a palm trees area in Conil de la Frontera—Cádiz (36°18'32.4" N 6°08'58.6" W) treated with of the insecticide chlorpyrifos. Soil samples were taken from the superficial horizon (0–20 cm) and were partially air-dried for 24 h and sieved (2 mm). Their properties are shown in Table 1. The pH was determined in a 1:2.5 soil/water extract. A Bouyoucos densimeter was used to evaluate the particle size distribution; OM was quantified by $K_2Cr_2O_7$ oxidation; and the total carbonate content was measured using the manometric method. The water holding capacity of the soils was 52.4%, 23.16% and 53.2% for LL, PLD and R, respectively.

Table 1. Some properties of the soils used.

Soils	pH	CO ₃ ⁻² (%)	OM (%)	Sand (%)	Silt (%)	Clay (%)	Textural Classification	Taxonomic Classification *
R	7.72	4.00	3.44	77	9.50	13.5	Sandy	Alfisol
PLD	8.24	9.70	1.67	47.0	18.5	34.5	Clay loam	Inceptisol
LL	7.84	4.00	0.87	79.6	9.30	11.1	Sandy	Alfisol

* USDA SOIL TAXONOMY, Soil maps, 2005. National Geographic Institute. Nature Database (Spanish Ministry of Environment).

2.2. Diuron Degradation Enrichment

Diuron degrader bacterial strains were isolated in our laboratory from the agricultural soil sample LL. 10 g of LL soil was added to sterilized 250 mL Erlenmeyer flasks (JP-Selecta, Barcelona, Spain) with one cycle at 120 °C, inlet pressure of 101 kPa, for 20 min with 50 mL of a mineral salt medium (sterilized MSM), which provided the macronutrients. MSM contained (g L⁻¹): 4.0 Na₂HPO₄; 2.0 KH₂PO₄; 0.8 MgSO₄; 0.8 NH₄SO₄. The aqueous solution was spiked with diuron to reach a concentration of 1 g L⁻¹, which was the only source of carbon and energy. 1 mL of a micronutrients solution (SNs), which contained: 12.5 NiCl₂ 6H₂O; 25.0 SnCl₂ 2H₂O; 12.5 ZnSO₄ 7H₂O; 12.5 Al₂(SO₄)₃ 18H₂O; 75.0 MnCl₂ 4H₂O; 12.5 CoCl₂ 2H₂O; 37.5 FeSO₄ 7H₂O; 10.0 CaSO₄ 2H₂O; 3.75 KBr; 3.75 KCl; 2.50 LiCl (mg L⁻¹) was also added [34]. The mixture of SNs and MSM (1:50) was named nutrient solution (NS). Incubation conditions were 150 rpm at 30 °C, and every 7 d (four times) an aliquot of 10 mL of the culture was removed and transferred to another flask containing the

same previously described composition. Aliquots of 100 μL of the final culture were added on agar plates, prepared with MSM medium and diuron with a concentration of 0.5 g L^{-1} . Successive isolations were performed, and different colonies, according to their size, colour, edge and elevation, were selected. Eleven strains (LL1D, LL2D, LL3D, LL4D, LL5D, LL6D, LL7D, LL8D, LL9D, LL10D and LL11D) were isolated. The selected isolated strains were kept in cryovials (Microbank[®], Pro-Lab Diagnostics, Bromborough, UK), which contained a specific culture medium and 20 porous spheres at 3 mm diameter. They were kept at $-80 \text{ }^\circ\text{C}$.

2.3. Inoculum Preparation

For each experiment, bacterial cryovials were defrosted and the bacteria were sowed in Luria–Bertani (LB) medium. The culture was incubated at $30 \text{ }^\circ\text{C}$ in an orbital shaker (150 rpm). The different bacterial strains were collected at the end of their respective exponential phase and later washed twice with a sterile MSM solution before beginning the experiments. The final density of the strains added in biodegradation assays was 10^8 CFU mL^{-1} .

2.4. Diuron Degrading Capacity of the Isolated Bacterial Strains

The capacity of the 11 isolated strains to degrade diuron in solution was tested. Each inoculum was added to NS solution containing diuron as the only carbon and nitrogen sources (10 mg L^{-1}). The samples were incubated in a thermostatic chamber at $30 \text{ }^\circ\text{C}$ for 20 d, and periodically the concentration of diuron was measured by HPLC as described in the Analytical Methods section.

2.5. Isolated Bacterial Strain Identification with 16S rDNA Amplification

The isolated strain which showed to be the most effective for diuron degradation in solution (Figure S1) was identified. DNA was extracted from the liquid cultured of the strain employing the G-spin[™] total DNA Extraction Kit (iNtRON Biotechnology, Seongnam, Korea), then 16S rRNA gene was amplified by polymerase chain reaction (PCR) using a high-fidelity polymerase (Velocity DNA polymerase from Bioline, Almería, Spain) with universal oligonucleotides primers: 16F27 (annealing at position 8–27 *E. coli* numbering) and 16R1488 [35]. Finally, the PCR product was purified using a NucleoSpin[®]PCR clean-up gel extraction kit and PCR clean-up (Macherey-Nagel, Cultek S. L., Madrid, Spain) prior to be sequenced.

2.6. Biodegradation Experiments in Solution

Phenylurea herbicide biodegradation experiments in solution were performed in 25 mL sterilized glass vials during 60 d of assay. At each incubation timepoint (1, 3, 7, 12, 21, 30 and 60 d), aliquots were taken from solutions in triplicate. Each vial contained 330 μL of the bacterial inoculum (final density of $4.5 \times 10^6 \text{ CFU mL}^{-1}$), 15 μL SNs and 15 mL of MSM, contaminated with 10 mg L^{-1} of each phenylurea herbicide. To monitor abiotic degradation, non-inoculated vials (control) were also set up. The vials were kept at a temperature of $30 \text{ }^\circ\text{C}$ in a laboratory incubator. Due to the importance and toxicity reported in the literature the main metabolite of diuron and linuron, 3,4-DCA, was analyzed at different time points. The samples were taken under sterile conditions in a vertical laminar flow cabin and filtered using a nylon filter ($0.45 \text{ } \mu\text{m}$, RephiQuik, Madrid, Spain) for analytical analysis (see Section 2.10).

2.7. Biodegradation Experiments in Soils

Diuron biodegradation experiments in soils were performed in a 250 mL sterilized flask containing 100 g of soil sample contaminated with 50 mg kg^{-1} diuron and the required volume of NS (MSM + SNs) to reach 40% of the soil water holding capacity [24]. The treatments were assigned completely at random so that each experimental unit had the same chance of receiving any one treatment. Different bioremediation strategies were

designed: (i) biostimulation: spiked soil + NS; (ii) bioaugmentation: spiked soil + NS + *O. anthropi* CD3 inoculum (1×10^8 CFU g^{-1}); (iii) bioavailability enhancing: spiked soil + HPBCD solution + NS. HPBCD was added with a concentration corresponding to 10 times that of the diuron initially spiked in the soil sample; (iv) Combined use of biostimulation, bioaugmentation and bioavailability enhancing: spiked soil + HPBCD + *O. anthropi* CD3 + NS. In parallel, abiotic controls were set up adding 200 mg L^{-1} of $HgCl_2$ to eliminate the soil endogenous microbiota in order to observe any diuron dissipation due to abiotic processes. All experiments were set at a temperature of 30 °C in a laboratory incubator for 100 d. Soil samples were taken at different times of the biodegradation assays (0, 6, 8, 13, 20, 27, 36, 47, 65, 87 and 100 d). Diuron degradation was measured through the relative decline of its residues in the soil samples. For this purpose, 1 g of each soil sample was exhaustively extracted with 5 mL of methanol. The extraction process was carried out through the following steps; (1) 24 h of agitation of the tubes in an orbital shaker at 100 rpm at 20 ± 1 °C, (2) 1 min of vortex mixer, (3) 30 min in an ultrasound bath and finally the supernatant was recovered by 10 min centrifugation at 8000 rpm. Once the samples were obtained, diuron and its metabolite 3,4-DCA were analysed by liquid chromatography as described in the Section 2.10.

2.8. Model of Biodegradation Kinetics

The biodegradation curves were fitted to a simple first-order kinetic model (SFO) and a biphasic first-order kinetic sequential model (Hockey- stick, HS) using an excel file provided by focus [36] and the Solver tool (Microsoft Statistical Package, New York, NY, USA), and the following equations:

$$[C]_t = [C]_0 e^{-kt} \text{ (SFO)}$$

$$DT_{50} = \ln 2/k \text{ (SFO)}$$

$$[C]_t = [C]_0 e^{-k_1 t_b} e^{-k_2(t-t_b)} \text{ (HS)}$$

$$DT_{50} = (\ln 100/100-50)/k_1 \text{ if } DT_{50} \leq t_b \text{ (HS)}$$

$$DT_{50} = t_b + (\ln (100/100-50) - k_1 t_b)/k_2 \text{ if } DT_{50} > t_b \text{ (HS)}$$

where $[C]_t$ and $[C]_0$ are the concentrations of diuron at time t and just at the beginning of the experiments. In SFO model, K is the degradation (d^{-1}). DT_{50} is the time required for the pollutant concentration to decline to half of its initial value. HS model has two different rate constants of degradation, k_1 and k_2 , and t_b , which is the time at which the rate constant changes. SFO and HS models were chosen for their simplicity and their capabilities to fit previously published measured dissipation kinetic datasets for herbicides [37]. The coefficient of determination and the chi-square (χ^2) test were calculated as indicators of the goodness of fit χ^2 values less than 15 imply a good fit [36].

2.9. Diuron Availability in Soil

Soil diuron extraction assays were carried out to check the effect of using NS (MSM + SNs) and HPBCD as extractants on diuron availability. One g of soil previously contaminated with 50 mg kg^{-1} of diuron was extracted with 5 mL of NS or NS combined with HPBCD (10 times the concentration of diuron initially spiked in soil). The centrifuge tubes were shaken on an orbital shaker for 72 h at 20 ± 1 °C, centrifuged (10 min, 7000 rpm) to separate the supernatant, which was filtered through a 0.22 μm glass fiber membrane (Millipore, Burlington, MA, USA). Diuron concentration was measured using the analytical methods described below.

2.10. Analytical Methods

Concentrations of the phenylurea herbicides and the metabolite 3,4-DCA during biodegradation experiments in soil and solution were analyzed on a LC-2010A HT HPLC

system equipped with a UV detector (Shimadzu Scientific Instruments, Columbia, MD, USA). Kromasil C18 reverse-phase was the chromatographic column used, the mobile phase was acetonitrile:water (60:40, *v/v*), and all the compounds were detected by UV absorbance at 230 nm. The calibration curves were linear over the concentration range studied, with correlation coefficients higher than 0.99, indicating good performance of the chromatographic method. The detection limits calculated using a noise-signal ratio of 3 were below 0.01 mg L^{-1} for the investigated compounds, and the standard deviations were around 5%. In soils, diuron extraction with methanol was effective with recovery rates from 88 ± 6 to $115 \pm 8\%$ and relative standard deviation (RSD) lower than 15%. For the acquisition and management of the data, the LC Shimadzu Solution Chromatography Data System computer program was used.

2.11. Toxicity Analysis

The Microtox[®] Test System (Modern Water Inc., New Castle, U.K.) based on the bioluminescent of the bacterium *Vibrio fischeri* was used to measure the toxicity of the soil, according to the standard protocol for the Microtox[®] basic test [38]. This toxicity test can be applied to liquid waste discharged in natural waters and also to leachates/percolates from contaminated soils. Soil toxicity was estimated through the determination of the calculated value EC_{50} of the soil lixivate, which is a hypothetical value that represents the soil extract concentration (% *v/v*) having a toxic effect that would produce 50% reduction in luminescence in *V. fischeri*. Briefly, 2 g of each soil sample was added to 3 mL of NaCl at 2% solution. These suspensions were hatched in an orbital shaker for 10 min, centrifuged for 2 min at 10,000 rpm and serially diluted (1:2) with NaCl at 2% solution (6.25%, 12.5%, 25% and 50% (*v/v*)) and compared with the control [39]. Freeze-dried *V. fischeri* were rehydrated forthwith prior to use in testing. Tests were performed in a temperature-controlled photometer at 15 °C (Microbics Inc., Lenexa, KS, USA, 1992). It was assumed that the control containing only *V. fischeri* is considered to have 0% of inhibition, and the lixivate containing only NaCl (2%) is considered to have 100% of inhibition. EC_{50} parameter is given by the Microtox[®]Text System for each sample analysed, and the toxic units (TU) were estimated using the formula $TU = 100/EC_{50}$. TU results were classified according to Persoone et al. [40] ($TU < 0.4$ no acute toxicity; $0.4 < TU < 1$ light acute toxicity; $1 < TU < 10$ acute toxicity; $10 < TU < 100$ high acute toxicity; $TU > 100$ very high acute toxicity).

2.12. Statistical Analysis

One-way ANOVA with three replicates was performed to study the significant differences among treatments. Statistical analysis was conducted by means of the SPSS v. 21, (SPSS Inc., Chicago, IL, USA) statistical package.

3. Results and Discussion

3.1. Identification of the Diuron Degrading Bacterial Strain

A soil that had been treated with a great variety of pesticides for a long time, including phenylurea herbicides, was enriched with diuron in order to isolate a diuron-degrading mixed consortium. 11 bacterial strains (LL1D, LL2D, LL3D, LL4D, LL5D, LL6D, LL7D, LL8D, LL9D, LL10D and LL11D) were isolated from the consortium initially obtained and only the strain LL6D showed ability to remove diuron in solution, achieving an extent of biodegradation 97.4% after 20 d of the assay (Figure S1). LL6D was identified and selected to perform biodegradation assays due to its capacity to use diuron as the only source of carbon and energy. 16S rRNA gene sequence showed a match of 100% to species from *O. anthropi* in the NCBI GenBank, and it was named *O. anthropi* CD3. This strain belongs to the phylum Proteobacteria. Other bacteria belonging to this phylum are in relation with phenylurea herbicides biodegradation. Pesce et al. [41] showed that Proteobacterium phylum was identified as the bacterial phylotype after being in contact with a realistic diuron exposure. *Sphingomonas* sp. strain SRS2, which belongs to the Proteobacterium phylum, combined with SRS1 strain

isolated from a soil contaminated by isoproturon, were able to mineralize the herbicide diuron. SRS2 was described as a possible new genus whose 16S rRNA gene sequence present 95.7% of similarity with *Polaromonas vacuolata* [42]. However, it is important to highlight that *O. anthropi* has never been described as a phenylurea degrader, although several studies have described strains belonging to *Ochrobactrum* species as degrader for other pesticides. *Ochrobactrum* sp. JAS2 was used to degrade chlorpyrifos, an organophosphate insecticide, in an aqueous medium [31]. *O. anthropi* strain NC-1, isolated from a pesticide-contaminated soil, was identified as a degrading strain (98.5% within 7 d) for phenmedipham [43]. *O. thiopenivorans* and *Sphingomonas melonis* were inoculated forming a consortium to remove the insecticide methomyl, achieving a bioremediation rate of 86% [44].

3.2. Phenylureas Biodegradation in Solution by *O. anthropi* CD3

O. anthropi CD3 isolated from an agricultural soil after enrichment with diuron, has been tested for biodegradation of five herbicides belonging to the phenylurea family. *O. anthropi* CD3 biodegraded, to different extents, all the studied herbicides (Figure 1). All the obtained biodegradation curves were fitted to a simple first-order kinetic model (SFO) (Table 2). In the case of isoproturon, a significant percentage of biotransformation was observed reaching a degradation of about 89% after the 60 d of the assay (Table 2). In the cases of diuron, linuron, chlorotoluron and fluometuron, their initial concentration was completely removed when *O. anthropi* CD3 was inoculated. Moreover, diuron, linuron and chlorotoluron were rapidly transformed, achieving 50% of biodegradation after only 1.1, 3.8 and 0.4 d, respectively (Table 2).




Figure 1. Biodegradation curves (60 d) of (♦) diuron, (▲) linuron, (+) isoproturon, (■) chlorotoluron and (●) fluometuron in solution in presence of *Ochrobactrum anthropi* CD3. Solid lines show model fitting to the experimental results (symbols). Standard deviation (vertical bars).

Table 2. Kinetic parameters obtained from phenylurea herbicides biodegradation in solution after inoculation with *Ochrobactrum anthropi* CD3.

Contaminant	R ²	χ ²	K (d ⁻¹)	DT ₅₀ (d)	Extent of Biodegradation (%)
Diuron	0.89	7.2	6.2 × 10 ⁻¹	1.1	100
Linuron	0.91	10.8	2.5 × 10 ⁻¹	3.8	100
Isoproturon	0.94	14.6	3.5 × 10 ⁻²	19.5	89
Chlorotoluron	0.99	13.5	1.8	0.4	100
Fluometuron	0.98	14.3	8.4 × 10 ⁻²	8.3	100

Biodegradation curve were fitted to single first order model. K: mineralization rate constants. DT₅₀: Time required for the concentration to decline to half of the initial value.

The relationship between pesticide structure and biodegradation is known. Alexander [45] showed that there is the specificity of the microorganism for a source of carbon is mainly related to the enzymatic function, which is largely restricted to performing only a single type of reaction on a range of substrates of very similar structures. In this work, *O. anthropi* CD3 was able of degrading five herbicides that belong to the same family and, therefore, with similar chemical structures.

The ability of individual strains to degrade different phenylurea herbicides have been previously demonstrated. Tixier et al. [17] carried out biodegradation experiments in solution employing the bacterial phenylureadegrader *Arthrobacter* sp. N2, isolated from a diuron-contaminated soil. The ability of *Arthrobacter* sp. N2 to degrade diuron, chlorotoluron and isoproturon in solution depended on the herbicide considered. Although the three phenylurea herbicides were completely degraded, the fastest biodegradation occurred with diuron (10 h), followed by chlorotoluron (30 h) and finally isoproturon (5 d). In the case of this last herbicide, the transformation rate seemed to be affected by the presence of the isopropyl group. Cullington and Walker [46] also studied this phenomenon during biodegradation assays of five different herbicides in solution using an unidentified soil bacterium, observing different efficacy in the rate of degradation: linuron > diuron > monolinuron >> metoxuron >>> isoproturon. Diuron and linuron were easily converted whereas the bacterium showed much lower activity towards isoproturon. Something similar happened in the present study, where isoproturon was the only phenylurea herbicide that was not completely degraded after 60 d of incubation, which pointed out that the presence of the isopropyl group influenced the transformation rate. The isopropyl group makes it difficult for microorganisms to use isoproturon as a carbon source, because it presents steric hindrance and hence microorganisms will prefer other easier to attack carbon sources.

Sharma et al. [19] reported that *Micrococcus* sp. PS-1 isolated from a diuron storage site could biodegrade diuron and other structural analogue phenylurea herbicides in the order: fenuron (99%) > monuron (98%) > diuron (96%) > monolinuron (80%) > linuron (74%) > chlorotoluron (65%). These results suggested that the size, degree of ring chlorination and the nature of the urea N'-substituents were the determining factors in the occurrence of degradation of phenylurea herbicides. Similarly, in our study, different biodegradation rates were observed, probably due to their structural differences.

The structure of phenylurea herbicides is not the only factor influencing their biodegradability as mentioned above, but also the type of bacterial strain plays an important role. The Gram-negative strain *Diaphorobacter* sp. LR2014-1 was capable of biodegrading five phenylurea herbicides with different structures [16]. The authors concluded that this large substrate spectrum points to the fact that strain LR2014-1 harbours an unusual enzyme or multiple enzymes to catabolize these phenylureas. It should be pointed out that those herbicides which presented the highest degradation (linuron and chlorotoluron) were those which gave the lowest degradation in the study of Sharma et al. [19], indicating the important role of the catabolizing enzymes present in the bacterial strain used.

As with the biodegradation observed with the bacterial strains reported previously, the results obtained indicate that the isolated bacterial strain *O. anthropi* CD3 has an exceptionally broad substrate spectrum for a variety of substituted phenylurea herbicides.

In this work, 3,4-DCA was the only metabolite quantifiable in the solution formed from diuron and linuron biodegradation after inoculation with the investigated phenylurea degrading strain. For chlorotoluron, fluometuron and isoproturon typical aniline metabolites (3-chloro-4-methylaniline, *m*-trifluoromethylaniline, isopropylaniline, respectively) were not detectable at the end of the biodegradation assay. Also, Tixier et al. [17] observed that the aniline metabolites from chlorotoluron and isoproturon were produced at a low concentration that they could not be also detected, whereas formation of 3,4-DCA from diuron was almost quantitative. 3,4-DCA can be produced during biotic and abiotic degradation processes of diuron and linuron, and shows a higher ecotoxicological impact on living organisms than the parent compounds and hence the importance of studying its possible formation. Huda Bhuiyan et al. [47] stated that 3,4-DCA can be considered as

one of the major aniline derivatives released into the environment, capable of provoking disruption of the endocrine and other biological systems in animals. Figure 2 shows the concentration of 3,4-DCA produced throughout the biodegradation of diuron and linuron in solution when *O. anthropi* CD3 was inoculated. The highest 3,4-DCA concentration was observed within 9 and 15 d of incubation in the case of diuron (4.7–4.9 mg L⁻¹), and in the case of linuron, the highest concentration (5.9 mg L⁻¹) was reached after 7 d of assay. 3,4-DCA concentration was maintained without any further degradation until the end of the experiment (60 d). In both cases a similar metabolite formation profile was observed.




Figure 2. Concentration of 3,4-dichloroaniline (3,4-DCA) in solution throughout the biodegradation process of diuron (◆) and linuron (*) in presence of *Ochrobactrum anthropi* CD3. Standard deviation (vertical bars).

3.3. Diuron and 3,4-Dichloroaniline Biodegradation in Soils

O. anthropi CD3 has been also selected for performing diuron biodegradation experiments in soils. As commented above, Blondel et al. and Agbaogun, et al. [7,8] observed that the K_{ow} values of phenylurea herbicides were as high as the number of chlorine atoms on the phenyl group, provoking an increase in their adsorption to soil particles. This property combined with the high level of persistence of the herbicide diuron makes it a hazard for soil environment.

Different bioremediation techniques have been proven for application in two different soils (PLD and R), which were artificially contaminated with diuron: (i) biostimulation, using a nutrient solution (NS) which accelerates the decontamination rate; (ii) use of HPBCD solution as bioavailability enhancer; (iii) bioaugmentation, using *O. anthropi* CD3. Figure 3 shows the different biodegradation curves obtained from these assays.

Diuron abiotic dissipation was evaluated in both soils, where a $HgCl_2$ solution was added to soil prior to begin of the assays with the aim of killing the soil endogenous microbiota (Figure 3). The global extent of biodegradation for the abiotic control was only 1.1% and 1.2% for R and PLD, respectively (Table 3). The role of soil microbiota in diuron biodegradation was evaluated through the addition of nutrients (NS), but no significant biodegradation was observed in any of the investigated soils (1.3% after 100 d), with DT_{50} values of 24.6 and 29 years for soils PLD and R, respectively, indicating the extremely high persistence of diuron in these soils. This observed result could be due to the fact that the soil endogenous microbiota is not adapted for diuron degradation; but another reason can be possible: the low bioavailability of the herbicide due to its strong adsorption on soil particles. These results point the need to use techniques of bioaugmentation with specific microbial degraders and/or the use of CDs as bioavailability enhancers.




Figure 3. Diuron biodegradation curves in R (A) and PLD (B) soils after the application of: NS (*), HPBCD + NS (+), *Ochrobactrum anthropi* CD3 + NS (◆), *Ochrobactrum anthropi* CD3 + HPBCD + NS (▲) and HgCl₂ (-). Standard deviation (vertical bars).

This latter option was tested by using HPBCD. In the case of R soil, diuron biodegradation curve obtained in the presence of NS and HPBCD indicated a slight enhancement in comparison with the biodegradation profile obtained when only biostimulation with NS was applied (Figure 3A). DT₅₀ decreased from more than 10,000 d until about 1000 d, and 20.3% of diuron biodegraded was reached after 100 d of experiment (Table 3). In the case of PLD soil, better results were observed, where the addition of HPBCD provoked a great increase in diuron biodegradation, decreasing DT₅₀ from almost 9000 d to about 37 d, and reaching an extent of biodegradation of 84%. These results might be due to the low OM content in PLD soil (1.67%) in comparison with the R soil (3.44%). Diuron has a high affinity for soil OM, causing a reduction of its bioavailability as higher is the percentage of OM in soil [48–50].

Table 3. Kinetic parameters obtained after different biodegradation treatments applied in R and PLD soils contaminated with diuron.

Soil	Treatment	Kinetic Model	R ²	χ ²	K ₁ (d ⁻¹)	K ₂ (d ⁻¹)	tb (d)	DT ₅₀ (d)	Extent of Biodegradation (%)
R	NS	SFO	0.99	4.2	5.2 × 10 ⁻⁴	-	-	10,569	1.3
	HPBCD + NS	HS	0.91	3.8	8.4 × 10 ⁻³	4.4 × 10 ⁻⁴	26.9	1077	20.3
	<i>O. anthropi</i> CD3 + NS	SFO	0.99	2.3	1.1 × 10 ⁻¹	-	-	6.1	100
	<i>O. anthropi</i> CD3 + HPBCD + NS	SFO	0.94	14.1	1.5 × 10 ⁻¹	-	-	4.7	100
	HgCl ₂	SFO	0.98	5.1	5.1 × 10 ⁻⁴	-	-	17,011	1.1
PLD	NS	SFO	0.99	7.9	4.2 × 10 ⁻⁴	-	-	8977	1.3
	HPBCD + NS	SFO	0.88	14.8	1.8 × 10 ⁻²	-	-	37.1	84.0
	<i>O. anthropi</i> CD3 + NS	SFO	0.9	14.3	2.2 × 10 ⁻¹	-	-	3.1	100
	<i>O. anthropi</i> CD3 + HPBCD + NS	SFO	0.92	14.8	1.1	-	-	0.7	100
	HgCl ₂	SFO	0.98	1.7	1.9 × 10 ⁻⁴	-	-	3501	1.2

NS: nutrients solution. K: mineralization rate constants. tb: Time at which rate constant changes. DT₅₀: Time required for the concentration to decline to half of the initial value.

HPBCD is widely known as an ecofriendly compound for organic contaminants bioremediation due to its ability to form inclusion complexes between HPBCD hydrophobic cavity and hydrophobic molecules, which improves their water solubility, making them more available for microorganisms [51–53]. In this case, the increase in soil availability of diuron mediated by HPBCD will be facilitated as lower is the OM content of the soil.

Reid et al. [54] conducted a non-exhaustive extraction technique employing HPBCD solutions for PAHs, and observed a significant correlation with mineralization of a phenanthrene-degrading bacteria in three different soils to assess its bioavailability. This method has been broadly used for assessing the availability of different types of organic contaminants in soil [23,55].

Soil diuron extraction experiments were performed, in order to test the effect of soil solution, NS and HPBCD solutions on diuron availability on the two investigated soils (Figure 4). The percentage of diuron extracted was lower when NS solution was used as extractant. When HPBCD was employed, in the case of the R soil, the amount of diuron extracted was similar to those extracted with the NS solution 17.4% and 22.3%, respectively, and in the case of PLD soil with a lower soil OM content, diuron was extracted from 25.4%, when NS was added, to 58% when HPBCD was used. The content of OM in the soil plays an important role in diuron bioavailability [27], which explains the significant decrease observed in DT₅₀ for diuron biodegradation in the case of PLD soil (Table 3).

The higher diuron degradation observed can also be caused by the stimulating effect of HPBCD on the soil microbial community, since it is an oligosaccharide, and may be considered an attractive carbon source for microorganisms [56]. The high percentage of degradation obtained in PLD soil after the addition of HPBCD indicates that the soil microbiota would act as an effective diuron degrader after being stimulated in the presence of HPBCD, because such microbiota had been previously adapted to the presence of diuron or other compounds with similar chemical structures, which would have produced an alteration of soil microbial communities, favoring the development of some microbial taxa which are able to use diuron as a source of carbon and energy [57].




Figure 4. Diuron extracted from R (A) and PLD (B) soil in presence of H₂O, nutrient solution (NS) and HPBCD. Standard deviation (vertical bars). The same letter in the same column means no significant differences among treatments.

The effect of bioaugmentation with *O. anthropi* CD3 on diuron biodegradation in both investigated soils can be also observed in Figure 3 and Table 3. In the case of R soil, the initial concentration of diuron was biodegraded completely after 50 d in the presence of *O. anthropi* CD3 + NS, and DT₅₀ was 6.1 d. When *O. anthropi* CD3 and HPBCD were jointly applied, 100% of diuron was removed after 35 d of the assay, with a slight decrease in the DT₅₀ (4.7 d). In the case of PLD soil, diuron was completely biodegraded when *O. anthropi* CD3 + NS was applied after 20 d of the assay (Figure 3), and the DT₅₀ observed was only 3.1 d. A particularly noteworthy improvement in biodegradation kinetics was observed when the bacterial strain was added together with HPBCD, decreasing DT₅₀ to only 0.7 d.

The main metabolite of diuron, 3,4-DCA, was monitored throughout the soil biodegradation assays when using *O. anthropi* in both soils (Figure 5 and Figure S2). Its maximum concentration in PLD soil was observed after 24 h of the assay (Figure 5B), reaching a higher value in the presence of HPBCD, and being completely degraded after 50 d. R soil has a higher soil OM content than PLD soil, which provoked a lower bioavailability of both, the herbicide and 3,4-DCA [58], and for this reason, the maximum concentration of 3,4-DCA was reached after 8 d, and its dissipation rate in this soil was much slower than in PLD soil, being not possible to achieve its complete biodegradation after 100 d of incubation (Figure 5).




Figure 5. Concentration of 3,4-dichloroaniline (3,4-DCA) in soils R (A) and PLD (B) throughout the biodegradation process of diuron in presence of *Ochrobactrum anthropi* CD3 + NS (▲) and *Ochrobactrum anthropi* CD3 + HPBCD + NS (■). Standard deviation (vertical bars).

O. anthropi CD3 has proved to be capable of removing completely the initial concentration of diuron in both studied soils. However, this bacterial strain was not capable of carrying out the degradation of 3,4-DCA in solution (Figure 2). The synergetic action of *O. anthropi* CD3 and the soil endogenous microbiota was able to reduce the metabolite concentration formed during diuron biodegradation assays. The microbial community of soil PLD showed to be more effective to degrade diuron and its metabolite than that of R soil, due to the contact of PLD soil with compounds with chemical structures similar to diuron for a longer time, as commented above.

The significance of bacterial consortia composed of several strains with synergistic catabolic co-operation for the degradation of multiple pesticides in solution has been previously mentioned [59].

3.4. Ecotoxicology

Soil toxicity was evaluated in the studied artificially contaminated soils (50 mg kg⁻¹ of diuron) before and after treatment with *O. anthropi* CD3 and/or HPBCD solution using a Microtox[®] test. The measured toxicity parameters are shown in Table 4. In the case of the R soil, after inoculation and inoculation + HPBCD only a slight decrease in toxicity at the

end of the experiment (100 d) was determined. TU value before inoculation was 4.0, and this toxicity was reduced to 3.2 and 3.3, respectively, after both treatments were applied. Even though diuron was completely biodegraded, a level of acute toxicity was maintained as a result of the presence of the formed 3,4-DCA remaining in the soil after being treated (Figure 5A). In R soil, its endogenous microbiota was not able to synergistically act with the investigated strain *O. anthropi* CD3 to completely remove the metabolite. On the contrary, in the case of PLD soil, significant differences could be observed in the values studied as indicators of toxicity. When *O. anthropi* CD3 was inoculated, TU decreased from 10.9 (high acute toxicity, $10 < TU < 100$) to 1.4 (acute toxicity), a value very close to slight acute toxicity ($0.4 < TU < 1$). Similar results were observed when HPBCD and *O. anthropi* CD3 were employed together for soil decontamination. This strong toxicity decrease is due to the fact that 3,4-DCA was completely removed from PLD soil sample (Figure 5B), but, in spite of this, the soil remained toxic, indicating that other toxic degradation compounds must be present, and, therefore, the soil sample would need a deeper bioremediation to decrease its toxicity to levels similar to the uncontaminated soil. Monitoring pollutant levels by analytical methods shows the degradation of the contaminants, but it does not reveal the toxicity of the potentially formed metabolites. For this reason, it is important to assess the soil ecotoxicology after any bioremediation technique application [60,61] to determine whether or not there is toxicity in the sample, through their comparison with the classification proposed by Persoone et al. [40].

Table 4. Acute toxicity test towards *V. fischeri* before and after 100 d of incubation of R and PLD soils contaminated with diuron and treated with *O. anthropi* CD3 and HPBCD.

Soil	Treatment	EC ₅₀ (%) *	TU	Class	Toxicity **
R	Uncontaminated	166	0.6	Class II	Slight acute toxicity
	Contaminated with diuron	24.7	4.0	Class III	Acute toxicity
	<i>O. anthropi</i> CD3	30.8	3.2	Class III	Acute toxicity
	<i>O. anthropi</i> CD3 + HPBCD	29.9	3.3	Class III	Acute toxicity
PLD	Uncontaminated	111	0.9	Class II	Slight acute toxicity
	Contaminated with diuron	9.2	10.9	Class IV	High Acute toxicity
	<i>O. anthropi</i> CD3	73.2	1.4	Class III	Acute toxicity
	<i>O. anthropi</i> CD3 + HPBCD	72.8	1.4	Class III	Acute toxicity

* For each soil sample the EC₅₀ value corresponds to the soil extract concentration (% v/v) having a toxic effect on 50% of *V. fischeri*. ** According to Persoone et al. (2003).

4. Conclusions

The Gram-negative bacterial strain *O. anthropi* CD3 isolated from an agricultural soil by selective enrichment with diuron, and which was not previously described as phenylurea degrader, was capable of utilizing some phenylurea herbicides as sole carbon sources. The relative degradation profile in aqueous solution of *O. anthropi* CD3 was in the order of chlorotoluron > diuron > linuron > fluomethuron > isoproturon, which was related to the different structural features of the molecules. The only metabolite detected throughout the biodegradation processes was 3,4-DCA from diuron and linuron, which could not be degraded by *O. anthropi* CD3 in solution.

The use of *O. anthropi* CD3 in soils contaminated by diuron led to its degradation, but at different levels depending on their soil organic matter content (the higher the OM the lower the biodegradation) and the characteristics of their endogenous microbiota, reaching the complete degradation of diuron and 3,4-DCA in that soil with a microbial population more adapted to degrade similar contaminants and able to synergistically act with *O. anthropi* CD3 to remove the metabolite. HPBCD was used as a bioavailability enhancer of diuron due to its capacity to form an inclusion compound with the herbicide, and also as biostimulant of the endogenous microbial community, increasing considerably the rate of diuron biodegradation in the soil with the least amount of OM.

Ecotoxicity studies showed a decrease in toxicity of the studied contaminated soils, but more drastically in the soil where both diuron and 3,4-DCA were completely removed. However, the complete elimination of toxicity could not be reached, indicating the presence of other toxic compounds. It indicates the need to carry out ecotoxicity studies to assess the feasibility of any bioremediation treatment application, even when the parent contaminant and its main metabolites had been reduced or completely eliminated.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/ijerph19031365/s1>. Figure S1: quantification of the diuron degraded in aqueous solution after 20 days by the bacterial strains isolated from an agricultural soil, Figure S2: HPLC chromatograms obtained from the extracts of PLD soil (A.1) and R soil (B.1) after one day of incubation; and from PLD soil (A.2) and R soil (B.2) after eight days of incubation, Table S1: Formulas, molecular structures and some properties of phenylurea herbicides studied in this work (PPDB. IUPAC Pesticides Properties DataBase).

Author Contributions: Conceptualization, L.-M.A. and M.E. and V.J.; methodology, L.-M.A. and V.J.; software, L.-M.A. and M.E. and V.J.; validation, L.-M.A. and M.E. and V.J.; formal analysis, L.-M.A.; investigation, L.-M.A. and V.J.; resources, M.E.; data curation, L.-M.A. and M.E. and V.J.; writing—original draft preparation, L.-M.A.; supervision, M.E. and V.J.; funding acquisition, M.E. and V.J. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Acknowledgments: This work was supported by the Spanish Ministry of Economy and Competitiveness under the research project CMT2017-82472-C2-1-R (AEI/FEDER, UE). Alba, Lara-Moreno acknowledges the Spanish Ministry of Education, Culture and Sports for her FPU fellowship (FPU15/03740).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Castelo-Grande, T.; Augusto, P.; Monteiro, P.; Estevez, A.M.; Barbosa, D. Remediation of soils contaminated with pesticides: A review. *Int. J. Environ. Anal. Chem.* **2010**, *90*, 438–467. [[CrossRef](#)]
2. Abiotic and Biological Technologies for the Remediation of Phenylurea Herbicides in Soils. Available online: <https://link.springer.com/chapter/10.1007/978-94-007-799-9> (accessed on 21 August 2021).
3. Greulich, K.; Hoque, E.; Pflugmacher, S. Uptake, metabolism, and effects on detoxication enzymes of isoproturon in spawn and tadpoles of amphibians. *Ecotoxicol. Environ. Saf.* **2002**, *52*, 256–266. [[CrossRef](#)] [[PubMed](#)]
4. Nélieu, S.; Bonnemoy, F.; Bonnet, J.L.; Lefevre, L.; Baudiffier, D.; Heydorff, M.; Quéméneur, A.; Azam, D.; Ducrot, P.H.; Lagadic, L.; et al. Ecotoxicological effects of diuron and chlorotoluron nitrate-induced photodegradation products: Monospecific and aquatic mesocosm-integrated studies. *Environ. Toxicol. Chem.* **2010**, *29*, 2644–2652. [[CrossRef](#)] [[PubMed](#)]
5. ElGouzi, S.; Mingorance, M.; Draoui, K.; Chtoun, E.; Peña, A. Assessment of phenylurea herbicides sorption on various Mediterranean soils affected by irrigation with wastewater. *Chemosphere* **2012**, *89*, 334–339. [[CrossRef](#)] [[PubMed](#)]
6. Burant, A.; Selbig, W.; Furlong, E.T.; Higgins, C.P. Trace organic contaminants in urban runoff: Associations with urban land-use. *Environ. Pollut.* **2018**, *242*, 2068–2077. [[CrossRef](#)]
7. Blondel, A.; Langeron, J.; Sayen, S.; Hénon, E.; Couderchet, M.; Guillon, E. Molecular properties affecting adsorption coefficient of phenylurea herbicides. *Environ. Sci. Pollut. Res.* **2013**, *20*, 6266–6281. [[CrossRef](#)]
8. Agbaogun, B.K.; Fischer, K. Adsorption of phenylurea herbicides by tropical soils. *Environ. Monit. Assess.* **2020**, *192*, 212–220. [[CrossRef](#)]
9. IUPAC FOOTPRINT Pesticide Properties Data-Base. Available online: <https://sitem.herts.ac.uk/aeru/iupac/index.htm> (accessed on 1 June 2020).
10. Boscolo, C.N.P.; Pereira, T.S.B.; Batalhão, I.G.; Dourado, P.L.R.; Schlenk, D.; de Almeida, E.A. Diuron metabolites act as endocrine disruptors and alter aggressive behavior in Nile tilapia (*Oreochromis niloticus*). *Chemosphere* **2018**, *191*, 832–838. [[CrossRef](#)]
11. European Commission. Water Framework Directive, Priority Substances and Certain Other Pollutants according to Annex II of Directive 2013/39/EU. List of Priority Substances in the Field of Water Policy. Available online: https://ec.europa.eu/environment/water/water-dangersub/pri_substances.htm (accessed on 17 August 2021).

12. Hussain, S.; Arshad, M.; Springael, D.; Sørensen, S.R.; Bending, G.; Devers-Lamrani, M.; Maqbool, Z.; Martin-Laurent, F. Abiotic and Biotic Processes Governing the Fate of Phenylurea Herbicides in Soils: A Review. *Crit. Rev. Environ. Sci. Technol.* **2015**, *45*, 1947–1998. [[CrossRef](#)]
13. Cycoń, M.; Mroziak, A.; Piotrowska-Seget, Z. Bioaugmentation as a strategy for the remediation of pesticide-polluted soil: A review. *Chemosphere* **2017**, *172*, 52–71. [[CrossRef](#)]
14. Morillo, E.; Villaverde, J. Review Advanced technologies for the remediation of pesticide-contaminated soils. *Sci. Total Environ.* **2017**, *586*, 576–597. [[CrossRef](#)] [[PubMed](#)]
15. Ellegaard-Jensen, L.; Knudsen, B.E.; Johansen, A.; Albers, C.; Aamand, J.; Rosendahl, S. Fungal–bacterial consortia increase diuron degradation in water-unsaturated systems. *Sci. Total Environ.* **2014**, *466–467*, 699–705. [[CrossRef](#)] [[PubMed](#)]
16. Zhang, L.; Hang, P.; Hu, Q.; Chen, X.L.; Zhou, X.Y.; Chen, K.; Jiang, J.D. Degradation of phenylurea herbicides by a novel bacterial consortium containing synergistically catabolic species and functionally complementary hydrolases. *J. Agric. Food Chem.* **2018**, *66*, 12479–12489. [[CrossRef](#)] [[PubMed](#)]
17. Tixier, C.; Sancelme, M.; Aït-Aïssa, S.; Widehem, P.; Bonnemoy, F.E.; Cuer, A.; Truffaut, N.; Veschambre, H. Biotransformation of phenylurea herbicides by a soil bacterial strain, *Arthrobacter* sp. N2: Structure, ecotoxicity and fate of diuron metabolite with soil fungi. *Chemosphere* **2002**, *46*, 519–526. [[CrossRef](#)]
18. Sørensen, S.R.; Albers, C.; Aamand, J. Rapid Mineralization of the Phenylurea Herbicide Diuron by *Variovorax* sp. Strain SRS16 in Pure Culture and within a Two-Member Consortium. *Appl. Environ. Microbiol.* **2008**, *74*, 2332–2340. [[CrossRef](#)]
19. Sharma, P.; Chopra, A.; Cameotra, S.S.; Suri, C.R. Efficient biotransformation of herbicide diuron by bacterial strain *Micrococcus* sp. PS-1. *Biodegradation* **2010**, *21*, 979–987. [[CrossRef](#)]
20. Muendo, B.M.; Shikuku, V.O.; Lalah, J.O.; Getenga, Z.M.; Wandiga, S.O.; Rothballer, M. Enhanced degradation of diuron by two *Bacillus* species isolated from diuron contaminated sugarcane and pineapple-cultivated soils in Kenya. *Appl. Soil Ecol.* **2020**, *157*, 103721. [[CrossRef](#)]
21. Silambarasan, S.; Logeswari, P.; Ruiz, A.; Cornejo, P.; Kannan, V.R. Influence of plant beneficial *Stenotrophomonas rhizophila* strain CASB3 on the degradation of diuron-contaminated saline soil and improvement of *Lactuca sativa* growth. *Environ. Sci. Pollut. Res.* **2020**, *27*, 35195–35207. [[CrossRef](#)]
22. Wang, Y.; Li, H.; Feng, G.; Du, L.; Zeng, D. Biodegradation of diuron by an endophytic fungus *Neurospora intermedia* DP8-1 isolated from sugarcane and its potential for remediating diuron-contaminated soils. *PLoS ONE* **2017**, *12*, e0182556. [[CrossRef](#)]
23. Villaverde, J.; Rubio-Bellido, M.; Posada-Baquero, R.; Madrid, F.; Morillo, E. Hydroxypropyl- β -cyclodextrin-based extraction for diuron bioaccessibility in an artificially contaminated soil. *Int. J. Environ. Anal. Chem.* **2013**, *93*, 1620–1627. [[CrossRef](#)]
24. Villaverde, J.; Rubio-Bellido, M.; Lara-Moreno, A.; Merchan, F.; Morillo, E. Combined use of microbial consortia isolated from different agricultural soils and cyclodextrin as a bioremediation technique for herbicide contaminated soils. *Chemosphere* **2018**, *193*, 118–125. [[CrossRef](#)] [[PubMed](#)]
25. Villaverde, J.; Posada-Baquero, R.; Rubio-Bellido, M.; Laiz, L.; Saiz-Jimenez, C.; Sanchez-Trujillo, M.A.; Morillo, E. Enhanced Mineralization of Diuron Using a Cyclodextrin-Based Bioremediation Technology. *J. Agric. Food Chem.* **2012**, *60*, 9941–9947. [[CrossRef](#)] [[PubMed](#)]
26. Sørensen, S.R.; Aamand, J. Rapid mineralisation of the herbicide isoproturon in soil from a previously treated Danish agricultural field. *Pest. Manag. Sci.* **2003**, *59*, 1118–1124. [[CrossRef](#)] [[PubMed](#)]
27. Villaverde, J.; Maqueda, C.; Morillo, E. Improvement of the desorption of the herbicide norflurazon from soils via complexation with β -cyclodextrin. *J. Agric. Food. Chem.* **2006**, *53*, 5366–5372. [[CrossRef](#)] [[PubMed](#)]
28. Yáñez, C.; Cañete-Rosales, P.; Castillo, J.P.; Catalán, N.; Undabeytia, T.; Morillo, E. Cyclodextrin inclusion complex to improve physicochemical properties of herbicide bentazon: Exploring better formulations. *PLoS ONE* **2012**, *7*, e41072. [[CrossRef](#)] [[PubMed](#)]
29. Morillo, E.; Madrid, F.; Lara-Moreno, A.; Villaverde, J. Soil bioremediation by cyclodextrins. A review. *Int. J. Pharm.* **2020**, *591*, 119943. [[CrossRef](#)]
30. Sánchez-Trujillo, M.A.; Lacorte, S.; Villaverde, J.; Barata, C.; Morillo, E. Decontamination of polycyclic aromatic hydrocarbons and nonylphenol from sewage sludge using hydroxypropyl- β -cyclodextrin and evaluation of the toxicity of leachates. *Environ. Sci. Pollut. Res.* **2014**, *21*, 507–517. [[CrossRef](#)]
31. Abraham, J.; Silambarasan, S. Biodegradation of chlorpyrifos and its hydrolysis product 3,5,6-trichloro-2-pyridinol using a novel bacterium *Ochrobactrum* sp. JAS2: A proposal of its metabolic pathway. *Pestic. Biochem. Physiol.* **2016**, *126*, 13–21. [[CrossRef](#)]
32. Egea, T.C.; Da Silva, R.; Boscolo, M.; Rignonato, J.; Monteiro, D.A.; Grünig, D.; da Silva, H.; der Wielen, F.; Helmus, R.; Parsons, J.R.; et al. Diuron degradation by bacteria from soil of sugarcane crops. *Heliyon* **2017**, *3*, e00471. [[CrossRef](#)]
33. Soil Maps. National Geographic Institute. Nature Database (Spanish Ministry of Environment). 2005. Available online: <https://www.ign.es/web/catalogo-cartoteca/recursos/html/030769.html> (accessed on 20 November 2021).
34. Fenlon, K.A.; Andreou, K.; Jones, K.C.; Semple, K.T. The extractability and mineralisation of cypermethrin aged in four UK soils. *Chemosphere* **2011**, *82*, 187–192. [[CrossRef](#)]
35. Lane, D.J. 16S/23S rRNA Sequencing. In *Nucleic Acid Techniques in Bacterial Systematics*; Stackebrandt, E., Goodfellow, M., Eds.; Wiley: Chichester, UK, 1991; Volume 2, pp. 115–175.
36. FOCUS. Guidance Document on Estimating Persistence and Degradation Kinetics from Environmental Fate Studies on Pesticides in EU Registration. Report of the FOCUS Work Group on Degradation Kinetics. EC Documents Reference

- Sanco/10058/2005 Version 2.0. 2006. Available online: https://esdac.jrc.ec.europa.eu/public_path/projects_data/focus/dk/docs/finalreportFOCDegKinetics.pdf (accessed on 17 August 2021).
37. Beulke, S.; Beinum, W.; Van Brown, C.D.; Mitchell, M.; Walker, A. Evaluation of Simplifying Assumptions on Pesticide Degradation in Soil. *J. Environ. Qual.* **2005**, *34*, 1933–1943. [[CrossRef](#)] [[PubMed](#)]
 38. UNE-EN ISO 11348-3:2009/A1:2019. Water Quality—Determination of the Inhibitory Effect of Water Samples on the Light Emission of *Vibrio Fischeri* (Luminescent Bacteria Test)—Part 3: Method Using Freeze-Dried Bacteria—Amendment 1 (ISO 11348-3:2007/Amd1:2018). Available online: [https://www.une.org/encuentra-tu-norma/busca-tu-norma/norma?c=N0062816#~:{}:text=2009%2FA1%3A2019-,Calidad%20del%20agua.,\(Ensayo%20de%20bacterias%20luminescentes](https://www.une.org/encuentra-tu-norma/busca-tu-norma/norma?c=N0062816#~:{}:text=2009%2FA1%3A2019-,Calidad%20del%20agua.,(Ensayo%20de%20bacterias%20luminescentes) (accessed on 17 August 2021).
 39. Molina, M.C.; González, N.; Bautista, L.F.; Sanz, R.; Simarro, R.; Sánchez, I.; Sanz, J.L. Isolation and genetic identification of PAH degrading bacteria from a microbial consortium. *Biodegradation* **2009**, *20*, 789–800. [[CrossRef](#)] [[PubMed](#)]
 40. Persoone, G.; Marsalek, B.; Blinova, I.; Törökne, A.; Zarina, D.; Manusadzianas, L.; Nalecz-Jawecki, G.; Tofan, L.; Stepanova, N.; Tothova, L.; et al. A practical and user-friendly toxicity classification system with microbiotests for natural waters and wastewaters. *Environ. Toxicol.* **2003**, *18*, 395–402. [[CrossRef](#)] [[PubMed](#)]
 41. Pesce, S.; Bardot, C.; Lehours, A.C.; Batisson, I.; Bohatier, J.; Fajon, C. Effects of diuron in microcosms on natural riverine bacterial community composition: New insight into phylogenetic approaches using PCR-TTGE analysis. *Aquat. Sci.* **2008**, *70*, 410–418. [[CrossRef](#)]
 42. Sørensen, S.R.; Ronen, Z.; Aamand, J. Growth in Coculture Stimulates Metabolism of the Phenylurea Herbicide Isoproturon by *Sphingomonas* sp. Strain SRS2. *Appl. Environ. Microbiol.* **2002**, *68*, 3478–3485. [[CrossRef](#)] [[PubMed](#)]
 43. Pujar, N.K.; Laad, S.; Premakshi, H.G.; Pattar, S.V.; Mirjankar, M.; Kamanavalli, C.M. Biodegradation of phenmedipham by novel *Ochrobactrum anthropi* NC-1. *3 Biotech* **2019**, *9*, 1–10. [[CrossRef](#)] [[PubMed](#)]
 44. Tatar, S.; Yildirim, N.C.; Serdar, O.; Erguven, G.O. Can toxicities induced by insecticide Methomyl be remediated via soil bacteria *Ochrobactrum thiophenivorans* and *Sphingomonas melonis*? *Curr. Microbiol.* **2020**, *77*, 1301–1307. [[CrossRef](#)]
 45. Alexander, M. *Biodegradation and Bioremediation*, 2nd ed.; 92101-4495; Academic Press: San Diego, CA, USA, 1999; ISBN 978-0-12-049861-1.
 46. Cullington, J.E.; Walker, A. Rapid biodegradation of diuron and other phenylurea herbicides by a soil bacterium. *Soil Biol. Biochem.* **1999**, *31*, 677–686. [[CrossRef](#)]
 47. Huda Bhuiyan, M.N.; Kang, H.; Kim, J.H.; Kim, S.; Kho, Y.; Choi, K. Endocrine disruption by several aniline derivatives and related mechanisms in a human adrenal H295R cell line and adult male zebrafish. *Ecotoxicol. Environ. Saf.* **2019**, *180*, 326–332. [[CrossRef](#)]
 48. Gholamalazadeh, A.; Smernik, R.J.; Chittleborough, D.J.; Kookana, R.S. Clear effects of soil organic matter chemistry, as determined by NMR spectroscopy, on the sorption of diuron. *Chemosphere* **2008**, *70*, 1153–1160.
 49. Albers, C.N.; Banta, G.T.; Erik, P.; Jacobsen, O.S. The influence of organic matter on sorption and fate of glyphosate in soil—Comparing different soils and humic substances. *Environ. Pollut.* **2009**, *157*, 2865–2870. [[CrossRef](#)] [[PubMed](#)]
 50. Echevarría, C.; Valderrama, C.; Cortina, J.L.; Martín, I.; Arnaldos, M.; Bernat, X.; De la Cal, A.; Boleda, M.R.; Vega, A.; Teuler, A.; et al. Techno-economic evaluation and comparison of PAC-MBR and ozonation-UV revamping for organic micro-pollutants removal from urban reclaimed wastewater. *Sci. Total Environ.* **2019**, *671*, 288–298. [[CrossRef](#)] [[PubMed](#)]
 51. Crini, C. Review: A History of Cyclodextrins. *Chem. Rev.* **2014**, *114*, 10940–10975. [[CrossRef](#)] [[PubMed](#)]
 52. Villaverde, J.; Rubio-Bellido, M.; Merchán, F.; Morillo, E. Bioremediation of diuron contaminated soils by a novel degrading microbial consortium. *J. Environ. Manag.* **2017**, *188*, 379–386. [[CrossRef](#)]
 53. Morillo, E.; Sánchez-Trujillo, M.A.; Moyano, J.R.; Villaverde, J.; Gómez-Pantoja, M.E.; Pérez-Martínez, J.I. Enhanced solubilisation of six PAHs by three synthetic cyclodextrins for remediation applications: Molecular modelling of the inclusion complexes. *PLoS ONE* **2012**, *7*, e44137.
 54. Reid, B.J.; Jones, K.C.; Semple, K.T. Bioavailability of persistent organic pollutants in soils and sediments a perspective on mechanisms, consequences and assessment. *Environ. Pollut.* **2000**, *108*, 103–112. [[CrossRef](#)]
 55. Rubio-Bellido, M.; Morillo, E.; Villaverde, J. Assessment of soil diuron bioavailability to plants and microorganisms through non-exhaustive chemical extractions of the herbicide. *Geoderma* **2018**, *312*, 130–138. [[CrossRef](#)]
 56. Fenyvesi, E.; Gruiz, K.; Verstichel, S.; De Wilde, B.; Leitgib, L.; Csabai, K.; Szaniszló, N. Biodegradation of cyclodextrins in soil. *Chemosphere* **2005**, *60*, 1001–1008. [[CrossRef](#)]
 57. Parajuli, A.; Grönroos, M.; Kauppi, S.; Płociniczak, T.; Roslund, M.I.; Galitskaya, P.; Laitinen, O.H.; Hyöty, H.; Jumpponen, A.; Strömmer, R.; et al. The abundance of health-associated bacteria is altered in PAH polluted soils—Implications for health in urban areas? *PLoS ONE* **2017**, *12*, 1–18. [[CrossRef](#)]
 58. Morillo, E.; Undabeytia, T.; Cabrera, A.; Villaverde, J.; Maqueda, C. Effect of soil type on adsorption-desorption, mobility, and activity of the herbicide norflurazon. *J. Agric. Food. Chem.* **2004**, *52*, 884–890. [[CrossRef](#)]
 59. Nzila, A. Update on the cometabolism of organic pollutants by bacteria. *Environ. Pollut.* **2013**, *178*, 474–482. [[CrossRef](#)] [[PubMed](#)]
 60. Jiang, Y.; Brassington, K.J.; Prpich, G.; Paton, G.I.; Semple, K.T.; Pollard, S.J.T.; Coulon, F. Insights into the biodegradation of weathered hydrocarbons in contaminated soils by bioaugmentation and nutrient stimulation. *Chemosphere* **2016**, *161*, 300–307. [[CrossRef](#)] [[PubMed](#)]
 61. Madrid, F.; Rubio-Bellido, M.; Villaverde, J.; Peña, A.; Morillo, E. Natural and assisted dissipation of polycyclic aromatic hydrocarbons in a long-term co-contaminated soil with creosote and potentially toxic elements. *Sci. Total Environ.* **2019**, *660*, 705–714. [[CrossRef](#)] [[PubMed](#)]