

Table S1. Association between FA symptoms and psychopathological scales (pt.1)

	Amount			Attempts			Time			Activities			Consequences			Tolerance		
	-	+	p	-	+	p	-	+	p	-	+	p	-	+	p	-	+	p
Internalising Problems (YSR)	40 (47.1)	9 (52.9)	0.658	33 (42.9)	16 (64.0)	0.066	25 (38.5)	24 (64.9)	0.010	26 (38.8)	23 (65.7)	0.010	29 (38.7)	20 (74.1)	0.002	34 (43.0)	15 (65.2)	0.061
Externalising Problems (YSR)	5 (5.9)	1 (5.9)	1.000	3 (3.9)	3 (12.0)	0.135	3 (4.6)	3 (8.1)	0.471	3 (4.5)	3 (8.6)	0.404	4 (5.3)	2 (7.4)	0.694	6 (7.6)	0 (0.0)	0.173
Total Problems (YSR)	27 (31.8)	5 (29.4)	0.849	22 (28.6)	10 (40.0)	0.285	15 (23.1)	17 (45.9)	0.017	18 (26.9)	14 (40.0)	0.175	22 (29.3)	10 (37.0)	0.459	24 (30.4)	8 (34.8)	0.689
Drive for Thinness (EDI-3)	44 (51.8)	14 (82.4)	0.020	41 (53.2)	17 (68.0)	0.196	33 (50.8)	25 (67.6)	0.100	30 (44.8)	28 (80.0)	0.001	38 (50.7)	20 (74.1)	0.035	40 (50.6)	18 (78.3)	0.019
Bulimia (EDI-3)	12 (14.1)	11 (64.7)	<0.001	10 (13.0)	13 (52.0)	<0.001	11 (16.9)	12 (32.4)	0.072	10 (14.9)	13 (37.1)	0.011	12 (16.0)	11 (40.7)	0.008	11 (13.9)	12 (52.2)	<0.001
Body dissatisfaction (EDI-3)	39 (45.9)	9 (52.9)	0.595	38 (49.4)	10 (40.0)	0.416	26 (40.0)	22 (59.5)	0.058	28 (41.8)	20 (57.1)	0.140	34 (45.3)	14 (51.9)	0.561	35 (44.3)	13 (56.5)	0.302
Low Self-esteem (EDI-3)	37 (43.5)	8 (47.1)	0.789	33 (42.9)	12 (48.0)	0.653	27 (41.5)	18 (48.6)	0.487	28 (41.8)	17 (48.6)	0.513	29 (38.7)	16 (59.3)	0.065	31 (39.2)	14 (60.9)	0.066
Personal Alienation (EDI-3)	37 (43.5)	8 (47.1)	0.789	32 (41.6)	13 (52.0)	0.361	27 (41.5)	18 (48.6)	0.487	27 (40.3)	18 (51.4)	0.282	30 (40.0)	15 (55.6)	0.163	32 (40.5)	13 (56.5)	0.173
Interpersonal Insecurity (EDI-3)	31 (36.5)	8 (47.1)	0.412	27 (35.1)	12 (48.0)	0.248	23 (35.4)	16 (43.2)	0.432	21 (31.3)	18 (51.4)	0.048	25 (33.3)	14 (51.9)	0.090	25 (31.6)	14 (60.9)	0.011
Interpersonal Alienation (EDI-3)	32 (37.6)	7 (41.2)	0.785	26 (33.8)	13 (52.0)	0.103	22 (33.8)	17 (45.9)	0.227	22 (32.8)	17 (48.6)	0.121	26 (34.7)	13 (48.1)	0.216	28 (35.4)	11 (47.8)	0.282
Interoceptive Deficits (EDI-3)	42 (49.4)	12 (70.6)	0.110	38 (49.4)	16 (64.0)	0.202	30 (46.2)	24 (64.9)	0.069	30 (44.8)	24 (68.6)	0.022	32 (42.7)	22 (81.5)	0.001	38 (48.1)	16 (69.6)	0.070
Emotional Dysregulation (EDI-3)	40 (47.1)	9 (52.9)	0.658	37 (48.1)	12 (48.0)	0.996	28 (43.1)	21 (56.8)	0.184	29 (43.3)	20 (57.1)	0.183	33 (44.0)	16 (59.3)	0.174	35 (44.3)	14 (60.9)	0.162
Perfectionism (EDI-3)	31 (36.5)	9 (52.9)	0.204	26 (33.8)	14 (56.0)	0.048	23 (35.4)	17 (45.9)	0.294	23 (34.3)	17 (48.6)	0.162	27 (36.0)	13 (48.1)	0.268	27 (34.2)	13 (56.5)	0.053
Asceticism (EDI-3)	36 (42.4)	10 (58.8)	0.213	28 (36.4)	18 (72.0)	0.002	25 (38.5)	21 (56.8)	0.074	25 (37.3)	21 (60.0)	0.029	31 (41.3)	15 (55.6)	0.203	31 (39.2)	15 (65.2)	0.028
Maturity Fears (EDI-3)	43 (50.6)	9 (52.9)	0.859	32 (41.6)	20 (80.0)	0.001	28 (43.1)	24 (64.9)	0.034	28 (41.8)	24 (68.6)	0.100	34 (45.3)	18 (66.7)	0.057	35 (44.3)	17 (74.0)	0.012
Total Problems (MASC 2)	11 (12.9)	2 (11.8)	0.894	9 (11.7)	4 (16.0)	0.574	5 (7.7)	8 (21.6)	0.043	6 (9.0)	7 (20.0)	0.112	6 (8.0)	7 (25.9)	0.017	8 (10.1)	5 (21.7)	0.142
Total score (CDI 2)	37 (43.5)	9 (52.9)	0.477	31 (40.3)	15 (60.0)	0.085	25 (38.5)	21 (56.8)	0.074	25 (37.3)	21 (60.0)	0.029	28 (37.3)	18 (66.7)	0.009	32 (40.5)	14 (60.9)	0.084

The Pearson's chi-square test was used to evaluate the association between the presence of FA's symptoms (+ present; - absent) and clinical scores at YSR, EDI-3, MASC 2 and CDI 2 scales. Statistical significance for p<0.05. CDI 2, Children Depression Inventory 2; EDI-3, Eating Disorder Inventory- 3; MASC 2, Multidimensional Anxiety Scale for Children 2; YSR, Youth Self Report.

Table S2. Association between FA symptoms and psychopathological scales (pt.2)

	Withdrawal			Problems			Obligations			Situations			Craving		
	-	+	p	-	+	p	-	+	p	-	+	p	-	+	p
Internalising Problems (YSR)	16 (32.0)	33 (63.5)	0.001	28 (39.4)	21 (67.7)	0.008	45 (47.9)	4 (50.0)	0.908	43 (47.3)	6 (54.5)	0.647	33 (42.9)	16 (64.0)	0.066
Externalising Problems (YSR)	1 (2.0)	5 (9.6)	0.102	4 (5.6)	2 (6.5)	0.872	5 (5.3)	1 (12.5)	0.407	6 (6.6)	0 (0.0)	0.380	3 (3.9)	3 (12.0)	0.135
Total Problems (YSR)	8 (16.0)	24 (46.2)	0.001	17 (23.9)	15 (48.4)	0.014	29 (30.9)	3 (37.5)	0.697	28 (30.8)	4 (36.4)	0.706	22 (28.6)	10 (40.0)	0.285
Drive for Thinness (EDI-3)	22 (44.0)	36 (69.2)	0.010	31 (43.7)	27 (87.1)	<0.001	51 (54.3)	7 (87.5)	0.068	51 (56.0)	7 (63.6)	0.631	41 (53.2)	17 (68.0)	0.196
Bulimia (EDI-3)	7 (14.0)	16 (30.8)	0.043	13 (18.3)	10 (32.3)	0.121	18 (19.1)	5 (62.5)	0.005	17 (18.7)	6 (54.5)	0.007	10 (13.0)	13 (52.0)	<0.001
Body dissatisfaction (EDI-3)	19 (38.0)	29 (55.8)	0.072	30 (42.3)	18 (58.1)	0.141	43 (45.7)	5 (62.5)	0.362	43 (47.3)	5 (45.5)	0.910	38 (49.4)	10 (40.0)	0.416
Low Self-esteem (EDI-3)	18 (36.0)	27 (51.9)	0.105	29 (40.8)	16 (51.6)	0.314	39 (41.5)	6 (75.0)	0.067	41 (45.1)	4 (36.4)	0.583	33 (42.9)	12 (48.0)	0.653
Personal Alienation (EDI-3)	17 (34.0)	28 (53.8)	0.044	28 (39.4)	17 (54.8)	0.150	42 (44.7)	3 (37.5)	0.695	39 (42.9)	6 (54.5)	0.461	32 (41.6)	13 (52.0)	0.361
Interpersonal Insecurity (EDI-3)	14 (28.0)	25 (48.1)	0.037	22 (31.0)	17 (54.8)	0.023	36 (38.3)	3 (37.5)	0.964	31 (34.1)	8 (72.7)	0.013	27 (35.1)	12 (48.0)	0.248
Interpersonal Alienation (EDI-3)	14 (28.0)	25 (48.1)	0.037	25 (35.2)	14 (45.2)	0.342	36 (38.3)	3 (37.5)	0.964	34 (37.4)	5 (45.5)	0.602	26 (33.8)	13 (52.0)	0.103
Interoceptive Deficits (EDI-3)	19 (38.0)	35 (67.3)	0.003	33 (46.5)	21 (67.7)	0.048	48 (51.1)	6 (75.0)	0.193	47 (51.6)	7 (63.6)	0.452	38 (49.4)	16 (64.0)	0.202
Emotional Dysregulation (EDI-3)	18 (36.0)	31 (59.6)	0.017	27 (38.0)	22 (71.0)	0.002	44 (46.8)	5 (62.5)	0.394	42 (46.2)	7 (63.6)	0.273	37 (48.1)	12 (48.0)	0.996
Perfectionism (EDI-3)	15 (30.0)	25 (48.1)	0.062	28 (39.4)	12 (38.7)	0.945	35 (37.2)	5 (62.5)	0.160	33 (36.3)	7 (63.6)	0.079	26 (33.8)	14 (56.0)	0.048
Asceticism (EDI-3)	17 (34.0)	29 (55.8)	0.027	26 (36.6)	20 (64.5)	0.009	40 (42.6)	6 (75.0)	0.077	39 (42.9)	7 (63.6)	0.191	28 (36.4)	18 (72.0)	0.002
Maturity Fears (EDI-3)	19 (38.0)	33 (63.4)	0.010	34 (47.9)	18 (58.1)	0.344	48 (51.1)	4 (50.0)	0.954	45 (49.5)	7 (63.6)	0.374	32 (41.6)	20 (80.0)	0.001
Total Problems (MASC 2)	4 (8.0)	9 (17.3)	0.159	8 (11.3)	5 (16.1)	0.498	12 (12.8)	1 (12.5)	0.983	11 (12.1)	2 (18.2)	0.567	9 (11.7)	4 (16.0)	0.574
Total score (CDI 2)	16 (32.0)	30 (57.7)	0.009	27 (38.0)	19 (61.3)	0.030	42 (44.7)	4 (50.0)	0.772	40 (44.0)	6 (54.5)	0.505	31 (40.3)	15 (60.0)	0.085

The Pearson's chi-square test was used to evaluate the association between the presence of FA's symptoms (+ present; - absent) and clinical scores at YSR, EDI-3, MASC 2 and CDI 2 scales. Statistical significance for p<0.05. CDI 2, Children Depression Inventory 2; EDI-3, Eating Disorder Inventory- 3; MASC 2, Multidimensional Anxiety Scale for Children 2; YSR, Youth Self Report.