

Supplementary Information

Valorization of Fine Recycled Aggregates Contaminated with Gypsum Residues: Characterization and Evaluation of the Risk for Secondary Ettringite Formation

Charlotte Colman ^{1,2,*}, David Bulteel ¹, Sébastien Rémond ³, Zengfeng Zhao ² and Luc Courard ²

¹ Laboratoire de Génie Civil et géo-Environnement, Université de Lille, IMT Lille Douai, ULR 4515 - LGCgE, F-59000 Lille, France; david.bulteel@imt-lille-douai.fr

² Urban and Environmental Engineering, GeMMe Building Materials, University of Liège, 4000 Liège, Belgium; zengfeng.zhao@uliege.be (Z.Z.); luc.courard@uliege.be (L.C.)

³ Université d'Orléans, Université de Tours, INSA CVL, EA 7494 LaMé, France; sebastien.remond@univ-orleans.fr

* Correspondence: charlotte.colman@uliege.be

Received: 2 October 2020; Accepted: 26 October 2020; Published: date

Table S1. T-test results of the swelling curves. A red color means that samples differed significantly from each other for $p < 0.05$.

	A- 0.19%	B- 0.08%	C- 0.29%	D- 0.62%	E- 0.61%	F- 0.18%	REF- 0.47%	REF- 3.08%	NS- 0.29%
A-0.19%									
B-0.08%									
C-0.29%									
D-0.62%									
E-0.61%									
F-0.18%									
REF-0.47%									
REF-3.08%									
NS-0.29%									




Figure S1. Swelling curve corresponding to Figure 3a, shown with error bars.

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).