

Features of the Preparation and Luminescence of Langmuir-Blodgett Films Based on the Tb(III) Complex with 3-Methyl-1-Phenyl-4-Stearoylpyrazol-5-One and 2,2'-Bipyridine

Julia Devterova ¹, Kirill Kirillov ², Anton Nikolaev ¹, Michail Sokolov ², Victor Shul'gin ³, Alexey Gusev ³, Victor Panyushkin ¹ and Wolfgang Linert ^{4,*}

¹ Department of Chemistry and High Technology, Kuban State University, 350040 Krasnodar, Russia; devterova8julia@gmail.com (J.D.); map_kompass@mail.ru (A.N.); panyushkin@chem.kubsu.ru (V.P.)

² Department of Physics and Technics, Kuban State University, 350040 Krasnodar, Russia; kirillka.kir@bk.ru (K.K.); sokolovme@mail.ru (M.S.)

³ Institute of Biochemical Technology, Ecology and Pharmacy, V.I. Vernadsky Crimean Federal University, 295007 Simferopol, Russia; shulvic@gmail.com (V.S.); galex0330@gmail.com (A.G.)

⁴ Institute of Applied Synthetic Chemistry, Vienna University of Technology, A-1060 Vienna, Austria

* Correspondence: wolfgang.linert@tuwien.ac.at; Tel.: +43-(1)-58801-163613

Figures




Figure S1. IR spectra of 1) HL (powder), 2) TbL₃·bipy (powder), 3) 10-layer TbL₃·bipy-based LBF obtained on subphase with C(Tb³⁺) = 5·10^{−4} M, 4) 10-layer TbL₃·bipy-based LBF obtained on subphase without terbium ions, C(Tb³⁺) = 0.0 M.

