

Visualization of Swift Ion Tracks in Suspended Local Diamondized Few-Layer Graphene

Nadezhda A. Nebogatikova ^{1,*}, Irina V. Antonova ^{1,2}, Anton K. Gutakovskii ^{1,3}, Dmitriy V. Smovzh ⁴, Vladimir A. Volodin ^{1,3} and Pavel B. Sorokin ^{5,*}

¹ Rzhanov Institute of Semiconductor Physics of the Siberian Branch of the RAS, Novosibirsk 630090, Russia; ifp@isp.nsc.ru (I.V.A.); inter@nstu.ru (A.K.G.); director@itp.nsc.ru (V.A.V.)

² Department of Semiconductor Devices and Microelectronics, Novosibirsk State Technical University, Novosibirsk 630087, Russia

³ Physical Department, Novosibirsk State University, Novosibirsk 630090, Russia

⁴ Kutateladze Institute of Thermophysics, Siberian Branch of the Russian Academy of Sciences, Novosibirsk 630090, Russia; rector@nsu.ru

⁵ Technological Institute for Superhard and Novel Carbon Materials, Moscow 108840, Russia

* Correspondence: nadonebo@gmail.com (N.A.N.); pbsorokin@tisnum.ru (P.B.S.)




Figure S1: Cluster of nanocrystals at the periphery of the ion track at the irradiation dose of 1×10^{11} ion/cm², ion energy is 167 MeV.




Figure S2: Cluster of nanocrystals at the periphery of the ion track at the irradiation dose of 1×10^{11} ion/cm², ion energy is 26 MeV.




Figure S3: Cluster of nanocrystals at the periphery of the ion track at the irradiation dose of 5×10^{11} ion/cm², ion energy is 167 MeV.