

Supplementary Materials

Canopy Temperature Differences between Liana-Infested and Non-liana Infested Areas at a Neotropical Dry Forest

Authors: Xu Yuan, Kati Laakso, Philip Marzahn and G. Arturo Sanchez-Azofeifa

Table S1. The results of the Levene's Test.


		Levene Statistic	Df1	Df2	Sig.
Temperature	Based on mean	1.899	2	1149	0.150


Table S2. The results of the Kolmogorov-Smirnov Test.

	Statistic	Df	Sig.
Residual for temperature	0.045	1152	0.000

Table S3. Statistical characteristics of the canopy temperatures of the liana-infested areas, non-liana infested areas and gaps of the subsample ($n = 384$), Max.: Maximum, Min.: Minimum, Std. dev.: Standard deviation.

	Mean /°C	Median /°C	Std. Dev.	Min. /°C	Max. /°C
Liana-infested areas	26.69	26.77	0.85	24.42	28.87
Non-liana infested areas	26.28	26.24	0.86	24.16	29.32
Gaps	26.16	26.14	0.79	24.40	28.34





(c)

Figure S1. Temperature distributions of the (a) liana-infested areas ($n = 225,647$), (b) non-liana infested areas ($n = 420,811$) and (c) gaps ($n = 307,126$).