

Article

Development and Characterization of Simple Sequence Repeat Markers for, and Genetic Diversity Analysis of *Liquidambar formosana*

Siyuan Chen ¹ , Mingliang Dong ¹, Yan Zhang ^{1,2}, Shuaizheng Qi ¹, Xuezeng Liu ³, Jinfeng Zhang ^{1,*} and Jian Zhao ^{1,*}

¹ Beijing Advanced Innovation Center for Tree Breeding by Molecular Design, National Engineering Laboratory for Tree Breeding, Key Laboratory of Genetics and Breeding in Forest Trees and Ornamental Plants of Ministry of Education, Key Laboratory of Forest Trees and Ornamental Plants Biological Engineering of State Forestry Administration, College of Biological Sciences and Biotechnology, Beijing Forestry University, Beijing 100083, China; bjfucsy@163.com (S.C.); dmllinxue@163.com (M.D.); zhangyan19890802@126.com (Y.Z.); 13051857171@163.com (S.Q.)

² College of Landscape Architecture, Beijing University of Agriculture, Beijing 102206, China

³ Henan Longyuan Flowers & Trees Co., Ltd., Yanling 461200, China; lxuezeng@126.com

* Correspondence: zjf@bjfu.edu.cn (F.Z.); zhaojian0703@bjfu.edu.cn (J.Z.); Tel.: +86-1352-266-8293 (F.Z.); +86-1520-144-2430 (J.Z.)

Received: 29 November 2019; Accepted: 24 January 2020; Published: 12 February 2020



Abstract: *Liquidambar formosana* (Hamamelidaceae) is a relatively fast-growing deciduous tree of high ornamental value that is indigenous to China. However, few molecular markers are available for the species or its close relatives; this has hindered genomic and genetic studies. Here, we develop a series of transferable expressed sequence tag-simple sequence repeats (EST-SSRs) for genomic analysis of *L. formosana*. We downloaded the sequence of the *L. formosana* transcriptome from the National Center of Biotechnology Information Database and identified SSR loci in the Unigene library. We found 3284 EST-SSRs by mining 34,491 assembled unigenes. We synthesized 100 random primer pairs for validation of eight *L. formosana* individuals; of the 100 pairs, 32 were polymorphic. We successfully transferred 12 EST-SSR markers across three related *Liquidambar* species; the markers exhibited excellent cross-species transferability and will facilitate genetic studies and breeding of *Liquidambar*. A total of 72 clones of three *Liquidambar* species were uniquely divided into three main clusters; principal coordinate analysis (PCoA) supported this division. Additionally, a set of 20 SSR markers that did not exhibit nonspecific amplification were used to genotype more than 53 *L. formosana* trees. The mean number of alleles (N_a) was 5.75 and the average polymorphism information content (PIC) was 0.578, which was higher than that of the natural *L. formosana* population (0.390). In other words, the genetic diversity of the plus *L. formosana* population increased, but excellent phenotypic features were maintained. The primers will be valuable for genomic mapping, germplasm characterization, gene tagging, and further genetic studies. Analyses of genetic diversity in *L. formosana* will provide a basis for efficient application of genetic materials and rational management of *L. formosana* breeding programs.

Keywords: EST-SSR markers; *Liquidambar formosana*; plus trees; transcriptome; cross-amplification; genetic diversity

1. Introduction

Liquidambar formosana is a large deciduous tree in the genus *Liquidambar*. There are four principal species in the genus: *L. formosana*, *L. styraciflua*, *L. acalycina*, and *L. orientalis*. *L. formosana* is distributed

principally in East Asia, including China, Laos, Vietnam, Japan, and the Korean Peninsula, from 18° to 38° north latitude [1]. The tree is generally found on plains and hills at 1000–1500 m above sea level, and requires abundant sunshine and moisture for growth. Given the increasing demand for wood, fast-growing trees yielding high-quality wood that are amenable to afforestation are urgently needed [2]. *Liquidambar*, especially *L. styraciflua* and *L. formosana*, generally meet these requirements [3,4]. *L. formosana* is of high economic, ornamental, and ecological value, and is attracting increasing attention in terms of wood production, landscaping, and medicinal use. The wood is compact, soft, and easy to process and handle; the roots, leaves, bark, and inflorescence resin can be used in medicines [5]. The resin (in particular) is used for detoxification, pain relief, to establish haemostasis, and to encourage muscle growth, and also serves as a perfume fixative [6]. The trunk is thick, straight, and imposing in appearance; the crown is wide and the leaves have an attractive red colour in the late autumn. *L. formosana* exhibits marked adaptability in the wild and is highly resistant to toxic gases, especially SO₂ and chlorine, and is thus often used to green factories and mines [7]. This fast-growing species is recognized as a “pioneer tree” for afforestation.

The cultivation, asexual propagation, afforestation, and genetic breeding of *Liquidambar* (especially *L. styraciflua*) have been extensively studied; however, breeding research is less advanced. Winstead (1971) reported that various seed sources differed in the amount of time required for germination; the germination rates differed significantly [8]. Suhaendi (1989) explored 12 provenances of *L. styraciflua* and found that the Honduran provenance (LosAkoos-46/83) exhibited the highest germination rate [9]. Kariuki (1989) reported significant differences in seed germination rate among trees from different provenances [10]. He (2005) tested 20 provenances of *L. formosana* from eight provinces in China, and reported highly significant differences in tree height and ground-level diameter [11]. These studies provided valuable materials and allowed researchers to gain experience useful for breeding improved *formosana* varieties. Shi et al. (1997) selected 90 superior *L. formosana* trees from Anhui Province, China [12]. Zhao et al. (2009) optimized the secondary forests of *L. formosana* in 10 counties of Guizhou Province, China. Compared to the control group, the average tree height, diameter at breast height, and volume increased by 12.7, 24.4, and 61.2%, respectively [13]. Individual trees, or groups, exhibiting valuable traits were selected for breeding; the useful traits were shown to be stably inheritable. However, the vast majority of tree species remain in a wild or semi-wild state. Genetic variation is thus large, such that there is great potential for the selection of valuable traits [14]. However, marker-assisted selection (MAS) of *L. formosana* and related species is difficult, given the lack of molecular markers. MAS enhances breeding efficiency; target genotypes are directly selected [15]. Little is known about species-specific diagnostic genetic relationships among, or the genetic diversity of populations of, *L. formosana*. Vendrame et al. (2001) used RAPD markers to show that the somatic embryo seedlings of hybrid sweetgum were heterozygous [16]. Bi et al. (2010) analyzed genetic diversity in the natural populations of *Liquidambar formosana* revealed by ISSR molecular markers [17]. Sun et al. (2016) developed and screened 14 pairs of simple sequence repeat (SSR) primers based on transcriptome data, and analyzed the genetic diversity of 25 populations and 691 individual plants over the entire distributional range of *L. formosana* [18]. Genetic diversity is the principal form of biodiversity, providing excellent evidence of species formation, molecular evolution, and geographical variation. The extent and distributional pattern of intraspecific genetic variation determine whether a species can cope with environmental mutations. The genetic diversity of plus trees allows us to understand the genetics of superiority, and guides rational selection of the second generation of such trees [19,20]. However, compared to other economically important tree species, few genetic studies on *L. formosana* have appeared.

SSRs, also termed microsatellites, are tandem repeat sequences in which at least one nucleotide serves as the basic unit; they are widely dispersed in the genomes of eukaryotes, being present in non-coding, coding, and other chromosomal regions. SSRs are currently divided into genomic SSRs and expressed sequence tagged-SSRs (EST-SSRs), and are universally accepted as powerful molecular tools facilitating MAS, genetic linkage mapping, analysis of genetic diversity, and cultivar

identification [21,22]. EST-SSRs are found in coding regions, affording the advantages of co-dominance, wide coverage, simple detection, convenient operation, rapid action, and high-level polymorphism. EST-SSR markers exhibit excellent cross-species transferability because they are designed using highly conserved transcriptome data [23,24]. For estimation of genetic advantage, EST-SSRs have been developed for many trees and crops (including *Gossypium*, *Populus*, and the *Leguminosae*) [25–27]; these have been used to create high-quality genetic maps, to perform comparative genomic studies, and to facilitate MAS. For *L. formosana*, the 14 currently available EST-SSRs pairs are inadequate for detailed genetic research [18]. Consequently, a set of polymorphic EST-SSR markers for *L. formosana* based on transcriptome data is urgently needed.

Here, we obtained transcriptome data from the National Center of Biotechnology Information, generated unigenes, and identified many useful EST-SSRs for *L. formosana*. We created a new series of EST-SSR markers and tested their cross-species transferability among *Liquidambar* species. We used the original EST-SSR markers to investigate the genetic diversity of plus *L. formosana* trees in Henan Province. The new EST-SSR markers will serve as valuable molecular tools for further genetic and breeding research on *L. formosana* and related species. Analysis of genetic diversity will provide significant information on the plus *L. formosana* population. Evaluation of genetic variation after selection could underpin efficient management and selection of hybrid parents among *Liquidambar* species.

2. Materials and Methods

2.1. Plant Materials

On Jigong Mountain (Xinyang, Henan Province, China, 31°52′ N, 114°05′ E), we selected branches from 60 plus *L. formosana* trees and hydroponically cultured them in the greenhouse of Beijing Forestry University (Haidian District, Beijing, P.R. China; 39°95′ N, 116°30′ E). In the selection process, the candidate plus tree are selected as the center, and within the radius of 10–25m, three plus trees rank only second to the candidate plus tree are selected. The height, diameter at breast height (DBH) and under branch height of the candidate plus tree should be greater than 105%, 120% and 110% of the average values of the three plus trees. In addition, the candidate plus trees should reach the mature stage, with large amount of flowering and fruit bearing, which is also required the trunk is straight, with obvious apical dominance. No plant diseases or insect pests. Finally, tender leaves of 53 individuals were collected and stored at −80 °C prior to DNA extraction due to some samples died during hydroponics. During primer screening, two individuals of *L. formosana* were randomly selected to detect amplification, and eight were used to detect EST-SSR polymorphisms. To test cross-species transferability, six *L. styraciflua* were collected from Shanghai Chenshan Botanical Garden (Songjiang District, Shanghai, China, 31°03′ N, 121°22′ E); 13 hybrid sweetgum (*L. styraciflua* × *L. formosana*) obtained via hybridization [28] were used to test SSR transferability. The *L. formosana* and *L. styraciflua* were native population while hybrid sweetgum was from cultivated population. In total, 72 sampled *Liquidambar* trees were used and details for the 72 individuals can be obtained from supplementary Table S1.

2.2. Transcriptome Assembly, SSR Mining, and Primer Design

The sequence of the *L. formosana* transcriptome was downloaded from <https://www.ncbi.nlm.nih.gov/> (accession numbers SRR1514949 and SRR1514913) [29]; the sequences were assembled and SSR loci were identified in the Unigene library using MISA (MIcroSATellite identification tool, <http://pgrc.ipk-gatersleben.de/misa/>). The search criteria were as follows: minimum repetitions of di-, tri-, tetra-, penta-, and hexa-nucleotides of nine, six, five, five, and four respectively (the SSR search criteria did not include single-nucleotide repetitions). Primer3 software [30] was used to design the EST-SSR primers after excluding fragments with inappropriate or excessively short flanking sequences. To avoid hairpin formation and mispairing, the primers with the highest scores were selected and stored in a primer library. The specific primer design parameters were as follows: (1) length, 18 to

22 bp; (2) annealing temperature, 58 to 61 °C (optimum, 60 °C); (3) GC content, 40 to 60%; and (4) expected PCR product length, 150 to 300 bp.

2.3. DNA Extraction, PCR Amplification, and Detection of Polymorphisms

Total genomic DNA was extracted using a modified cetyltrimethylammonium bromide method [31], and yields were determined with the aid of a NanoDrop 2000 spectrophotometer (Thermo Scientific, Wilmington, DE, USA). DNA was diluted to 50–100 ng/μL and DNA quality was evaluated via 0.8% (w/v) agarose gel electrophoresis prior to subsequent analysis. The PCR amplifications afforded by 100 synthetic primers in eight randomly selected *L. formosana* were evaluated prior to further genetic research. Each reaction mixture (20 μL) contained 2 μL template DNA (50–100 ng/μL), 10 μL 2× TaqMaster Mix (Biomed-Tech, Beijing, China), and 4 μL of primer solution (volume ratio of forward:reverse primers 1:4). We ensured that the fragments created using mixed primers were long, to guarantee that detection via fluorescent capillary electrophoresis (the next step) was not compromised by unwanted fluorescent signals. The PCR conditions were as follows: 95 °C for 5 min, followed by 30 cycles at 94 °C for 30 s, 56 °C for 30 s, and 72 °C for 30 s, 8 cycles at 94 °C for 30 s, 54 °C for 30 s, 72 °C for 30 s, and a final extension at 72 °C for 10 min. The amplified products were subjected to 2% (w/v) agarose gel electrophoresis. EST-SSR primers that yielded clear bands were employed for wholesale genotyping, and for analysis of genetic diversity and cross-species transferability, using M13 technology [32] with fluorescent labels (FAM, HEX, TAMRA, and ROX), and same dye always used with each locus. The sequence of m13 was TGTAACGACGGCCAGT. The PCR products were analyzed using a capillary-based ABI 3730XL DNA Analyzer, and amplicon sizes were measured by a Gene Mapper (Applied Biosystems, Foster City, CA, USA).

2.4. Genetic Diversity of Plus Trees of *L. formosana*, and Statistical Analysis

Of the 32 pairs of polymorphic primers developed, 16 pairs that did not show non-specific amplification were selected. Additionally, the four SSR pairs developed by Sun et al. [18] were employed; these exhibited high-level capacity to detect polymorphisms and were well-amplified. A total of 20 SSRs were thus used for PCR amplification of 53 plus *L. formosana* individuals. All fluorescent capillary electrophoresis data were recorded in Microsoft Excel 2007 (Microsoft Corp., Redmond, WA, USA). Convert software (ver. 1.3) [33] was used to transform the data into formats that could be analyzed by other biological software. POPGENE (ver. 1.3.2) [34] was employed to estimate genetic parameters, such as the number of alleles (N_a), effective allele number (N_e), Shannon information index (I), observed heterozygosity (H_o), and expected heterozygosity (H_e). Each locus was subjected to Hardy–Weinberg equilibrium (HWE) testing using POPGENE (ver. 1.32) and the chi-squared test. The polymorphism information content (PIC) of each locus was calculated using Pic_Calc (ver. 0.6) [35]. The population genetic structures of three *Liquidambar* species were analyzed by Structure software (ver. 2.3.3) and the true value of clusters (K) were harvested using log probability of data according to the method of G. Evanno (2005), when delta K is maximum, the k value is suitable for numbers of groups [36] running a mixed model. PowerMarker (ver. 3.25) [37] was employed to calculate the relationship between any two clones based on the Nei genetic distance, and principal component analysis (PCA) was performed using GenALEX software (ver. 6.2) [38].

3. Results

3.1. The Frequency and Distribution of SSR Loci

Unprocessed data obtained via high-throughput sequencing were pre-processed and redundant low-quality sequences were removed. Finally, a total of 34,491 unigenes were defined in the *L. formosana* transcriptome, with an average length 699 bp. SSR loci were then searched for in MISA. Table 1 showed that the total sequence length was 24,122,122 bp; 3284 SSR loci were found (frequency, 9%). On average, one SSR locus was seen every 7.3 kb. Of these, 298 unigene loci containing more than one SSR locus accounted for 10% of the total, and 109 complex SSR loci constituted 3% of the total.

Table 1. Character of Unigenes in *L. formosana*.

Sequence Type	Total
Co-detected Unigenes Length (kbp)	24,122
Average Length of Unigenes (bp)	699
Number of SSR loci detected	3284
Number of Unigenes Containing SSR Loci	2949
Number of Unigenes with more than one locus	298
Number of complex SSR loci	109
Frequency of SSR loci	1/7.3 kbp

As high-throughput transcriptome sequences sometimes contain errors, we did not consider single-nucleotide SSR sites. Table 2 shows that the remaining 2742 SSR loci included five SSR repeat motifs (di-, tri-, tetra-, penta- and hexa-nucleotide repeats), of which the principal type was the dinucleotide (52.56%), followed by the trinucleotide (33.80%). The most abundant SSR loci contained six serial repeats (16.67% of all SSR loci), followed by loci with nine repeats (15.50%). Of the dinucleotide repeats, AG/CT exhibited the highest frequency (Figure 1) (92.51%), followed by AC/GT (5.27%) and AT/AT (2.22%). Of the trinucleotide repeats, AAG/CTT was the most abundant (31.18%), followed by ACC/GGT (20.50%) and AGC/CTG (13.48%). Of the tetranucleotide repeats, AAAT/ATTT exhibited the highest frequency (29.69%), followed by AAAG/CTTT (25.00%) and ACAT/ATGT (9.38%). Of the pentanucleotide repeats, AAAAG/CTTTT was the most common (17.24%), followed by AAGAG/CTCTT and AAAGG/CCTTT (each, 10.34%). The frequency of the hexanucleotide repeat AGAGGG/CCCTCT was 6.76%.

Table 2. Frequency of different repeat motifs of SSRs in the transcriptome of *L. formosana*.

Repeat Motif	Repeat Number										Percentage (%)
	4	5	6	7	8	9	10	11	>11	Total	
Dinucleotide						344	298	209	590	1441	52.56
Trinucleotide			427	234	104	80	43	7	32	927	33.8
Tetranucleotide		47	10	4	3					64	2.33
Pentanucleotid-e		20	9							29	0.88
Hexonucleotid-e	205	57	11	6	1	1				281	10.58
Total	205	124	457	244	108	425	341	216	622	2742	100
Percentage (%)	7.48	4.52	16.67	8.9	3.94	15.5	12.44	7.88	22.68	100	52.56




Figure 1. Frequency distribution of different SSR (simple sequence repeat) motifs of the specific SSR repeat motifs in the transcriptome of *L. formosana*.

3.2. Synthesis, Screening, and Polymorphisms of EST-SSR Primers

Primer software (ver. 3.0) was used to design 2742 SSR loci; some loci were unusable because the flanking sequences were inadequate or they did not meet primer design standards. We finally selected 100 pairs of synthetic primers. The amplifications obtained were preliminarily screened using two randomly selected individual DNAs of *L. formosana*. Seventy-two pairs of primers yielding amplified bands were selected (effective amplification rate, 72%). Eight randomly selected DNA samples of *L. formosana* were amplified by PCR and products were detected via fluorescence capillary electrophoresis; 32 pairs of polymorphic primers (32%) were found. The information afforded by each pair is shown in Table 3. The polymorphisms amplified by primer pair Liq_eSSR10, Liq_eSSR17 and Liq_eSSR59 in *L. formosana* samples are selected randomly to discuss here (Figure 2).

Table 3. Information on the 32 polymorphic SSR (simple sequence repeat) markers developed for *L. formosana*.

Code	Locus	Primer Sequence	Repeat Motif	Tm (°C)	Product Size (bp)	Dye
1	Liq_eSSR1	F: TGTAACGACGGCCA GTCATCATCAGACACGGTTTGG R: GTCCAGCTCCGTACATCCAT	(TTAGGG)4	60	217	HEX
2	Liq_eSSR2	F: TGTAACGACGGCCAG TATGGGACGATGACTGTGTT R: CGTTGCAAATCTCATTGACG	(ACGGCA)4	60	200	ROX
3	Liq_eSSR4	F: TGTAACGACGGCCAG TAGTGGTTGTTGCTGCTGTTG R: CTTATGCAGCCCCAAATGTT	(CTGGTG)7	60	206	FAM
4	Liq_eSSR7 ¹	F: TGTAACGACGGCCAG TCAAGTAGTCCAGGGCGTCTC R: CCTAACGAGAAGGGAGGCTT	(CTCTTC)4	60	192	ROX
5	Liq_eSSR10 ¹	F: TGTAACGACGGCCAG TAGACCACATGCTCGTTACCC R: TGCTGGAAGTGGTGTAGTG	(CTAGCA)5	60	142	FAM
6	Liq_eSSR13	F: TGTAACGACGGCCAG TTCTGGGTCAGGTTACAGGTTTC R: GCCATATCCCTCACCATGTC	(GGTTCA)4	60	24	TAMRA
7	Liq_eSSR16	F: TGTAACGACGGCCAG TTGGCAGATCTAGGGGATTG R: AACCTCCATCTCCACGATTG	(GAGGTG)4	60	203	FAM
8	Liq_eSSR17 ¹	F: TGTAACGACGGCCAG TAGAGGTGAATGGGACACCAG R: CTGCCAACCTGGGAATAGAA	(GGATGA)4	60	206	HEX
9	Liq_eSSR25 ¹	F: TGTAACGACGGCCAG TTCAACTTGATGGGGTCAGTG R: GACTGCCCGTACCCAGAATA	(TATG)6	60	194	ROX
10	Liq_eSSR28	F: TGTAACGACGGCCAG TACATCAGAAGGAGCCGAAGA R: CGGATTGGCAAGAGAAGAAG	(GGT)6	60	175	HEX
11	Liq_eSSR30 ¹	F: TGTAACGACGGCCAG TAGGCAGGGTTAGTCCCTTGT R: GCTTGCGGACTTGTCTCTC	(CTG)6	60	224	ROX
12	Liq_eSSR33 ¹	F: TGTAACGACGGCCAG TGAGGAGATTACACACAGGA R: TCATCATCAAGTCCGCCATA	(GCA)8	60	212	HEX
13	Liq_eSSR34 ¹	F: TGTAACGACGGCCAG TGCTGCCTCCAAAGAAGTGTC R: AGGGCTCCTCAGCTAGTTCC	(GCT)7	60	199	ROX
14	Liq_eSSR35 ¹	F: TGTAACGACGGCCAG TTTACCACCGTCGTCGTCAT R: GAACCAAGAGAAGCACCAGC	(TCC)8	60	165	FAM

Table 3. Cont.

Code	Locus	Primer Sequence	Repeat Motif	Tm (°C)	Product Size (bp)	Dye
15	Liq_eSSR42	F: TGTAACACGACGGCCAG TCACATCGGAAGCTCCTTCTC R: GCGACCCACAATACTAGGGA	(TC)10	60	201	HEX
16	Liq_eSSR43	F: TGTAACACGACGGCCAG TCCAAAGCTTCCTCCTCCTCT R: ACATTTCTTCCGCATTGACC	(CT)12	60	24	ROX
17	Liq_eSSR44	F: TGTAACACGACGGCCAG TCTTGGAATCACCTTCGCATT R: CGATCGAATTGTAAAGCGGT	(CA)9	60	198	FAM
18	Liq_eSSR46	F: TGTAACACGACGGCCAG TTCTCCATGGTCTTTCTTGG R: CCAAACAGGGTGGAGAGAAA	(AGAGA)6	60	162	ROX
19	Liq_eSSR48	F: TGTAACACGACGGCCAG TTCGATGATGAGAACACTGGAA R: ATCCGATAATAACGTCCCCC	(GATTT)6	60	163	FAM
20	Liq_eSSR51	F: TGTAACACGACGGCCAG TGAGAATGGCAGAGGAAGCAC R: AAGCCATGAAGAGAAGGCAA	(TGT)9	60	197	ROX
21	Liq_eSSR55 ¹	F: TGTAACACGACGGCCAG TCCTTCTTTGGATCTCCACCA R: TTGCACTTTGACTGAGACCG	(CTC)8	60	194	ROX
22	Liq_eSSR59 ¹	F: TGTAACACGACGGCCAG TCAAACGAAAGATGAGCGACA R: CCGTTTTTCATGGACCACTCT	(CAA)6	60	201	ROX
23	Liq_eSSR60 ¹	F: TGTAACACGACGGCCAG TGAGACCTGGTTGGACTTGGGA R: CGGAGAATGACCGGTCTAAA	(GGA)6	60	216	HEX
24	Liq_eSSR62 ¹	F: TGTAACACGACGGCCAG TCCTCGAACTGTTTCTCCAGC R: CGCTCCTTAAACCCTAACCC	(AGG)6	60	202	FAM
25	Liq_eSSR64	F: TGTAACACGACGGCCAG TCGGTACATGTGATCAGCCAC R: CTTCTGCTTCTCCAAATGGC	(GAA)7	60	169	ROX
26	Liq_eSSR67 ¹	F: TGTAACACGACGGCCAG TCCAAGGAGAGAGTGACGAGG R: ATTGATTTTTGCGGACTTGG	(CCT)8	60	273	ROX
27	Liq_eSSR70	F: TGTAACACGACGGCCAG TTGGGGGAATGAGAGTTTGAC R: ATCTCAGTGGGCTCTCCTCA	(GTG)7	60	174	HEX
28	Liq_eSSR86 ¹	F: TGTAACACGACGGCCAG TCTGAAAATAAGGCGGGTGAA R: GTCTGGGTCATCTTCGTCGT	(TC)9	60	172	HEX

Table 3. Cont.

Code	Locus	Primer Sequence	Repeat Motif	T _m (°C)	Product Size (bp)	Dye
29	Liq_eSSR89	F: TGTAACACGACGGCCAG TATCGTCCAGGCTTATCATCG R: GCCTCTGCTGACACTGACAA	(GTT) ⁷ gatc (GGCTGT) ⁵	60	201	FAM
30	Liq_eSSR90 ¹	F: TGTAACACGACGGCCAG TCCATTGTTCCCCCTACCTCT R: TGCAAGAACGAGAGCAGAGA	(ACC) ⁶ acttctctatca gcaaagccaccg cg(ACC) ⁷	60	196	ROX
31	Liq_eSSR92	F: TGTAACACGACGGCCAG TAGAGGACACCCAAGGGAAGT R: CCATTGTTCCCCCTACCTCT	(CCCTTG) ⁴ gagagagaaag cgatag(GT) ¹¹ atagcgagag agaa(AG) ⁹ (ACC) ⁶ ac ttctctatcagc aaagccacc gcg(ACC) ⁷	60 60	181	HEX
32	Liq_eSSR98 ¹	F: TGTAACACGACGGCCA GTTCCCAATCTCAATCTCGACC R: CCTCCGTAATCCGAAGAACA	(CCTCAA) ⁴	60	220	ROX

¹ EST-SSR markers were selected for the analysis of genetic diversity and cross-species transferability.




Figure 2. The polymorphisms amplified by primer pair Liq_eSSR10 (blue), Liq_eSSR17 (green) and Liq_eSSR59 (red) in *L. formosana* samples.

3.3. Genetic Diversity of Plus Trees of *L. Formosana*

A total of 115 alleles (Na) from 20 SSR loci were amplified from 53 individuals of *L. formosana*. The number of Liq_eSSR35 alleles was highest (13), while the numbers of Liq_eSSR17 and Liq_eSSR34 alleles were lowest (3). An average of 5.75 alleles were amplified from each locus; 3.16 effective alleles (Ne) were amplified from 20 SSR loci; the maximum number (Liq_eSSR35) was 8.87 and the minimum number (Liq_eSSR34) 1.28. The average number of effective alleles/locus was 3.17 (Table 4). The Shannon diversity index (I) was highest at Liq_eSSR35 (2.317); the average value was 1.250. The observed heterozygosity (Ho) and expected heterozygosity (He) were highest for Liq_eSSR35 (0.978 and 0.887, respectively). The average Ho was 0.653 and the average He was 0.621. PIC is useful for measuring SSR locus polymorphisms, reflecting whether a locus can distinguish among populations. If the PIC is >0.5, the locus is highly polymorphic; if it is between 0.25 and 0.5, it is moderately polymorphic; and if it is <0.25, it is poorly polymorphic; the average PIC was 0.578. We analyzed plus *L. formosana* trees; the PIC of each locus reflected the genetic diversity. The average PIC was 0.578, significantly higher than the average PIC (0.309) of a natural population of *L. formosana*. Thus, the genetic diversity of the dominant population did not decrease; in fact, it increased, although phenotypic dominance was maintained. In the HWE test, five SSR loci (Liq_eSSR33, Liq_eSSR35, Liq_eSSR7, Liq_eSSR90, and Liq_eSSR25) deviated significantly from equilibrium ($P < 0.05$); all exhibited heterozygote overabundance (Table 4).

Table 4. Diversity information parameters for 20 SSR loci in 53 individuals of *L. formosana*.

Locus	Na ¹	Ne ²	Ho ³	He ⁴	I ⁵	PIC ⁶	HW ⁷
Liq_eSSR35	13	8.872	0.978	0.887	2.317	0.877	*
Liq_eSSR86	10	5.25	0.925	0.81	1.882	0.785	ns
Liq_eSSR67	7	4.86	0.906	0.794	1.712	0.765	ns
Liq_eSSR90	7	3.959	0.698	0.747	1.565	0.71	*
Liq_eSSR98	6	3.822	0.736	0.738	1.469	0.694	ns
Liq_eSSR25	6	3.451	0.491	0.71	1.459	0.671	*
Liq_eSSR10	5	3.216	0.736	0.689	1.3	0.632	ns
Liq_eSSR30	7	2.814	0.711	0.645	1.339	0.608	ns
LF39 ⁸	6	2.681	0.577	0.627	1.337	0.599	ns
LF17 ⁸	4	2.825	0.774	0.646	1.097	0.572	ns
Liq_eSSR33	4	2.688	0.962	0.628	1.133	0.568	***
Liq_eSSR62	5	2.636	0.642	0.621	1.137	0.558	ns
LF37 ⁸	6	2.452	0.604	0.592	1.149	0.545	ns
Liq_eSSR60	4	2.465	0.547	0.594	1.094	0.543	ns
Liq_eSSR59	3	2.485	0.64	0.598	0.998	0.528	ns
LF15 ⁸	5	2.016	0.566	0.504	1	0.47	ns
Liq_eSSR17	3	2.158	0.679	0.537	0.883	0.458	ns
Liq_eSSR55	7	1.683	0.457	0.406	0.911	0.388	ns
Liq_eSSR7	4	1.751	0.188	0.429	0.755	0.372	*
Liq_eSSR34	3	1.286	0.245	0.222	0.454	0.211	ns
Mean	5.75	3.169	0.653	0.621	1.25	0.578	ns

¹ Observed number of alleles, ² effective number of alleles, ³ observed heterozygosity, ⁴ expected heterozygosity,

⁵ Shannon's information index, ⁶ PIC polymorphism information content, ⁷ Chi-squared tests for Hardy–Weinberg equilibrium, ns not significant, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, ⁸ EST-SSR markers were selected from previous study [6].

3.4. Cross-Transferability of EST-SSR Primers

Twenty EST-SSR primers were further tested in two related *Liquidambar* species; *L. styraciflua* ($N = 6$) and hybrid sweetgum ($N = 13$). Na and PIC were evaluated to measure the cross-transferability of the 20 primer pairs (Table 5.). Nineteen of the 20 primer pairs showed amplification in hybrid sweetgum (95%), as did 20 pairs in *L. styraciflua* (100%); five pairs showed monomorphic amplification. 12 primer pairs of the newly developed EST-SSRs in this study presented highly polymorphism among

the two related species, with average PIC values 0.502 in hybrid sweetgum and 0.452 in *L. styraciflua*; average Na 3.89 in hybrid sweetgum and 0.452 in *L. styraciflua*. Thus, these primers could be used to analyze other species of *Liquidambar*.

Table 5. Cross-species amplification of 20 SSR loci in other *Liquidambar* species as demonstrated by two genetic diversity indices (Na and PIC).

Locus	hybrid Sweetgum			<i>L. styraciflua</i>		
	PIC ¹	Na ²	Transferability ³	PIC ¹	Na ²	Transferability ³
LF15	0.577	4	++	0.000	1	+
LF17	0.492	4	++	0.476	3	++
LF19	0.552	4	++	0.346	2	++
LF37	0.419	3	++	0.240	2	++
Liq_eSSR 7	-	0	-	0.000	1	+
Liq_eSSR 10 ⁴	0.519	4	++	0.240	2	++
Liq_eSSR 17	0.335	2	++	0.000	1	+
Liq_eSSR 25 ⁴	0.640	4	++	0.536	3	++
Liq_eSSR 30 ⁴	0.580	3	++	0.346	2	++
Liq_eSSR 33	0.496	4	++	0.000	1	+
Liq_eSSR 34 ⁴	0.666	4	++	0.476	3	++
Liq_eSSR 35 ⁴	0.445	4	++	0.620	4	++
Liq_eSSR 55 ⁴	0.558	4	++	0.460	3	++
Liq_eSSR 59	0.205	4	++	0.000	1	+
Liq_eSSR 60 ⁴	0.686	6	++	0.643	5	++
Liq_eSSR 62 ⁴	0.575	4	++	0.346	2	++
Liq_eSSR 67 ⁴	0.555	3	++	0.240	2	++
Liq_eSSR 86 ⁴	0.645	5	++	0.393	4	++
Liq_eSSR 90 ⁴	0.746	6	++	0.643	5	++
Liq_eSSR 98 ⁴	0.417	3	++	0.476	4	++
Mean	0.505	3.72		0.324	2.55	

¹ Observed numbers of alleles, ² polymorphism information content, ³ polymorphism was expressed as ++, monomorphism as +, and absence of primer amplification as -, ⁴ 12 pairs of original EST-SSR markers with high level of polymorphism in two *Liquidambar* species.

3.5. Population Genetic Structure Analysis, Cluster Analysis, and PCA of Three Species of *Liquidambar*

STRUCTURE software (ver. 2.3.3) was used to analyze genetic structure of 72 individuals of three species of *Liquidambar*. According to Evanno, the maximum K represents the optimal number of clusters and K = 2 was the largest value; the *Liquidambar* individuals should thus be divided into two groups (Figure 3). 72 individuals were divided into two main gene pools. Here, the male parent of the hybrid sweetgum was *L. formosana* (more red) and the female parent was *L. styraciflua* (more green). As the proportion of green in the hybrid sweetgum was much higher than that of red, the hybrid was grouped with *L. styraciflua*. Cluster analysis based on the neighbour-joining (NJ) method was established to demonstrate the relationships between 72 individuals of three species of *Liquidambar*; *L. formosana* was clearly clustered apart from hybrid sweetgum and *L. styraciflua*, when we focused on hybrid sweetgum and *L. styraciflua*, cluster analysis can clearly separate *L. styraciflua* from hybrid sweetgum. (Figure 4). *L. styraciflua* and hybrid sweetgum were closely related, confirming the results of the genetic structure analysis. PCoA analysis demonstrated the dispersion of the three *Liquidambar* species through a two-dimensional scatter plot, which divided the 72 individuals into three groups, In particular, *L. formosana* was well-grouped and the closer relationship between hybrid sweetgum and *L. styraciflua* was proved as well. The results (Figure 5) showed that coordinates 1 differentiate Group I from Group II and Group III and explained for 41.22% of the total variation. Coordinates 2 differentiate Group II from Group III and accounted 15.63% of the total variation. The results were in agreement with those of cluster analysis and population genetic structure analysis.




Figure 3. Results of STRUCTURE analysis for 72 *Liquidambar* individuals based on 20 EST-SSR markers. (a) Estimation of population using log probability of data-derived delta K with cluster number (K) ranged from one to eight. (b) The 72 individuals are separated into two main gene pools (K = 2), Proportion of green (*L. styraciflua*) in the hybrid sweetgum was much higher than that of red (*L. formosana*), the hybrid was grouped with *L. styraciflua*.




Figure 4. Genetic relationship among 72 individuals of three species *Liquidambar* based on 20 EST-SSR markers using neighbour-joining clustering. The 72 individuals were divided into three clusters, which are depicted with green (*L. formosana*), blue (hybrid sweetgum), and red (*L. styraciflua*) solid, respectively. The details for 72 individuals can be obtained from Table S1.




Figure 5. Principal coordinate analysis (PCoA) of 72 individuals of three species *Liquidambar* based on 20 EST-SSR markers. The blue squares represent *L. formosana*, the green filled circles represent hybrid sweetgum and the red triangles represent *L. styraciflua*. The first two coordinates explain percentages of total variation for 41.22% and 15.63%, respectively.

4. Discussion

L. formosana is highly adaptable, grows rapidly in forests, exhibits a straight trunk and wide crown, and is economically, ecologically, and medically valuable; thus, it is a genuine “multi-purpose tree” [1,2]. However, molecular genetic data are lacking for *L. formosana* because the molecular tools required are not available. SSRs are efficient genetic markers widely used for MAS, distinguishing among tree varieties, constructing genetic atlases, and researching genetic diversity [21]. The development of SSR markers based on transcriptome sequencing is one of the currently recognized efficient methods [24,39], which provides an effective way to alleviate the shortage of SSR markers available in *Liquidambar*. In previous study, however, only 14 EST-SSR markers [18] were developed for this economic tree species and thus the identification of more co-dominant polymorphic EST-SSR markers with transferable ability from transcriptome would aid genetic improvement of the tree and knowledge transfer to other *Liquidambar* species. Here, we identified 3284 putative SSRs from 2949 unigenes of the transcriptome. The SSR locus frequency was 1/7.3 kb, lower than that of *Camellia sinensis* (1/4.99 kb) [40] but much higher than those of *Pinus dabeshanensis* (1/23.08 kb) [41] and *Larix principis-rupprechtii* (1/26.8 kb) [42]. The predominant repeat type was the dinucleotide (52.56%), as was true for *Robinia pseudoacacia* L. [43], *Populus tomentosa* [44], and *Hevea brasiliensis* [45], but not *Persea americana* Mill [46] and *Dalbergia odorifera* T. Chen. [47]. The numbers and SSR motif types vary among species. Of the dinucleotide repeats, AG/CT units predominated; of the trinucleotide repeats, AAG/CTT was dominant. Differences among studies may be attributable to disparities in genomic structure, the methods used, the parameters selected, and dataset size.

We used 100 primer pairs, selected randomly from a multitude of primer pairs identified via transcriptome analysis, to evaluate the quality and effectiveness of the new EST-SSR markers. Seventy-two primer pairs (72%) generated clear bands from *L. formosana* genomic DNA; the amplification rate was significantly higher than that of the SSR markers of the transcriptomes of *Robinia* (25%) [48] and *Taxodium ‘zhongshansa’* (51.1%) [49]. Eighteen primer pairs did not yield amplicons, despite adjustment for annealing temperature. Sequencing errors may have been a factor, and primers for protein binding sites may have been unable to bind to their templates [50]. Of the 100 primer pairs tested, 32 (32%) amplified EST-SSR markers exhibiting high-level polymorphism; this proportion is much higher than that of *Neolitsea sericea* (9.9%) [51], perhaps reflecting the complexity of genome structure and the level of species diversity [52]. We estimated the genetic diversity parameters of 53 *L. formosana* specimens at 20 loci. The PICs of the 16 new EST-SSR markers ranged from 0.21 to

0.87; 12 markers were highly polymorphic ($PIC > 0.5$). The PICs reflecting moderate polymorphism ($0.2 < PIC < 0.5$) were in the order Liq_eSSR35, Liq_eSSR86, Liq_eSSR67, Liq_eSSR90, Liq_eSSR98, Liq_eSSR25, Liq_eSSR10, Liq_eSSR30, Liq_eSSR33, Liq_eSSR62, Liq_eSSR60, Liq_eSSR59; Liq_eSSR17, Liq_eSSR55, and Liq_eSSR7; Liq_eSSR33 was minimally polymorphic ($PIC < 0.25$) [53]. The average PIC (0.585) was higher than that of a previous study on *L. formosana* [18]. The PIC is affected by all of SSR numbers, the test method used, and SSR motif repeats [54]. As high-level polymorphism is not the only determinant of primer quality, the stability and cross-species transferability of SSR markers were also important resources for genetic analysis. Therefore, we also evaluated the genetic relationships among 72 trees of three *Liquidambar* species; the new EST-SSR markers clearly divided *Liquidambar* into three major clusters. Thus, the markers effectively identified *Liquidambar* species, as confirmed by PCA and genetic structural analysis. The 16 original EST-SSR markers were transferable to *L. styraciflua* and hybrid sweetgum; 15 primer pairs yielded clear bands in the two related species and the transferability rate was 93.75%. This is because EST-SSR markers exhibit high transferability between genetically related species, generally. [55]. Thus, our innovative EST-SSR markers showed high-level polymorphism and can be used in genetic studies of *L. formosana* and related species for interspecific hybrids, comparative mapping analysis, and phylogenetic relationships among species.

Evaluation of genetic diversity is a core issue in forestry. Knowledge of such diversity allows us to better understand adaptation to environmental changes and the evolution of specific species, thus facilitating species protection. Genetic diversity is reflected both phenotypically and at the molecular level. Analysis of such diversity using molecular markers provides insight into the genetic background [56]. We evaluated the genetic diversity of a plus *L. formosana* population of Henan Province using 20 polymorphic EST-SSR markers. The mean H_o was 0.653, the mean H_e was 0.621, and the mean PIC was 0.578. Compared to the diversity of a natural *L. formosana* population [18], we found, surprisingly, that the plus population exhibited greater genetic diversity even after selection. The potential reason may be due to different primers or the influence of selection. The study of superior trees allows us to understand production potential, determine selection criteria, and plan breeding strategies [14]. Heterosis was clearly a factor during interspecific hybridization of *Liquidambar* species [57]. Therefore, the plus *L. formosana* population exhibiting abundant genetic variation and a superior phenotype is an ideal hybridization parent, thereby providing valuable materials for further genetic improvement of *Liquidambar* species.

Cluster analysis and principal component analysis (PCA) clearly divided the three species of *Liquidambar* into three categories according to the species classification standard, although previous studies showed that the genetic relationship between *L. formosana* and *L. styraciflua* was relatively close. The results also showed the high efficiency of our original primer. The results of genetic structure analysis showed that the three *Liquidambar* species were divided into two main gene pools, the first gene pool (red) was mainly composed of *L. formosana*, the second gene pool (green) was mainly composed of *L. styraciflua*, while the hybrid sweetgum contained two gene pools, which indicated that the genetic composition of hybrid sweetgum was affected by both male and female parent, and hybrid sweetgum had a closer relationship with *L. styraciflua*. [36].

5. Conclusions

We used transcriptome data to develop novel EST-SSR markers exhibiting high-level cross-species transferability. These markers will serve as powerful molecular tools for evaluation of genetic diversity, MAS, and gene map construction, and for use in genetic studies. The genetic diversity of the plus *L. formosana* population will guide the creation of second-generation plus *L. formosana* trees, and improve *L. formosana* breeding programs.

Supplementary Materials: The following are available online at <http://www.mdpi.com/1999-4907/11/2/203/s1>, Table S1: Details for the 72 individuals of *Liquidambar*.

Author Contributions: Experiment designed, S.C., J.Z. (Jinfeng Zhang) and M.D.; experiment performed, S.C. and M.D.; data analyzed, S.C., M.D. and Y.Z.; materials collected, Y.Z., S.Q. and X.L.; writing—original draft

preparation, S.C.; writing—review and editing, J.Z. (Jian Zhao), Y.Z., M.D. All authors have read and agreed to the published version of the manuscript.

Funding: This paper was supported by Medium and Long Scientific Research Project for Young Teachers in Beijing Forestry University (2015ZCQ-SW-02), “948” Project of China (2014-4-59), and major science and technology special project of Xuchang, Henan province, China (20170112006).

Conflicts of Interest: The authors declare that they have no conflicts of interest.

References

1. Zhou, G.Y. Chinese sweetgum. *J. Ningbo Univ.* **1995**, *2*, 34–41.
2. Zheng, Y.Q.; Pan, B.; Itohl, T. Chemical induction of traumatic gum ducts in Chinese sweetgum, *Liquidambar formosana*. *IAWA J.* **2015**, *36*, 58–68. [[CrossRef](#)]
3. Merkle, S.A.; Neu, K.A.; Battle, P.J.; Bailey, R.L. Somatic embryogenesis and plantlet regeneration from immature and mature tissues of sweetgum (*Liquidambar styraciflua*). *Plant Sci.* **1998**, *132*, 169–178. [[CrossRef](#)]
4. Kathleen, B.P.; Ickert-Bond, S.M.; Wen, J. Anatomically preserved *Liquidambar* (Altingiaceae) from the middle Miocene of Yakima Canyon, Washington state, USA, and its biogeographic implications. *Am. J. Bot.* **2004**, *91*, 499–509.
5. Jiang, P.C. A survey of studies on *Liquidambar formosana*. *Guangxi Med. J.* **1998**, *20*, 241–242.
6. Zeng, T.; Song, X. Study on the Methods of the Tapping from Chinese SweetGum. *China For. Sci. Technol.* **2010**, *24*, 84–87.
7. Weng, L.L.; Jiang, J.D. The research progresses and prospects of local species *Liquidamba formosana*. *J. Fujian For. Sci. Technol.* **2007**, *34*, 184–189.
8. Winstead, J.E. Populational Differences in Seed Germination and Stratification Requirements of Sweetgum. *For. Sci.* **1971**, *17*, 34–36.
9. Suhaendi, H. Seed germination and seedling characteristics of twelve greenhouse grown provenances of *L. Styraciflua*. *Buletin Penelitian Hutan* **1989**, *520*, 1–18.
10. Kariuki, J.G. Interim results of a provenancetrial of liquidambar styraciflua in Kenya. *Kenya For. Res. Inst.* **1989**, *5*, 1–17.
11. He, G.P.; Chen, Y.T.; Tang, X.Y.; Shen, G.P.; Gao, S.M. A Study on Variations of Young Forest Growing Traits in *Liquidambar formosana* Geographic Provenances. *Acta Agric. Univ. Jiangxiensis* **2005**, *27*, 585–589.
12. Shi, J.L.; Zhu, S.L.; Ma, K.X.; Zhu, J.P. Study on multi-objective breeding of *Liquidambar formosana*. *For. Sci. Technol.* **1997**, *10*, 13–14.
13. Zhao, Y.; Wang, X.R.; Tang, R.H. The elite tree selection of *Liquidambar formosana* in Guizhou Province. *J. Mt. Agric. Biol.* **2009**, *28*, 530–534.
14. Cornelius, J. The effectiveness of plus-tree selection for yield. *For. Ecol. Manag.* **1994**, *67*, 23–34. [[CrossRef](#)]
15. Xu, Y.B.; Crouch, J.H. Marker-assisted selection in plant breeding: From publications to practice. *Crop. Sci.* **2008**, *48*, 391–407. [[CrossRef](#)]
16. Vendrame, W.A.; Holliday, C.P.; Merkle, S.A. Clonal propagation of hybrid sweetgum (*Liquidambar styraciflua* × *L. formosana*) by somatic embryogenesis. *Plant Cell Rep.* **2001**, *20*, 691–695. [[CrossRef](#)]
17. Bi, X.Q.; Jin, X.Z.; Li, H.J. Genetic diversity in the natural populations of *Liquidambar formosana* revealed by ISSR molecular markers. *Bull. Bot. Res.* **2010**, *30*, 120–125.
18. Sun, R.; Lin, F.; Huang, P.; Zheng, Y. Moderate Genetic Diversity and Genetic Differentiation in the Relict Tree *Liquidambar formosana* Hance Revealed by Genic Simple Sequence Repeat Markers. *Front. Plant Sci.* **2016**, *7*, 1411. [[CrossRef](#)]
19. Yin-Gang, L.I.; Zhou, X.P.; Liu, X.H.; Hua, Z.H.; Liu, W.; Liu, J.L.; Xie, J.Q. Preliminary Study on Growth of *Styrax tonkinensis* Plantation and Plus Tree Seletion. *J. Zhejiang For. Sci. Technol.* **2010**, *30*, 24–28.
20. Kuang, H.; Richardson, T.; Carson, S.; Wilcox, P.; Bongarten, B. Genetic analysis of inbreeding depression in plus tree 850.55 of *Pinus radiata* D. Don. I. Genetic map with distorted markers. *Theor. Appl. Genet.* **1999**, *98*, 697–703. [[CrossRef](#)]
21. Li, D.; Deng, Z.; Qin, B.; Liu, X.; Men, Z. De novo assembly and characterization of bark transcriptome using Illumina sequencing and development of EST-SSR markers in rubber tree (*Hevea brasiliensis* Muell. Arg.). *BMC Genom.* **2012**, *13*, 192. [[CrossRef](#)] [[PubMed](#)]

22. Colburn, B.C.; Mehlenbacher, S.A.; Sathuvalli, V.R. Development and mapping of microsatellite markers from transcriptome sequences of European hazelnut (*Corylus avellana* L.) and use for germplasm characterization. *Mol. Breed.* **2017**, *37*, 16. [CrossRef]
23. Aggarwal, R.K.; Hendre, P.S.; Varshney, R.K.; Bhat, P.R.; Krishnakumar, V.; Singh, L. Identification, characterization and utilization of EST-derived genic microsatellite markers for genome analyses of coffee and related species. *Theor. Appl. Genet.* **2007**, *114*, 359–372. [CrossRef] [PubMed]
24. Zalapa, J.E.; Cuevas, H.; Zhu, H.; Steffan, S.; Senalik, D.; Zeldin, E.; McCown, B.; Harbut, R.; Simon, P. Using next-generation sequencing approaches to isolate simple sequence repeat (SSR) loci in the plant sciences. *Am. J. Bot.* **2012**, *99*, 193–208. [CrossRef]
25. Wei, W.; Wang, C.B.; Fang, L.; Chen, H.D.; Lin, W.; Wang, C.Y.; Zhang, X.D.; Wang, Y.H.; Kun-Bo, A.W. Development and Evaluation of New Non-Redundant EST-SSR Markers from *Gossypium*. *Acta Agron. Sin.* **2012**, *38*, 1443–1451.
26. Xu, S.C.; Gong, Y.-M.; Mao, W.-H.; Hu, Q.-Z.; Zhang, G.-W.; Fu, W.; Xian, Q.-Q. Development and characterization of 41 novel EST-SSR markers for *Pisum sativum* (Leguminosae). *Am. J. Bot.* **2012**, *99*, e149–e153. [CrossRef]
27. Yadong, Z.; Hu, X.; Zhang, Y.D.; Song, C.; Hu, Y.X.; Song, C.W. Identification of Populus Varieties from Hubei Province by EST-SSR Marker. *Mol. Plant. Breed.* **2009**, *7*, 105–109.
28. Zhang, Y.; Wang, Z.W.; Qi, S.Z.; Wang, X.Q.; Zhao, J.; Zhang, J.F.; Li, B.L.; Zhang, Y.D.; Liu, X.Z.; Yuan, W. In vitro tetraploid induction from leaf and petiole explants of hybrid sweetgum (*Liquidambar styraciflua* × *Liquidambar formosana*). *Forests* **2017**, *8*, 264. [CrossRef]
29. Grabherr, M.G.; Haas, B.J.; Yassour, M.; Levin, J.Z.; Thompson, D.A.; Amit, I.; Adiconis, X.; Fan, L.; Raychowdhury, R.; Zeng, Q. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat. Biotechnol.* **2011**, *29*, 644–652. Available online: <https://www.ncbi.nlm.nih.gov/> (accessed on 27 April 2017). [CrossRef]
30. Rozen, S.; Skaletsky, H. Primer3 on the WWW for general users and for biologist programmers. *Methods Mol. Biol.* **2000**, *132*, 365–386.
31. Doyle, J.J.; Doyle, J.L. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem. Bull.* **1987**, *19*, 11–15.
32. Schuelke, M. An economic method for the fluorescent labeling of PCR fragments. *Nat. Biotechnol.* **2000**, *18*, 233–234. [CrossRef] [PubMed]
33. Jeffrey, C. Glaubitz convert: A user-friendly program to reformat diploid genotypic data for commonly used population genetic software packages. *Mol. Ecol. Resour.* **2004**, *4*, 309–310.
34. Popgene. Version 1.32. The User-Friendly Shareware for Population Genetic Analysis molecular Biology and Biotechnology Center, University of Alberta Edmonton. Available online: [http://www.ualberta.ca/~\[protect\]T1/textbraceleftprotectT1textbracerightfyeh](http://www.ualberta.ca/~[protect]T1/textbraceleftprotectT1textbracerightfyeh) (accessed on 23 November 2017).
35. Nagy, S.; Pocza, P.; Cernák, I.; Taller, J. PICcalc: An Online Program to Calculate Polymorphic Information Content for Molecular Genetic Studies. *Biochem. Genet.* **2012**, *50*, 670–672. [CrossRef] [PubMed]
36. Evanno, G.; Regnaut, S.; Goudet, J. Detecting the number of clusters of individuals using the software structure: A simulation study. *Mol. Ecol.* **2005**, *14*, 2611–2620. [CrossRef] [PubMed]
37. Liu, K.; Muse, S.V. PowerMarker: an integrated analysis environment for genetic marker analysis. *Bioinformatics* **2005**, *21*, 2128–2129. [CrossRef]
38. Peakall, R.; Smouse, P.E. genalex 6: genetic analysis in Excel. Population genetic software for teaching and research. *Mol. Ecol. Notes* **2005**, *6*, 288–295. [CrossRef]
39. Xu, W.; Yang, Q.; Huai, H.; Liu, A. Development of EST-SSR markers and investigation of genetic relatedness in tung tree. *Tree Genet. Genomes* **2012**, *8*, 933–940. [CrossRef]
40. Wu, H.; Chen, D.; Li, J.; Yu, B.; Qiao, X. De Novo Characterization of Leaf Transcriptome Using 454 Sequencing and Development of EST-SSR Markers in Tea (*Camellia sinensis*). *Plant. Mol. Biol. Rep.* **2013**, *31*, 524–538. [CrossRef]
41. Xiang, X.; Zhang, Z.; Wang, Z.; Zhang, X.; Wu, G. Transcriptome sequencing and development of EST-SSR markers in *Pinus dabeshanensis*, an endangered conifer endemic to China. *Mol. Breed.* **2015**, *35*, 158. [CrossRef]

42. Dong, M.; Wang, Z.; He, Q.; Zhao, J.; Fan, Z.; Zhang, J. Development of EST-SSR markers in *Larix principis-rupprechtii* Mayr and evaluation of their polymorphism and cross-species amplification. *Trees* **2018**, *32*, 1559–1571. [\[CrossRef\]](#)
43. Dong, L.; Sun, Y.H.; Zhao, J.; Zhang, Y.W.; Li, X.Y. Development and Application of EST-SSR Markers for DNA Fingerprinting and Genetic Diversity Analysis of the Main Cultivars of Black Locust (*Robinia pseudoacacia* L.) in China. *Forests* **2019**, *10*, 644. [\[CrossRef\]](#)
44. Du, Q.; Wang, B.; Wei, Z.; Zhang, D.; Li, B. Genetic Diversity and Population Structure of Chinese White Poplar (*Populus tomentosa*) Revealed by SSR Markers. *J. Hered.* **2012**, *103*, 853–862. [\[CrossRef\]](#) [\[PubMed\]](#)
45. Nirapathongporn, K.; Kongsawadworakul, P.; Viboonjun, U.; Teerawattanasuk, K.; Chrestin, H.; Segiun, M.; Clément-Dement, A.; Narangajavana, J. Development and mapping of functional expressed sequence tag-derived simple sequence repeat markers in a rubber tree RRIM600 × PB217 population. *Mol. Breed.* **2016**, *36*, 39. [\[CrossRef\]](#)
46. Ge, Y.; Tan, L.; Wu, B.; Wang, T. Transcriptome Sequencing of Different Avocado Ecotypes: de novo Transcriptome Assembly, Annotation, Identification and Validation of EST-SSR Markers. *Forests* **2019**, *10*, 411. [\[CrossRef\]](#)
47. Liu, F.M.; Hong, Z.; Yang, Z.J.; Zhang, N.N. De Novo Transcriptome Analysis of *Dalbergia odorifera* T. Chen (Fabaceae) and Transferability of SSR Markers Developed from the Transcriptome. *Forests* **2019**, *10*, 99.
48. Guo, Q.; Wang, J.-X.; Su, L.-Z.; Lv, W.; Sun, Y.-H.; Li, Y. Development and evaluation of a novel set of EST-SSR markers based on transcriptome sequences of Black Locust (*Robinia pseudoacacia* L.). *Genes* **2017**, *8*, 177. [\[CrossRef\]](#)
49. Cheng, Y.L.; Yang, Y.; Wang, Z.Y.; Qi, B.Y.; Yin, Y.L.; Li, H.G. Development and Characterization of EST-SSR Markers in *Taxodium 'zhongshansa'*. *Plant. Mol. Biol. Rep.* **2015**, *33*, 1–11. [\[CrossRef\]](#)
50. Jia, X.; Deng, Y.; Sun, X.; Liang, L.; Su, J. De novo assembly of the transcriptome of *Neottopteris nidus* using Illumina paired-end sequencing and development of EST-SSR markers. *Mol. Breed.* **2016**, *36*, 94. [\[CrossRef\]](#)
51. Chen, L.Y.; Cao, Y.N.; Yuan, N.; Koh, N.; Wang, G.M.; Qiu, Y.X. Characterization of transcriptome and development of novel EST-SSR makers based on next-generation sequencing technology in *Neolitsea sericea* (Lauraceae) endemic to East Asian land-bridge islands. *Mol. Breed.* **2015**, *35*, 187. [\[CrossRef\]](#)
52. Postolache, D.; Leonarduzzi, C.; Piotti, A.; Spanu, I.; Vendramin, G.G. Transcriptome versus Genomic Microsatellite Markers: Highly Informative Multiplexes for Genotyping *Abies alba* Mill. and Congeneric Species. *Plant. Mol. Biol. Rep.* **2013**, *32*. [\[CrossRef\]](#)
53. Botstein, D.; White, R.L.; Skolnick, M.; Davis, R.W. Construction of a genetic linkage map in man using restriction fragment length polymorphisms. *Am. J. Hum. Genet.* **1980**, *32*, 314–331. [\[PubMed\]](#)
54. Liu, S.; Liu, H.; Wu, A.; Hou, Y.; An, Y.; Wei, C. Construction of fingerprinting for tea plant (*Camellia sinensis*) accessions using new genomic SSR markers. *Mol. Breed.* **2017**, *37*, 93. [\[CrossRef\]](#)
55. Durand, J.; Bodénès, C.; Chancerel, E.; Frigerio, J.-M.; Vendramin, G.; Sebastiani, F.; Buonamici, A.; Gailing, O.; Koelewijn, H.-P.; Villani, F.; et al. A fast and cost-effective approach to develop and map EST-SSR markers: oak as a case study. *BMC Genom.* **2010**, *11*, 570. [\[CrossRef\]](#) [\[PubMed\]](#)
56. Wen, M.; Wang, H.; Xia, Z.; Zou, M.; Lu, C.; Wang, W. Development of EST-SSR and genomic-SSR markers to assess genetic diversity in *Jatropha Curcas* L. *BMC Res. Notes* **2010**, *3*, 42. [\[CrossRef\]](#) [\[PubMed\]](#)
57. Merkle, S.; Montello, P.; Kormanik, T.; Le, H. Propagation of novel hybrid sweetgum phenotypes for ornamental use via somatic embryogenesis. *Propag. Ornam. Plant.* **2010**, *10*, 220–226.

