

Table S4. Results of Chi-square tests for the comparison of host-feeding groups between the six most abundant mosquito species with adjusted p-values collected at four sampling sites in the DDBR, Romania (2014-2016).

Mosquito species	Mosquito species	adjusted p value	χ^2
<i>Aedes caspius</i>	<i>Aedes vexans</i>	0.036648349	12.0288742
<i>Aedes caspius</i>	<i>Anopheles hyrcanus</i>	1	0.876218151
<i>Aedes caspius</i>	<i>Anopheles maculipennis</i> s.l.	1	1.050654796
<i>Aedes caspius</i>	<i>Coquillettidia richiardii</i>	0.5154945	6.741357691
<i>Aedes caspius</i>	<i>Culex pipiens</i> s.l./torrentium	1.88E-07	36.3867405
<i>Aedes vexans</i>	<i>Anopheles hyrcanus</i>	9.19E-06	28.61119323
<i>Aedes vexans</i>	<i>Anopheles maculipennis</i> s.l.	0.00018112	22.64880073
<i>Aedes vexans</i>	<i>Coquillettidia richiardii</i>	0.227764729	8.374984545
<i>Aedes vexans</i>	<i>Culex pipiens</i> s.l./torrentium	1.80E-18	87.13728235
<i>Anopheles hyrcanus</i>	<i>Anopheles maculipennis</i> s.l.	1	3.276589246
<i>Anopheles hyrcanus</i>	<i>Coquillettidia richiardii</i>	8.18E-09	42.65875941
<i>Anopheles hyrcanus</i>	<i>Culex pipiens</i> s.l./torrentium	2.16E-31	146.6318183
<i>Anopheles maculipennis</i> s.l.	<i>Coquillettidia richiardii</i>	0.006237562	15.57043205
<i>Anopheles maculipennis</i> s.l.	<i>Culex pipiens</i> s.l./torrentium	2.48E-12	58.85872968
<i>Coquillettidia richiardii</i>	<i>Culex pipiens</i> s.l./torrentium	3.07E-23	109.094442