

Communication

Tomato Twisted Leaf Virus: A Novel Indigenous New World Monopartite Begomovirus Infecting Tomato in Venezuela

Gustavo Romay ^{1,*}, Francis Geraud-Pouey ², Dorys T. Chirinos ³, Mathieu Mahillon ¹, Annika Gillis ^{4,†}, Jacques Mahillon ⁴ and Claude Bragard ^{1,*}

¹ UCLouvain, Earth and Life Institute, Applied Microbiology-Phytopathology, Croix du Sud 2-L07.05.03, 1348 Louvain-la-Neuve, Belgium; mathieu.mahillon@uclouvain.be

² La Universidad del Zulia (LUZ), Unidad Técnica Fitosanitaria, Maracaibo 4005, Estado Zulia, Venezuela; fgeraudp@gmail.com

³ Facultad de Ingeniería Agronómica, Universidad Técnica de Manabí, Manabí 130105, Ecuador; dtchirinos@gmail.com

⁴ UCLouvain, Earth and Life Institute, Applied Microbiology-Laboratory of Food and Environmental Microbiology, Croix du Sud 2-L7.05.12, 1348 Louvain-la-Neuve, Belgium; annika.gillis@uclouvain.be (A.G.); jacques.mahillon@uclouvain.be (J.M.)

[†] Present address: Section of Microbiology and Medical Research Council Centre for Molecular Bacteriology and Infection, Imperial College London, London SW72AZ, United Kingdom.

* Correspondence: gustavo.romay@uclouvain.be (G.R.); claude.bragard@uclouvain.be (C.B.)




Figure S1. (A) *Nicotiana benthamiana* plant with mosaic symptoms at four weeks after agroinoculation with the ToTLV infectious clone. (B) *N. benthamiana* plant with no symptoms at four weeks after agroinoculation with the pCAMBIA1300 empty vector as a negative control