

SUPPLEMENT DATA

SUPPLEMENTARY TABLE

Table S1: Comparison of the differences between mean \pm standard deviation of log₁₀ titers at each time point

Viruses	Mean \pm SD of log ₁₀ titers at different time points (hours post inoculation)						Overall
	4	8	12	16	20	24	
O189	0.8 ^a \pm 1.4	3.3 ^a \pm 0.3	3.7 ^a \pm 0.3	3.6 ^a \pm 0.1	4.0 ^b \pm 0.3	5.4 ^a \pm 0.1	3.5 ^a \pm 1.5
Asia I	1.7 ^a \pm 1.4	2.7 ^b \pm 0.0	3.3 ^a \pm 0.3	3.5 ^a \pm 0.5	4.1 ^{ab} \pm 0.4	4.0 ^b \pm 0.4	3.2 ^a \pm 1.0
A/lopburi	0.8 ^a \pm 1.4	3.4 ^a \pm 0.1	2.6 ^{ab} \pm 2.2	3.4 ^a \pm 0.1	4.6 ^a \pm 0.1	3.9 ^b \pm 0.4	3.1 ^{ab} \pm 1.5
O/41	2.5 ^a \pm 0.0	2.5 ^b \pm 0.0	2.5 ^{ab} \pm 0.0	2.5 ^b \pm 0.0	2.5 ^c \pm 0.0	2.5 ^c \pm 0.0	2.5 ^b \pm 0.0
Non-virus control	0.0 ^a \pm 0.0	0.0 ^c \pm 0.0	0.0 ^{ab} \pm 0.0	0.0 ^c \pm 0.0	0.0 ^d \pm 0.0	0.0 ^d \pm 0.0	0.0 ^c \pm 0.0

Note: The different letters over the mean titers at each time point indicate statistically significant differences.