

Table S3. Diagnostic performance of the clinical scoring system in differentiating severe fever with thrombocytopenia syndrome.

Clinical Score* (n=247)	SFTS (n=112)	Scrub Typhus (n=135)	Sensitivity (95% CI)	Specificity (95% CI)	Positive Likelihood Ratio (95% CI)	Negative Likelihood Ratio (95% CI)	Positive Predictive Value	Negative Predictive Value
>0	112	37	100.0 (96.8– 100.0)	72.6 (64.3–79.9)	3.7 (2.8–4.8)	0	75.2 (69.7–79.9)	100.0
>1	109	3	97.3 (92.4–99.4)	97.8 (93.6–99.5)	43.8 (14.3–134.1)	0.03 (0.009–0.08)	97.3 (92.2–99.1)	97.8 (93.5–99.3)
>2	67	0	59.8 (50.1–69.0)	100.0 (97.3–100.0)	NA	0.40 (0.3–0.5)	100.0	75.0 (70.5–79.0)
>3	24	0	21.4 (14.2–30.2)	100.0 (97.3–100.0)	NA	0.79 (0.7–0.9)	100.0	60.5 (58.2–62.8)

* Scoring system = (1 x leukopenia) + (1 x prolonged aPTT) + (1 x normal CRP level \leq 3.0 mg/dL) + (1 x elevated CK level). Abbreviations: CI, confidence interval; SFTS, severe fever with thrombocytopenia syndrome; aPTT, activated partial thromboplastin time; CRP, C-reactive protein; CK, creatine kinas